

INVITED REVIEW

HUANGLONGBING: A DESTRUCTIVE, NEWLY-EMERGING,
CENTURY-OLD DISEASE OF CITRUS¹

J.M. Bové

Laboratoire de Biologie Cellulaire et Moléculaire, Institut de Biologie Végétale Montpellier (IBVM), Unité Mixte de Recherche en Génétique, Développement et Pouvoir Pathogène, Centre de Recherche INRA de Bordeaux, 71, Ave. Edouard Bourlès, 33883 Villenave d'Ornon Cedex, France, and Fundecitrus, 201, Av. Dr. Albeniz Pereira de Barros, Aracajuano S.P., Brazil

SUMMARY

A detailed account is given of the history, aetiology, biology, epidemiology, detection, geographical distribution, and control of huanglongbing (HLB), a destructive disease of citrus that represents a major threat to the world citrus industry, and is slowly invading new citrus-growing areas. HLB, whose name in Chinese means 'yellow dragon disease', was first reported from southern China in 1919 and is now known to occur or next to 40 different Asian, African, Oceanian, South and North American countries. The agent is a phloem-restricted, non cultured, Gram-negative bacterium causing crippling diseases denoted "greening" in South Africa, "mottle leaf" in the Philippines, "dieback" in India, "vein phloem degeneration" in Indonesia. The HLB bacterium belongs to the genus *Candidatus Liberibacter*, three species of which are currently known, *Candidatus Liberibacter asiaticus*, occurring in Asian countries and, to a lesser extent, in Brazil and the USA (Florida), *Candidatus Liberibacter africanus* with its subspecies "casensis", recorded from African countries, and *Candidatus Liberibacter americanus* present in Brazil. The suggestion is that each liberibacter species has evolved in the continent after which it is named. HLB symptoms are virtually the same wherever the disease occurs. Infected trees show a bumpy mottle condition of the leaves that results in the development of yellow shoots, the early and very characteristic symptom of the disease. Trees are stunted, declining and bear a few, small sized, and deformed (lop-sided) fruits, that are poorly coloured (greening) and with coloration starting at the peduncular end (colour inversion). HLB can be transmitted by grafting from citrus to citrus and by dipterid to psyllid-like. The psyllids *Trialeurodes erythrae* and *Diuraphis citri* are natural vectors. Two different types of

HLB are known: the heat sensitive African form transmitted by *T. erythrae*, which develops at temperatures of 22–25°C, and the heat tolerant Asian form, transmitted by *D. citri*, which stands temperatures well above 30°C. Although the HLB pathogen can be identified by electron microscopy, other laboratory methods are used for routine detection. ELISA with monoclonal antibodies is not recommended. Better systems are dot blot hybridization with a DNA probe, and various PCR formats (one step, nested, multiplex) using species specific primers based on 16S rRNA or *rflKA/JL-opsBC* operon sequences. Because no curative methods of HLB are available, control is preventive and largely based on inoculum elimination by removal of infected trees and chemical treatments against vectors. Strict quarantine measures must be implemented to impair further international spread of HLB agents and their vectors.

Key words: Citrus, huanglongbing, liberibacter, psyllid vectors, diagnosis, control

INTRODUCTION

Huanglongbing (HLB), a destructive disease of citrus, is caused by endogenous, slow-acting restricted bacteria, named liberibacters, which are transmitted from tree to tree by citrus psyllid insect vectors: *Diuraphis citri* in Asia and America, and *Trialeurodes erythrae* in Africa. Practically all commercial citrus species and cultivars are sensitive, regardless of rootstocks. Another destructive affliction of citrus is tristeza disease, due to *Citrus tristeza virus* (CTV), which induces quick decline and death of citrus trees grafted on sour orange (*Citrus aurantium* L.) rootstock, Carrizo rootstock. Control of tristeza disease is achieved by replacing sour orange by rootstocks giving tolerant combinations with the scions. Many important citrus regions, where CTV is endemic, have learned to live with it. For HLB however, no control is known, except preventing the trees from becoming infected. Therefore, HLB is probably the most serious disease of citrus, much more serious than tristeza, and it represents a dangerous threat for regions still free of the disease, such as the Mediterranean basin, Western Asia,

Corresponding author: J.M. Bové
Fax: +33 557 122369
E-mail: joseph.bove@wanadoo.fr

¹ This review is dedicated to the memory of Professor S.R. Capoor (1912–1993), one of the most eminent phytopathologists of India, who carried out most of his work on virus and virus-like diseases of citrus and other crops at the Virus Research Centre in Poona, Maharashtra, India.

Australia and Pacific Ocean islands.

Until recently, America was also free of HLB, but in March 2004 and August 2005, symptoms of the disease were recognized, respectively in the State of São Paulo, Brazil, and in Florida, USA, two of the largest citrus growing regions in the world. This discovery has given a renewed interest in HLB, as witnessed by the organisation of an international workshop on HLB in November 2005 in Florida.

Even though HLB is newly emerging in America, it is probably one of the oldest known diseases of citrus. Reviews on HLB in Asia and Africa have appeared (Garnier and Bové, 1993; da Graça, 1991; da Graça and Kester, 2004; Halbert and Manjunath, 2004). In this review, America will be covered as well. Until 1995, the disease was generally known under the South African name "greening". Today, the official designation is "huanglongbing", the most appropriate Chinese name (see below). Here, the abbreviation "HLB" will be used throughout as the generic name of the disease, regardless of the country where it occurs.

The mycoplasmal agent of citrus stubborn disease (Grewing and Calavan, 1970; Ladrière and Bové, 1970b) and the bacterial agent of HLB (Lafôche and Bové, 1970a) were discovered by electron microscopy (EM) in the same year, 1970. The stubborn agent was available in culture by 1971, characterized as *Spiroplasma citri* by 1973 (Saglio *et al.*, 1973), and its genome sequence completed by 2005. Even today, the HLB bacterium has not yet been obtained in culture. This explains why characterisation of the HLB agent has not progressed as fast as that of *S. citri*. Molecular techniques had to become available to finally characterise the organism at the phylogenetic and taxonomic level, and this review will focus in particular on the nature of the HLB bacteria and their detection for HLB identification and confirmation. As the phytoplasmata and the HLB liberibacters have not yet been cultured, Koch's postulates could not be fulfilled. However, from overwhelming but indirect evidence, it is assumed that they are the causal agents of the diseases with which they are associated. This assumption will be followed here.

HISTORICAL BACKGROUND

Huanglongbing in China. Reinking (1919), evaluating diseases of economic plants in southern China, used English to report, in 1919, on "yellow shoot" of citrus, a disease which he thought to be of little importance in those days. However, later surveys showed that by 1936 the disease had spread to become a serious problem. The most extensive work on HLB in southern China was to be conducted from 1941 to 1955 by Lin Kung Hsiang (Fig. 1A). Born in 1910 in Ganxia, within the Fujian citrus belt, he obtained his Ph.D. at Cornell Uni-

versity, USA, in the late 1930s (see Lin Kung Shun' *et al.*, 1996). On his return to China, he joined the Christian University of Langnan in Guangzhou, later to become the South China Agricultural University, and devoted most of his time to the study of HLB. Between 1941 and 1955, he carried out several surveys and field visits (Fig. 2). For instance, in 1943 he visited Taiwan, which he found affected by HLB, or rather "likobin", the name for HLB in the island. From the discussions he had with the farmers of the Chaoshou county in Guangdong province (see Fig. 2), he learned that HLB had been there since the 1870s, and on the basis of various observations he estimated that HLB in South China originated from that area.

In the Chaoshou district, the name given by the farmers to the disease was "huang long bing", "bing" standing for disease, "huang" meaning yellow, and "long", dragon, hence: yellow dragon disease. However, in other districts, different names were used. Lin considered that "huang long" was the most appropriate name, because "huang long", i. e. yellow dragon or yellow shoot, is the name given by the farmers to the new flush of growth on infected trees, and represents a characteristic, early symptom of the disease. Fig. 6 shows such yellow shoots or dragons. The most significant result obtained by Lin was the demonstration, by precise experimental work, that HLB is a graft-transmissible, infectious disease, and should neither be attributed to physiological disorders such as mineral deficiencies or water logging, nor to soil-borne diseases such as nematode infestation or *Fusarium* infection (Lin, 1956). In Lin's experiments, transmission of the HLB pathogen was rightly obtained by graft-inoculation. Previously, Chen Chi-yan (1943) had also obtained "transmission", but only by graft-propagation of HLB-affected shoots. As pointed out by Lin (1956), Chen failed to provide evidence for the systemic, infectious nature of HLB.

Lin's work was carried out during difficult times in China. The results were published in 1956 (Lin, 1956). It is often said that Lin's results remained practically unknown to the scientific community out of China. This is probably so, but there was at least one "Western" phytopathologist, Prof. Antonio Cecarone, who not only knew the work of Lin, for having visited the Guangzhou area in 1956, but who also tried to make it known by publishing a note on "yellow shoot" in the Italian journal *Rivista di Agronomia* 2: 45-50, 1957, in which he gave the reference to Lin's 1956 publication. In his note, Antonio Cecarone pointed out rightly that in Lin's transmission experiments, which were all done in the open, some of the uninoculated control plants showed HLB symptoms. Lin was well aware of this situation, and this prompted him to undertake, early 1951, a second trans-



Fig. 1. Pioneering studies on HLB were conducted by: (A) Lin Kung Hsiang (1910-1986) in China (Ph.D. from Cornell University, USA); (B) A.P.D. McClean (1900-1999) in South Africa (Ph.D. from Natal University, South Africa); (C) S.P. Capoor (1912-1995) in India (Ph.D. from London University, UK); and (D) Monique Garnier (1949-2001) in France (Ph.D. from Bordeaux 2 University, France).



Fig. 2. Places visited by Lin Kung Hsiang between 1941 and 1955 in southern China and Taiwan. Formal offer of HLB-191 with no clear evidence for HLB-101.

mission experiment involving a Taikan mandarin block and a Ponkan mandarin block. At the end of 1954, in the Ponkan block for instance, none of the 112 uninoculated control trees and none of the 50 unpruned guard row trees were symptomatic, while as many as 83 of 94 inoculated trees showed HLB symptoms, and only 4 of 22 unpruned guard row trees were symptomatic. These data clearly demonstrated transmission of the HLB pathogen by graft inoculation. In addition, in order to explain the fact that some uninoculated trees became affected, Lin assumed vector transmission to occur.

It has become evident that Lin was the first to demonstrate, by graft inoculation, the infectious nature of the disease, for which he appropriately used the name "huanglongbing". For these reasons, the International Organization of Citrus Virologists (IOCV) proposed in 1995 at the 13th conference of the Organization in Fuzhou (Fujian, China), that the official name of the disease be huanglongbing (HLB), and this proposal was accepted. Today, HLB is widely used for the African, American, and Asian forms of the disease.

Greening in South Africa. A disease similar to HLB was observed in 1928 under the name "yellow shoot" in the western Transvaal, while the name "greening" prevailed in the eastern Transvaal (Oberholzer *et al.*, 1965). However, the true nature of the disease was not immediately recognized, and, in the first description of "greening" in 1937, the problem was still assumed to be mineral toxicity (van der Merwe and Andersen, 1957). It was the collaboration between a phytopathologist, A.P.D. McClean (Fig. 1B), and a horticulturist, P.C.I. Oberholzer, that resulted in 1965 in the demonstration that greening was transmissible by graft inoculation (McClean and Oberholzer, 1965a) as well as by the African citrus psylla, *T. erythrae* (McClean and Oberholzer, 1965b). The infectious nature of "greening" was thus established, however ten years after that of "huanglongbing".

Mottle leaf in the Philippines. The disease was described in 1921, and thought to be related to zinc deficiency (Lan, 1921). It became a serious problem in the late 1950s. In 1966, Saliba and Cortez (1968) and Martinez and Wallace (1968) stressed the similarities between the symptoms of "mottle leaf" and those of HLB in China and Taiwan, and "greening" in South Africa. The major contribution of the Philippines to HLB was the demonstration, in 1967, that "mottle leaf" could be transmitted by the Asian citrus psylla, *D. citri* (Martinez and Wallace, 1967). In the same year, it was also reported in India that HLB could be transmitted by *D. citri* from trees affected by "citrus dieback".

Dieback and HLB in India. India is apparently another country where HLB seems to have a long history. As indicated by Kapoor (1963), citrus in India has been

known to suffer seriously from certain disorders resulting in low production, twig dieback, slow death and even sudden wilting. These symptoms had been attributed to "dieback", a disease that was first observed by Roghaji Bhonsale (cited by Kapoor, 1963) in the 18th Century in the Central Provinces, soon after the introduction of citrus in India. It was also observed by Bonavia (1883) in Assam. However, the problem with Indian "dieback" comes from the fact that, as pointed out by Asana (1955), "dieback in citrus is not a specific disease". Similarly, for Kapoor (1963), dieback was merely a symptom picture, and many factors, including soil disorders, nutritional deficiencies, twig fungi, and viruses, were evoked to account for it. In some cases, the major cause was CTV. Indeed, in the State of Bombay it was clearly proven by the Virus Research Center at Poona that some type of dieback was due to CTV (Masodeva and Kapoor, 1958; Kapoor, 1964). In other cases, HLB might have been involved. This was most certainly the case in the Coorg region, north of Mysore. As indicated by Asana (1955), in that area "the most pronounced symptom of the disease is the characteristic mottling of leaves". This was a very pertinent observation, as mottling is indeed, as known today, the most characteristic symptom of HLB. Several years later, positive dot blot hybridisation confirmed HLB in mottled Coorg mandarin leaves (Varma *et al.*, 1993), and members of the 12th conference of the IOCV in India in 1992 were able to see very severe HLB on the "Coorg" mandarin trees and other citrus cultivars in Gujratnagar and Chudli.

In 1966, further support to the HLB hypothesis came from the work of Fraser and co-workers (Fraser *et al.*, 1966; Fraser and Singh, 1968). Questioning tristeza as the general cause of dieback, they indicated that CTV was not invariably present in dieback affected trees. Many of the citrus species affected in India are tolerant to tristeza in other countries. Moreover, deficiencies of zinc and other minor elements were often implicated in dieback on a symptom basis, but applications of these elements failed to cure the dieback condition, although sometimes slight and temporary improvements were obtained.

These observations suggested that HLB might have been involved in dieback, and a survey for "dieback" in all major citrus areas of India was made. A striking feature of dieback was observed during the survey, namely the fact that, in early stages of the disease, leaf symptoms were often restricted to one or a few limbs. This symptom had already been described for "greening" in South Africa by McClean and Oberholzer (1965), and for "huanglongbing" by Lin (1956) in China, but Lin's work had remained unknown, as mentioned above. It was thus concluded that dieback was "caused by the virus responsible for greening disease of citrus in South Africa". However, this conclusion was based only on symptom observations, and was not confirmed by experimental work.

Unquestionable proof for the presence of HLB in India was eventually obtained at the virus Research Center at Poona by Capoor (Fig. 1 C) and co-workers, when they succeeded in transmitting the HLB pathogen by the Asian psylla, *D. citri* (Capoor *et al.*, 1957). With this technique at hand, they were able to show that trees with dieback symptoms invariably proved positive for HLB. Among the trees positive for HLB, some carried CTV others did not. Finally, in 1971, Bové and co-workers were able to detect the HLB bacterium in a Musambi sweet orange seedling, which had been experimentally infected by *D. citri* nymphs with the "Poona" strain of HLB (and was thus free of CTV), and had been kindly provided by Prof. Capoor in 1969 (Lafèche and Bové, 1970b; Bové and Saglio, 1974). Many results on HLB in the Bordeaux laboratory have been obtained with the Poona strain.

Phloem necrosis and Vein phloem degenerations in Indonesia. In Indonesia, HLB is a major problem (Aubert *et al.*, 1985; Bové *et al.*, 2000b). Interestingly, the disease is called "Vein phloem degeneration" (Tirawidjaja *et al.*, 1955).

In an anatomical study of greening affected sweet orange shoots from South Africa, Schneider (1968) found localized pockets of necrotic phloem scattered throughout the vascular system of mature leaves, that blocked the translocation stream. Other anatomical aberrations were observed and believed to be reactions to the blockage, i.e. massive accumulation of starch in the plastids, disordered cambial activity with excessive formation of phloem, soon to become necrotic. Leaf mottle symptoms associated with HLB are very probably the consequence of these alterations.

Destruction of citrus by HLB in Thailand. HLB first appeared in Thailand in the 1960s and was so severe that the length of time between the onset of the disease and debilitation of the entire tree was about two years (Schwarz *et al.*, 1973). Roistacher (1996) has stressed the seriousness of the disease and its destruction of citrus in Thailand. HLB destroys 10 to 35% of mango trees each year and, in the northern regions, many citrus areas have gone out of production. In addition to being spread by the Asian psyllid vector, *D. citri*, the disease is also propagated by infected nursery productions. Indeed, Thailand was known as one of the rare countries in the world to use marcooting (or air layering) on a large scale to produce nursery trees.

ELECTRON MICROSCOPY AND THE BACTERIAL NATURE OF THE HLB AGENT

The HLB pathogen is a Gram negative bacterium. By 1967, at a time when the nature of HLB was still

used, it became established that "greening" was transmissible by graft inoculation as well as by the two citrus psyllids. These results suggested that the pathogen causing greening was a virus, the only plant agent known at that time to be transmitted in these ways, and it became fashionable to speak of the "greening virus". For the same reasons, the expression "stubborn virus" entered the literature (see for instance Fraser and Singh, 1968; Salbe and Cortez, 1968). It was even thought that greening and stubborn, which had some common symptoms, were caused by different strains of the same virus.

In 1967, mycoplasma-like organisms (MLOs) were found to be associated with plant diseases, the aetiologies of which were previously thought to be of virus nature (De *et al.*, 1967). Mycoplasmas are special bacteria, which lack a cell wall and are surrounded only by a single cytoplasmic membrane. Most of the "MLO diseases" were of the "yellows" type, and their symptoms resembled those of greening and/or stubborn. For these reasons, search for such MLOs in greening, and stubborn affected sweet orange leaves was initiated by electron microscopy (EM), first in Versailles, and later in Bordeaux, France.

In both cases, microorganisms were detected in the phloem sieve tubes of symptomatic leaves, but not in those of healthy leaves. In the case of stubborn, the micro-organism was soon obtained in culture, and turned out to be a new type of mycoplasma, one with helical morphology and motility, and named *Sytoplasma citri* (Saglio *et al.*, 1973). In the case of the greening organisms, which could not be obtained in culture, it was initially thought that it was also a mycoplasma (Lafèche and Bové, 1970a, b), but the agent was soon found to be enclosed by a 25 nm thick envelope, which was much thicker than the 1 to 10 nm thick cytoplasmic membrane envelope, characteristic of mycoplasmas, including *S. citri* (Fig. 3 E) (Saglio *et al.*, 1971; Garnier *et al.*, 1976). These properties suggested that the HLB organism possessed, in addition to its cytoplasmic membrane, a bacterial cell wall. This was indeed found to be the case, and it could be shown that the cell wall was of the Gram negative type, composed of a membrane and a peptidoglycan layer (Garnier and Bové, 1977; Garnier *et al.*, 1984a). Thus, the HLB agent was a Gram negative bacterium (Garnier *et al.*, 1984b). As shown by Garnier and Bové (1977), in classic specimen fixation procedures for EM, the cell wall of the HLB bacterium appears as a characteristic, electron-dense layer surrounding the bacterial cell (Fig. 3 B, D), and this suffices for HLB identification. With special fixation methods, cell membranes reveal their triple layered structure, and the HLB bacterium is seen as surrounded by two such triple layered membranes (Fig. 3 C), i.e. the inner, cytoplasmic membrane, and the outer membrane, as part of the Gram negative cell wall (Garnier and Bové, 1977; see also Moll and Martin, 1974). Under




Fig. 3. Electron micrograph of liberibacter cells in a sieve tube of sweet orange leaf (Saudi Arabia) (A). At higher magnification, the cells are seen surrounded by a cell wall, which shows up as an electron dense layer (B, D). At even higher magnification and with special techniques, one sees the triple-layered ultrastructure of both the outer cell wall membrane and the inner cytoplasmic membrane (C: $\rightarrow \rightarrow \rightarrow$), while mycoplasmas, such as *Spiroplasma citri* seen here (E: $\rightarrow \rightarrow$) have no cell wall, and are surrounded by a cytoplasmic membrane only. With special techniques, the peptidoglycane layer of the Gram negative cell wall can be visualized (F: \rightarrow on F) in between the inner membrane cytoplasmic membrane (CM on F) and the outer cell wall membrane (OM on F). The cell wall of Gram positive bacteria has no membrane, but only a heavy layer of peptidoglycane (PG on G), surrounding the cytoplasmic membrane (CM on G). All electron micrographs are from M. Garnier.




Fig. 4. Phylogenetic tree constructed with MEGA version 2.1 using 118 rDNA sequence. (GenBank accession numbers in parentheses).

these conditions, the peptidoglycane layer (PG) is generally not seen, but sometimes, the inner zone of the cell wall membrane is somewhat thicker, more electron-dense than the outer zone, and is reminiscent of a PG. EM detection of PG (Fig. 3 E, G) required the use of cytochemical techniques involving papain treatments to separate the PG from the cell wall membrane, and hexamine treatments to digest the PG and prove that it is indeed peptidoglycane (Garnier *et al.*, 1984a, b). In control experiments, identical results were obtained with the Gram negative bacterium *Escherichia coli*, but not with the Gram positive bacterium *Staphylococcus aureus* (Fig. 3 G).

Indirect evidence for the Gram negative nature of the HLB bacterium had already been obtained prior to the above EM experiments, by studying the effects of penicillin on HLB affected trees. Penicillin inhibits a late step (transpeptidation) in the biosynthesis of peptidoglycane. The antibiotic was supplied to glasshouse grown sweet orange plants through the roots (Bové *et al.*, 1980), and to field grown sweet orange trees on Trévoir orange in Reunion island by trunk injections (Aubert and Bové, 1980). In both cases, penicillin had a beneficial effect on HLB-affected citrus. Penicillin-treated citrus plants outgrew symptoms, had a much better root system, and produced larger symptomless shoots and leaves than untreated controls. No HLB bacteria could be seen in the sieve tubes of new, symptomless leaves. However, the bacteria and the symptoms reappeared when the treatments were discontinued. As expected, penicillin had no effect on *Spiroplasma citri*-affected citrus, since mycoplasmas, lacking a cell wall and having no peptidoglycane, are insensitive to penicillin.

The HLB bacterium was detected by EM for the first time in sweet orange leaves infected with South Africa "greening" (Lafliche and Bové, 1970a), but soon similar organisms could also be seen by EM in leaves infected with Reunion "greening", India "dieback", Taiwan "likubin", and Philippines "mottle leaf" (Lafliche and Bové, 1970a; Bové and Saglio, 1974). These data were confirmed for likubin by Chen *et al.* (1971) and Tanaka and Doi (1973), and for mottle leaf by Tanaka and Doi (1973). In 1979, the same organisms were also found to be associated with HLB in Southern China (Chen *et al.*, 1979a, b; Ke *et al.*, 1979). Finally, the HLB bacterium was also seen in *T. erythrae* (Moll and Martin, 1973) and in *C. citri* (Chen *et al.*, 1979).

The HLB bacterium is probably the first walled, sieve tube-restricted bacterium seen in plants. Organisms similar to the HLB organism occur in plants other than citrus, and are involved in more than 20 diseases. Papaya bunchy top (Davis *et al.*, 1996), watermelon yellow vine (Bruton *et al.*, 1998), strawberry marginal chlorosis (Zelik *et al.*, 1998), and low sugar syndrome of sugar beets (Gatineau *et al.*, 2002) are such diseases. The associated bacteria are restricted to the phloem sieve tubes,

none of them has been obtained in culture, and they are transmitted by leafhoppers or psyllids.

The publication on HLB as a mycoplasma-like organism appeared in 1970 (Lafliche and Bové, 1970a, b). However, as shown above, Bové and his group questioned the mycoplasma nature of the HLB agent as early as 1971 (Saglio *et al.*, 1971), and provided many arguments against the mycoplasma hypothesis during the 1970s, until the Gram negative nature of the HLB was finally demonstrated in 1984 (Garnier and Bové, 1984a, b). Nevertheless, the erroneous term "mycoplasma-like" was still in use in several publications until 1980.

Electron microscopy: first laboratory technique for identification and confirmation of HLB. Specific HLB symptoms do not exist. Some symptoms, such as yellow shoots, leaf blotchy mottle, and lopsided fruits with col or inversion and aborted seeds, are characteristic, but they do not always occur together on the same tree, they can be distorted or masked by symptoms of other diseases, or induced by causes other than HLB.

For instance, HLB-induced zinc deficiency is indistinguishable from genuine zinc deficiency, yellow shoots can be the result of phytophthora gummosis, and removal of a ring of bark (girdling) for air-layering produces mottle on the leaves. Stubborn and HLB have common symptoms. No countries are known where the two diseases occur together, but this situation may change in the future. For these various reasons, a technique for unequivocal identification of HLB was required.

In our laboratory, and in the hands of Monique Garnier (1949-2003) (Fig. 1 D), EM detection of the HLB bacterium has fulfilled this purpose for many years (see for instance Aubert *et al.*, 1988; Bové and Garnier, 1984; Curing *et al.*, 1978; Garnier and Bové, 1996). The reliability and specificity of EM is based on two properties of the HLB bacterium: its exclusive location in the sieve tubes (Fig. 3 A), and the presence of a cell wall (Fig. 3 B, D). In citrus, no bacteria other than the HLB agent fit these properties. For EM detection of the HLB bacterium, leaf midribs are used. If the number of bacteria per sieve tube is low, it is recommended to use longitudinal sections of sieve tube cells.

Several years of experience with EM detection of Asian and African HLB have shown that the number of bacteria in sieve tubes is higher in leaves with strong mottle than in those with mild mottle. Therefore, leaves with strong mottle are to be preferred. Symptomless leaves are useless. For indisputable identification of HLB by EM, it is necessary that at least one bacterium in one section shows the electron dense cell wall layer surrounding the cell. Most often, the layer is seen only in certain parts of the cell (Fig. 3, B, D). Sometimes, several sections have to be examined before a bacterium with a "good" cell wall is seen (Fig. 3, B, D). Finally, based on many EM examinations, no morphological differences

could be found to distinguish the Asian HLB bacterium from its African counterpart.

VARIOUS FORMS OF HLB, ASIAN, AFRICAN, AND AMERICAN

Symptoms of citrus stubborn disease and HLB are influenced by temperatures at which affected trees grow. Symptoms of stubborn are severe in hot, dry areas of California, Arizona, Morocco, Iran, and Iraq. In South Africa, HLB and the African psyllid vector, *T. erythrae*, occur in the cool, "highveld" areas of Transvaal and Swaziland, but not in the hotter, "lowveld" areas of Swaziland. However, HLB and *T. erythrae* are present in the Cape Town region (Garnier *et al.*, 2000a), where the southern latitude compensates for the low altitude, and is responsible for a temperate climate.

In Madagascar, HLB and *T. erythrae* only occur on the central, high plateau (1200 to 1300 m a.s.l.), but not on the low lying coastal areas. In Senegal, HLB and *T. erythrae* are not seen below an altitude of 600 to 700 m. On the contrary, in Asia, HLB and the Asian psyllid vector, *D. citri*, are found in hot low altitude areas. For instance, in Indonesia, in 1984, HLB and *D. citri* were present in the whole of Java coastal lowlands, but many trees in orchards located between 800 and 1200 m a.s.l. were still healthy (Aubert *et al.*, 1985). In North Bali, HLB was commonly present below an altitude of 600 m, but was rarely seen above 1000 m, and its distribution reflected that of *D. citri* (Bové *et al.*, 2000b). Similarly, in the Ningnan county of Sichuan (South China), 100% of trees were affected by HLB in orchards located at altitudes of 1000 to 1400 m, where *D. citri* was abundant, while between 1385 and 1620 m, the percentage went down to 3% and no psyllids could be found (Zhao, 1981).

In the frame of an international cooperation experiment started in 1969, the influence of temperature on HLB symptoms was also studied under phytotron conditions in two temperature controlled chambers: a "cool" chamber (24°C with a 16-hr light period and 32°C with a 8-hr dark period) and a "warm" chamber (32°C with a 16-hr light period and 27°C with a 8-hr dark period) (Bové *et al.*, 1974). Sweet orange plants graft inoculated with the following HLB sources were used: South Africa (Nelspruit) "greening", India (Ponza, "dieback"), and Philippines (Lipa City) "mottle leaf". The presence of HLB bacteria in inoculated plants was established by EM.

With South African HLB, severe symptoms were obtained after 30 weeks in the cool chamber, and the plants averaged 35 cm in height; no symptoms developed in the warm chamber, and the plants reached 190 cm. When, after 40 weeks in the cool chamber, severely affected plants were transferred to the warm conditions, they quickly produced new, vigorous growth, recovered, and

remained symptomless during the remaining 10 months of the experiment. With both African HLB and Philippine HLB, symptoms were as pronounced at 22-24°C as at 27-32°C. After 30 weeks in the warm chamber, symptomatic seedlings infected with Indian HLB and Philippine HLB measured 25 and 40 cm, respectively, while healthy controls had grown to 180 and 140 cm.

This experiment as well as the field observations show that HLB in Africa is heat-sensitive, and occurs only in cool areas, with temperatures remaining below 30-32°C. Similarly, the African psyllid vector, *T. erythrae*, thrives only in cool environments, and is also sensitive to high temperature combined with low relative humidity (Carling, 1969c). On the contrary, HLB in Asia is heat-tolerant, and symptoms occur even when temperatures are well above 30°C. The Asian psyllid vector, *D. citri*, has similar properties, and is also heat-tolerant.

The South African Nelspruit strain, and the Indian Ponza strain of the HLB bacterium could be transmitted to periwinkle (*Catharanthus roseus*) plants (see below). Temperature experiments conducted with infected periwinkle plants gave the same results than those obtained above with HLB-affected citrus seedlings: with the Indian HLB bacterium, symptom expression occurred at both 25 and 32°C, but only at 25°C with the South African strain (Garnier and Bové, 1985). The effect of temperature on symptom expression is thus the same in periwinkle and citrus. Therefore, the temperature effect is due to the HLB bacterium and not to the plant, the African HLB bacterium being heat-sensitive and the Asian HLB bacterium, heat-tolerant. This biological difference indicates that the African and Asian HLB bacteria are not identical. Indeed, as shown below, they represent different bacterial species: *Candidatus Liberibacter africanus* and *Candidatus Liberibacter asiaticus*.

The fact that the African HLB bacterium and *T. erythrae* are both heat-sensitive, and the Asian HLB bacterium and *D. citri*, both heat-tolerant, seems to be a good example of adaptation between vector and pathogen. It has to be remembered however, that, experimentally at least, each one of the two vectors can transmit each one of the two HLB bacterial species (Massoné *et al.*, 1976; Lallemant *et al.*, 1986).

In the State of São Paulo, Brazil, *Ca. L. asiaticus* has been found in less than 10% of the HLB-affected trees. Indeed, a third bacterial species has been discovered: *Candidatus Liberibacter americanus* (see below), which affects more than 90% of the trees. In São Paulo State, the Asian psyllid vector, *D. citri*, was reported as early as 1942, and transmits not only *Ca. L. asiaticus*, but probably also *Ca. L. americanus*. Indeed, the new liberibacter could be detected by PCR within *D. citri* psyllids collected on *Ca. L. americanus*-infested trees. Furthermore, in the glasshouse, sweet orange seedlings, graft inoculated with *Ca. L. americanus*, showed severe HLB leaf mottle at both cool (22-24°C) and warm (27-32°C) conditions (Teixeira

et al., 2005c). These observations suggest that the South American (Brazil) HLB is of the heat-tolerant form.

In Florida, the presence of *D. citri* was reported in 1998 (Halbert, 1998). HLB has been observed in the Southern part of the State in August 2005, and only *C. L. asiaticus* has been detected in HLB-affected trees. Therefore, the North American (USA) HLB is probably also of the heat-tolerant form.

TRANSMISSION OF THE HLB BACTERIUM FROM CITRUS TO PERIWINKLE BY DODDER

In citrus, the HLB bacterium is present only in small numbers, but in dodder (*Cuscuta campestris*) the organism reaches high titres (Ghosh *et al.*, 1977). Periwinkle (*Catharanthus roseus*) is a host plant in which also many mycoplasma-like organisms multiply actively in sieve tubes. These observations suggested to attempt transmission of the HLB bacterium from citrus to periwinkle through dodder. Positive transmissions were indeed obtained (Garnier and Bove, 1983). In the case of Indian HLB, one of four periwinkle plants connected via dodder to a HLB-affected sweet orange seedling developed peculiar yellowing symptoms, and contained HLB-like bacteria in sieve tubes, some of which were particularly rich in bacteria. Shoots from the symptomatic periwinkle were top-grafted onto healthy periwinkle plants, which developed the characteristic symptoms after 3 months at 25°C. In this way, large numbers of symptomatic, HLB-infected periwinkle plants were produced. Similar results were obtained with South African, Chinese, and Philippine strains of the HLB organism. It must be stressed that periwinkle plants naturally infected with the HLB agent have never been observed. Citrus psyllids unlike periwinkle, are unable to transmit the HLB agent to this plant, and die when forced to stay on it.

Ultrathin sections from HLB-infected periwinkle plants were used in EM to demonstrate the Gram-negative nature of the bacterium (Garnier *et al.*, 1984a, b) (see above). As the HLB bacterium is not available in culture, great numbers of infected periwinkles have been used as convenient sources for the phylogenetic and taxonomic characterization of the HLB bacterium, the production of monoclonal antibodies, and the development of molecular detection techniques (see below).

Transmission of the HLB bacterium to tobacco (*Nicotiana glauca* Xanth.) by dodder has also been obtained (Garnier and Bove, 1993).

PHYLOGENETIC AND TAXONOMIC CHARACTERIZATION OF THE AFRICAN AND ASIAN HLB BACTERIA

16S rDNA. To determine the phylogenetic position of the HLB bacterium, the 16S ribosomal DNAs (16S

rDNA) of the South African Nelspruit strain and the Asian Punoa strain were obtained from total DNA of HLB-infected periwinkle plants, by polymerase chain reaction (PCR)-amplification, using the universal PCR-primers (DL/r-P1 for amplification of prokaryotic 16S rDNA (Weisburg *et al.*, 1991). In these experiments, care must be taken to prevent interference by periwinkle mitochondrial and chloroplast 16S rDNAs. This can be achieved by cutting mitochondrial 16S rDNA with endonuclease *Bcl*I before PCR amplification. The amplified DNA, about 1500 base pairs (bp) in size, contains bacterial as well as chloroplast 16S rDNA. The presence of bacterial DNA can be ascertained by *Eco*RI, which cuts it into two fragments (~650bp and ~850bp), while chloroplast DNA is uncut. The 16S rDNAs were cloned from the ~1500bp amplified products, and sequenced. Hybridization and PCR experiments performed with oligonucleotides specific for the amplified sequences revealed that the DNAs obtained were indeed the 16S rDNAs of the HLB bacterium, and not the DNA of a contaminating organism (Gagnoux *et al.*, 1994).

Comparisons with 16S rDNA sequences obtained from the GenBank database showed that the African and Asian HLB bacteria belonged to the α subdivision of the class *Proteobacteria*. Even though their closest relatives were members of the α -2 subgroup, the HLB bacteria were distinct from this subgroup, as the level of 16S rDNA sequence identity was only 87.5%. Therefore, the two HLB bacteria represented a new lineage in the α subdivision of the class *Proteobacteria*. As the members of this class are Gram-negative bacteria, these results confirmed the Gram-negative nature of the HLB bacterium, a property which had already been deduced from earlier EM studies by Garnier *et al.* (1984a, b).

The α subdivision of *Proteobacteria* is a diverse group of microbes that includes both plant pathogens or symbionts with some distinctive properties (*Agrobacterium tumefaciens*, *Bradyrhizobium* spp.) and human pathogens (*Rochalimna* spp., *Bartonella bacilliformis*, *Brevetia abortus*, *Aflpie* spp., etc.). Organisms in this group live in intimate association with eukaryotic cells and, in many cases, have acquired the ability to survive and grow within an arthropod vector. The HLB organism fits this description remarkably well. Indeed, it grows in a specialized niche in its eukaryotic plant host, the phloem sieve tubes, and it is transmitted by two arthropod vectors, the psyllids *T. erytraea* and *D. citri*, in which it multiplies in the hemolymph and within the salivary glands.

RpoKJL-rpoBC operon and DNA probes. Using total DNA from periwinkle plants infected with the Asian Punoa strain, several DNA fragments of the bacterial genome could be obtained by random cloning (McLachlan *et al.*, 1992).

Fragment In-2.6 (2.6 kbp), when used as a probe in Southern- or dot-hybridisations, hybridised at high stringency with all tested strains of the Asian HLB bacterium but not with the African strain. At low or intermediate stringency, some hybridisation was also seen with the African strain. The Asian strains that gave positive hybridisation signals with probe Poona In-2.6 were those from Poona (India) (homologous strain), Nakhom Pathon (Thailand), Lipa city (Philippines), Fujian (China), Taiwan, and Bali (Indonesia).

By sequencing, In-2.6 was found to be part of the *spkAHL-ropBC* gene cluster, the well-known, bacterial β -operon, which codes for ribosomal proteins K, A, J, and L, and RNA polymerase subunits β and β' (Vilchevaux *et al.*, 1993). From the sequence of In-2.6, two PCR primers were designed, F-1898 and r-1897, and used to amplify part of the *spkAHL-ropBC* operon of the African Nelspruit HLB bacterium (Planet *et al.*, 1993). A clear DNA band of about 1900 bp was obtained. Upon cloning and re-sequencing, the DNA from the African strain (1676 bp) was indeed found to correspond to part of the expected β operon. It was called AS-1.7. The AS-1.7 DNA hybridised at high stringency with DNA from periwinkle or citrus plants infected with the African Nelspruit strain, but no hybridisation was observed in the case of the Asian strains tested (Planet *et al.*, 1993). As indicated above, opposite results were obtained when In-2.6 was used as the probe: no hybridisation with the African strain, but strong hybridisation with all Asian strains tested. The overall nucleotide identity between In-2.6 and AS-1.7 was 74.2%. This relatively low homology for similar organisms explains why no hybridisation was observed between In-2.6 and DNA from periwinkle plants infected with the African Nelspruit HLB bacterium, and vice versa. This low homology also suggested that the African strains of the HLB bacterium and the Asian strains were members of two different species of the same genus (Jaguoux *et al.*, 1994; Planet *et al.*, 1993).

Monoclonal Antibodies. Since 1987, only thirteen different monoclonal antibodies (MA) specific for the HLB bacteria have been produced (Garnier *et al.*, 1991; Gao *et al.*, 1993). This low number is explained by the fact that the HLB bacteria are not available in culture. The first ten MAs were raised using as immunogen homogenates of phloem tissue from HLB-affected periwinkle plants. Of these MAs, two (including MA 10A6) were against the Indian Poona strain, five against a strain from China (Fujian), and three against the South African Nelspruit strain. The use of these MAs for the detection of HLB bacteria has shown that each MA is very specific for the strain used for immunisation and, therefore, they cannot be used for generalised diagnosis of HLB (Garnier *et al.*, 1991).

In attempting to produce antibodies recognizing most

or all strains of the HLB bacterium, an antigenic protein of the Indian Poona strain was purified by immunofluorescence chromatography using MA 10A6 directed against this protein, and used for *in vitro* immunisation of spleen cells (Gao *et al.*, 1993). Three MAs were obtained, one of which (1A3) recognized all Asian strains tested except the Chinese one, whereas the other two recognized most of the Asian, but not the Chinese strain. None of the three MAs reacted with the South African strain. These results agreed with those obtained with DNA probes, confirming that Asian and African HLB bacteria are members of two different bacterial species.

MA 10A6, coupled to CNBr-activated sepharose 4B, has also been successfully used to purify the Poona BLO by immunofluorescence (Vilchevaux *et al.*, 1993). In this way, purified cells of the HLB bacterium could be observed for the first time in the electron microscope.

Candidatus genus and species: Candidatus Liberibacter africanus and Candidatus Liberibacter asiaticus. Bacteriologists have had a conservative attitude when it came to give Latin binomial names to non-cultured organisms. However, with the development of DNA amplification by PCR and DNA sequencing, it became possible to characterise such organisms at the molecular and phylogenetic level. On the basis of such considerations, Murray and Schleifer (1994) proposed the "Candidatus" designation as an interim taxonomic status, to provide a proper allocation of sequence-based potential new taxa at the genus and species level. One of the first non-cultured bacteria to benefit from the *Candidatus* proposal was the HLB bacterium, shown by 16S rDNA sequence comparisons to be the first member of a new subgroup in the α subdivision of the *Proteobacteria* (see above). The trivial name *liberibacter* (Jaguoux *et al.*, 1994), but replaced by *liberibacter* (Garnier *et al.*, 2000b) (from the Latin *liber* (bark) and *bacter* (bacterium)), was given to organisms in this new subgroup.

As indicated above, the HLB *liberibacter* strains from Africa can be distinguished from those in Asia on the basis of temperature sensitivity (Bové *et al.*, 1974), DNA hybridizations and genomic properties (Vilchevaux *et al.*, 1992, 1993), and serology (Garnier *et al.*, 1991; Gao *et al.*, 1993). For these reasons, they represent two different species. Therefore, following the *Candidatus* proposal of Murray and Schleifer, the HLB *liberibacter* from Asia should be denoted *Candidatus Liberibacter asiaticus*, and the HLB *liberibacter* from Africa *Candidatus Liberibacter africanus* (Jaguoux *et al.*, 1994; Garnier *et al.*, 2000b).

The sequence variability of the region between the 16S rRNA gene and the 23S rRNA gene in the ribosomal operons is useful for differentiating species within genera. This 16S/23S intergenic region was obtained for two Asian strains of the HLB *liberibacter*, the Indian Poona strain and a strain from Fuzhou, Sichuan (China), as well

as for the African Nelsonii liberibacter (Jugoueix *et al.*, 1997). The intergenic regions of the two Asian Liberibacters had 100% sequence identity, even though they belong to different serotypes and were isolated from geographically distant areas. However, the sequence identity between an Asian liberibacter strain and the African liberibacter was only 79.46%, confirming the fact that the African liberibacter and the Asian liberibacter belong to two different species. Similar results have also been obtained by Subandiyah *et al.* (2000).

Candidatus Liberibacter africanus subsp. *capensis*

A third liberibacter was detected by PCR in an ornamental nutmegs tree, Cape chestnut (*Colodendrum capense*), in the Cape region of South Africa. Leaves of the affected tree showed characteristic mottling. The new liberibacter was characterized by serology and the sequences of its 16S rDNA, the intergenic 16S/23S rDNA region, and ribosomal protein genes of the β operon. Phylogenetic analysis showed the new liberibacter to be more closely related to *C. L. asiaticus* than to *C. L. asiaticus*, and a subspecies status was assigned to it, *C. L. africanus* (*Liberibacter africanus* subsp. *capensis*) (Garnier *et al.*, 2000a).

16S rDNA-based phylogeny tree. Fig. 4 shows the 16S rDNA-based phylogeny tree of liberibacters. The tree was constructed from 16S rDNA sequences obtained from *C. L. asiaticus*. It can be seen that *C. L. asiaticus* strains cluster together, and the "Asian" cluster is close to the "African" cluster. The "Asian" cluster and the "African" cluster form the "Liberibacter" and "Asian" liberibacter group. The new liberibacter from St. Paul's State, *C. L. africanus* (see below), is not part of this group, but forms a separate branch, indicating that it is a species different from *C. L. asiaticus* and *C. L. africanus*. All liberibacters are members of the α subdivision of the class *Proteobacteria*, and their closest relatives are members of the $\alpha 2$ subgroup. The tree also shows two members of the γ subdivision.

Recently a phylogenetic tree was constructed from the *ospA* gene sequences of the African and Asian liberibacters, and confirmed the phylogeny based on 16S rDNA sequences (Bastiani *et al.*, 2005). The *ospA* gene sequences were also used to study the diversity of several strains of *C. L. asiaticus*. Each strain could be characterized by a specific PCR-RFLP profile.

IDENTIFICATION OF HLB BY DETECTION OF THE AFRICAN AND ASIAN LIBERIBACTERS

Until 1992, electron microscopy visualization of the walled HLB organelles in the sieve tubes of citrus leaves showing blotchy mottle was the only reliable method of detection and was widely used (Garnier and Bové,

1996). However, the technique was heavy, and unable to distinguish between Asian and African liberibacters. This differentiation has now become possible, with the development of molecular techniques such as DNA hybridization and PCR. As mentioned, monoclonal antibodies are too specific to be used for diagnosis.

DNA Hybridisation. In dot hybridisation, probe Ja 2.6 (see above) gives positive hybridisation signals with DNA isolated from citrus leaves infected with Asian liberibacter strains, while probe AS L7 reacts positively with African liberibacter strains (Vilchevaux *et al.*, 1992, 1993; Planet *et al.*, 1995). These probes can also be used very efficiently to detect Liberibacters in psyllid insect vectors. Individual insects are crushed onto a nylon membrane and the membrane or "crush blot" is submitted to hybridisation with one or the other probe (Bové *et al.*, 1994). Non-reactive probes have been developed (Horquellat *et al.*, 1997).

PCR. Two PCR systems have been used. The first is based on the amplification of a 1160 bp fragment of liberibacter 16S rDNA (Jugoueix *et al.*, 1996). The primer pair OD/OD2 is able to amplify the rDNA of both liberibacter species, while the pair OA1/OD2 amplifies preferentially the African liberibacter rDNA. In countries where the two liberibacter species are known or suspected to be present, it is advisable to use the two forward primers, OD1 + OA1, and the common reverse OD2 primer in the same PCR mixture. Sequence analysis shows that the rDNA amplified from the Asian liberibacter has no *NheI* restriction site and yields, once *NheI* treatment, two fragments 520 bp and 640 bp in size, respectively. The rDNA amplified from the African liberibacter has an additional site and yields three fragments with a size of 520 bp, 506 bp, and 130 bp. Hence, by *NheI* treatment of the amplified DNA, it is easy to identify the liberibacter species present in a given sample (Jugoueix *et al.*, 1996).

The second PCR system is based on the sequence of the *spkA*11, *spB*C operon, which is slightly different from one liberibacter species to the other. In particular, the intergenic region between genes *spA* and *spB* is 34 bp larger in the Asian than in the African liberibacter. With forward primer (*spA*2), selected in the *spA* gene, and reverse primer (*spB*) from the *spB* gene, a 703 bp DNA is amplified from the Asian liberibacter, while a 669 bp DNA is obtained with the African liberibacter. When both liberibacter species are present in the same sample, amplification of the two DNAs is obtained, and upon agarose gel electrophoresis two DNA bands are seen, the upper (703 bp) corresponding to the Asian liberibacter, and the lower (669 bp), to the African liberibacter (Horquellat *et al.*, 1999). Finally, a PCR assay was developed, based on the β operon sequence and specific for the detection of *Candidatus Liberibacter*

strawus subsp. *canensis* (Garnier *et al.*, 2000b).

By the use of these molecular techniques, the presence of HLB has been clearly established in several African and Asian countries (Bové *et al.*, 1993; Bové *et al.*, 1996; Bové *et al.*, 2000a; Dee-Doe *et al.*, 2003; Garnier and Bové, 1996; Garnier and Bové, 2000; Garnier *et al.*, 2003; Keister *et al.*, 1996; Regmi *et al.*, 1996; Varma *et al.*, 1993). The presence of both liberibacter species, sometimes in the same trees, was emphasized in Scotland and Mauritius islands (Garnier *et al.*, 1996).

HLB IN SÃO PAULO STATE, BRAZIL: OLD AND NEW LIBERIBACTERS

Even though HLB was reported in São Paulo State (SPS) only in 2004, the Asian psyllid vector of HLB, *D. citri*, has been present in Brazil at least since 1942, when it was reported for the first time (Lima, 1942). Specimens of *D. citri* from Brazil were present at the British Museum in 15 of August, 1969, (personal communication to Cating, 1970). In certain years, high populations of the insect could be seen in SPS in sweet orange groves.

Discovery of a new liberibacter: *Candidatus Liberibacter americanus*. In March 2004, symptoms of HLB were observed in sweet orange trees near the city of Aracaju in SPS (Anonymous, 2004; Colletta Filho *et al.*, 2004; Aguiar *et al.*, 2005; Teixeira *et al.*, 2005a, b). This was the first reported case of HLB from the American Continent. Affected trees could be spotted from a distance because of their conspicuous yellow shoots. Leaf symptoms of HLB in SPS were very similar, if not identical, to those in Africa and Asia, with characteristic and pronounced blotchy mottle, as described by McClean and Schwarz (1970) on both small and large leaves (Fig. 6–7). Fruits were small and lopsided, exhibited strong colour inversion, seed abortion, and brown/orange-stained vascular bundles (Fig. 8, 9). These fruit symptoms were more similar to those seen in China than to those observed in South Africa. A survey conducted in September 2004, only six months after the disease was recognized, showed HLB to be already present in 40 municipalities of SPS. A few orchards had more than 50% of affected trees. These observations suggested that the disease had been present for almost ten years, but without being properly diagnosed. History repeats itself.

In April 2004, the 16S rDNA based PCR technique with forward primers OAI+OD1 and reverse primer OI2c, described by Jaeger *et al.* (1996), was applied in sweet orange leaves with strong mottle from suspicious trees in SPS to confirm the presence of HLB in the area and identify the liberibacter involved. This PCR method had been assigned in many Asian and African countries for the detection of the two HLB liberibacters (see for

instance Bové *et al.*, 1996, 2000b; Garnier and Bové, 1996; Garnier *et al.*, 1996). Whenever leaves with the classic blotchy mottle symptoms were used, positive PCR reactions were obtained, and yielded the characteristic 1160 bp amplicon. Unexpectedly in SPS, only negative PCR reactions were obtained from mottled leaf samples from 10 affected trees, many of which had also severe fruit symptoms. Under the same testing conditions symptomatic control citrus leaves infected with *Ca. L. asiaticus* or *Ca. L. africanus* from the HLB collection in Maribon, gave positive PCR reactions (Teixeira *et al.*, 2005a, b). However at the same time, the using practically the same PCR technique, *Ca. L. asiaticus* was detected in 2 of 10 leaf samples by other workers (Colletta Filho *et al.*, 2004). Because of the repeatedly negative PCR reactions, the presence of a new bacterial pathogen in many of the symptomatic, blotchy mottle leaves from SPS was suspected and investigated.

Evidence for the presence of a new HLB bacterium (SPS HLB bacterium) was obtained by PCR amplification with universal primers 133 and rPL for prokaryotic 16S rDNA amplification. The 16S rDNA of the SPS HLB bacterium was cloned and sequenced, and the sequencer was used to design the specific primer pair E-GB17c/GB17c PCR amplification, with these primers made it possible to detect the SPS HLB bacterium in all HLB leaf samples testing negative for *Ca. L. asiaticus* and *Ca. L. africanus*. Most leaf samples (98%) were infected with the SPS HLB bacterium, the remaining samples (2%) carried *Ca. L. asiaticus*. Primer GB17c, complementary to GB17c, was used in conjunction with reverse primer 23S1 to amplify, clone, and sequence the 16S/23S ribosomal intergenic region (RIR) (a total, a 1473 bp sequence of 16S rDNA (almost the complete 16S rDNA sequence) followed by the complete 583 bp sequence of the RIR, was available for characterization of the SPS HLB bacterium by comparison with similar sequences of various isolates of *Ca. L. asiaticus*, and the Nelspruit isolate of *Ca. L. africanus* (Teixeira *et al.*, 2005a).

These comparisons clearly showed that the SPS HLB bacterium was not only a member of the genus *Candidatus Liberibacter*, having all the oligonucleotide signatures of the liberibacters, but was a new species of this genus. In particular, in the 16S rDNA phylogenetic tree (Fig. 4), all isolates of *Ca. L. asiaticus* clustered together within the *Ca. L. asiaticus/Ca. L. swazicus* group, but the SPS HLB bacterium did not, and formed a separate branch. Also, the RIR sequences of different isolates of *Ca. L. asiaticus* were identical or almost identical (100% identity). However, the RIR of the SPS HLB bacterium and that of *Ca. L. asiaticus* had only 78% sequence identity. With *Ca. L. africanus*, the sequence identity was even lower: 66%.

For the above reasons the SPS HLB bacterium was denoted *Candidatus Liberibacter americanus*, sp. nov. (Teixeira *et al.*, 2005a, b, c). This designation refers to

the fact that the new liberibacter species was detected for the first time in the American Continent, and that it represented the major *Ca. Liberibacter* species associated with HLB in the affected SPS region. The designation is in line with the other *Ca. Liberibacter* names, which also refer to the continents where they occur: Africa for *Ca. L. africanus*, and Asia for *Ca. L. asiaticus*.

Additional properties of the SPS-HLB bacterium fit those of the other two liberibacters. Transmission to healthy orange seedlings by graft inoculation was obtained, and EM observations showed the American liberibacter to be restricted to sieve tubes (Teixeira *et al.*, 2005a; Tanaka *et al.*, 2004).

PCR detection of *Ca. L. americanus* and *Ca. L. asiaticus* in citrus leaves. For the specific PCR detection of *Ca. L. americanus*, forward primer GB1 and reverse primer GB3 were designed from the 16S rDNA sequence of the new liberibacter (accession number AY742824). These primers, as well as those specific for *Ca. L. africanus* and *Ca. L. asiaticus* (OA1+OI/OI2c), were used for detection of the three liberibacters in each leaf sample according to Teixeira *et al.* (2005b). A first aliquot of the DNA from a leaf sample was used for the detection of *Ca. L. americanus* with primers GB1/GB3, yielding an amplicon of 1027 bp, and a second aliquot served for the detection of *Ca. L. africanus* and *Ca. L. asiaticus* with primers OA1+OI/OI2c, giving an amplicon of 1160 bp.

In this way, from August 2004 to September 2005, 1525 leaf samples gave positive PCR reactions. Each sample came from a single tree with leaves showing blotchy mottle, ranging from mild to strong. Of 1525 samples, 1411 (92.5%) were found to be infected with *Ca. L. americanus*, 82 contained *Ca. L. asiaticus* (5.4%), and both liberibacter species were found in 32 samples (2.1%). *Ca. L. africanus* was not detected. Leaf samples for liberibacter detection and identification were collected from many citrus farms representing all 79 HLB-affected SPS municipalities. For these reasons, the results obtained are meaningful, and indicate that the major liberibacter in SPS is *Ca. L. americanus*, and that both *Ca. L. americanus* and *Ca. L. asiaticus* can be found in the same tree. Even though as many as 79 municipalities host HLB-affected trees, 95% of these trees are in only 10 municipalities, indicating that most municipalities have a low rate of infection, with less than 0.1% diseased trees in the affected farms.

In another set of assays, of 216 leaf samples infected with *Ca. L. americanus*, 208 were from sweet orange trees ('Chenour', 'Hamler', 'Lima', 'Nara', 'Pera', 'Valencia', and 'Westin'), 5 from 'Ponkan' mandarin trees, 1 from a 'Murcott' tangerine tree, and 2 from 'Cravo' mandarin trees. These proportions reflect essentially the fact that sweet orange is by far the major cultivar in SPS, but they also indicate that HLB is not restricted to sweet orange.

In the above experiments, blotchy mottled leaves from HLB-affected trees, i.e. trees showing leaf and fruit symptoms, never failed to give positive PCR reactions, which emphasizes the importance of using symptomatic leaves for HLB diagnosis.

The presence of *Ca. L. americanus* and/or *Ca. L. asiaticus* in symptomless leaves has also been investigated. Symptomless leaves were collected from symptomless parts of infected trees, from symptomless trees adjacent to symptomatic trees, and from trees in a region not affected by HLB. As expected from previous results with *Ca. L. africanus* and *Ca. L. asiaticus*, all symptomless leaves tested PCR-negative. This result is not due so much to lack of sensitivity of PCR, as similar results were also obtained with the more sensitive nested PCR, but probably reflects the uneven distribution of liberibacters in recently infected trees, and the difficulty of sampling liberibacter-infected leaves when no symptoms are present to guide the choice. For the same reasons, indexing by PCR symptomless nursery or orchard trees for HLB presence seems meaningless.

Several new developments in diagnosis have taken place. Thus, "multiplex" PCR is being developed for routine detection of all three liberibacters in one test tube with a single reaction mixture containing the two sets of primers described above: (i) f-GB1 and r-GB3 for 16S rDNA amplification of *Ca. L. americanus*, and (ii) f-*gPA2* and r-*sp15* for *rpm* gene amplification of *Ca. L. asiaticus* and/or *Ca. L. africanus* (Teixeira D. do Carmo, unpublished results). Nested PCR is also being developed (Li and Ke, 2002; Ding Fang *et al.*, 2004, 2006; Li *et al.*, 2005; Hartung *et al.* (2005a) have compared several methods for the detection of the Asian liberibacter.

PCR detection of *Ca. L. americanus* and *Ca. L. asiaticus* in *Diaphorina citri* psyllids. By the end of 2004, *Ca. L. americanus* was present in 46 municipalities of SPS, and one year later the number had increased to 79, indicating a rapid spread of the liberibacter. As mentioned, the psyllid vector of *Ca. L. asiaticus*, *Diaphorina citri*, was established in Brazil since the 1940s. The rapid spread of *Ca. L. americanus* suggested that a vector was involved, and that it might be *D. citri*. Thus, psyllids were collected in August 2004 on three Pera sweet orange trees with severe symptoms of HLB and shown by PCR to be infected only with *Ca. L. americanus*. The insects were subdivided into 22 batches of 10 individuals each. The 22 batches gave negative PCR reactions with the primers specific for *Ca. L. africanus* and *Ca. L. asiaticus*. However, 6 batches gave positive PCR signals with the primers specific for *Ca. L. americanus* (Teixeira *et al.*, 2005b).

In an additional experiment, psyllids were collected in a severely HLB-affected sweet orange orchard from: (i) symptomatic branches of symptomatic trees (S/S psyllids); (ii) asymptomatic branches of symptomatic trees

AS/S psyllids), and (iii) asymptomatic trees (AS/AS psyllids). The psyllids were subdivided in batches of 10 insects. *Ca. L. americanus* was detected in 5 of 56 batches of AS/AS psyllids, 13 of 56 batches of AS/S psyllids, and 27 of 76 batches of S/S psyllids. *Ca. L. asiaticus* was detected in only 2 batches of S/S psyllids. These data indicate that in a severely affected orchard (c.50% HLB trees), symptomless branches of symptomatic trees carry as many inoculation-competent psyllids as symptomatic branches, probably because of psyllid movement, and that infected psyllids, probably coming from symptomatic trees, can be found on trees with no symptoms. Also, *Ca. L. americanus* was found in a total of 45 psyllid batches, and *Ca. L. asiaticus* in 2. This proportion is quite similar to the ratio of *Ca. L. americanus* infected trees to *Ca. L. asiaticus* infected trees in SPS.

Finally, the American liberibacter was also detected in one batch of 10 psyllids collected on a *Moraya pumilata* plant close to a HLB-affected citrus orchard. In China, *Ca. L. asiaticus* has also been reported to infect *M. citrifolia* (Li and Ke, 2002).

These results, as a whole, clearly suggest that *D. citri* is a vector of *Ca. L. americanus* in SPS.

PCR detection of *Ca. L. americanus* in *Moraya paniculata* leaves. *M. paniculata* (buxus orange), an ornamental woody shrub or tree, has a wide distribution throughout SPS, and represents the principal host of *D. citri*. *M. paniculata* plants with suspicious foliar symptoms have been observed in HLB-affected citrus orchards in SPS. *Ca. L. americanus* was detected in symptomatic leaves from 3 of 13 such plants, but not in asymptomatic leaves of the same plants (Jones *et al.*, 2005). In China, *Ca. L. asiaticus* was reported to infect also *M. paniculata* by Li and Ke (2002) but not by other researchers (Jones *et al.*, 2005).

In view of these conflicting results, it is essential to confirm if *M. paniculata* is a host of *Ca. L. americanus* and *Ca. L. asiaticus*. Should this be confirmed, *M. paniculata* would become an additional source of liberibacter inoculation available for psyllids, thus calling for eradication of *M. paniculata* plants present in citrus farms. Removal of such plants from public parks, streets and avenues, as well as bus stop gardens, must also be seriously discussed. Planting and growing new *M. paniculata* plants should probably be prohibited. Relatedly, other citrus relatives, *Strychnos benzoides* (Burmese box orange) and *Lycopersicon lycium* (wood apple) (Huang *et al.*, 2006), and *Clausena laurolepis* (Chinese sumpter) (Ding *et al.*, 2006) have been shown to be hosts of *Ca. L. asiaticus*.

DIAPHYORINA CITRI AND HLB IN FLORIDA

The Asian HLB-vector, *D. citri*, was detected in Palm Beach County, Florida, in June 1995 in *huckiyanti* plant

ings of *M. paniculata*, and by 2001, it was found in as many as 31 counties (Halbert, 1998, 2005; Halbert *et al.*, 2003, 2004). It is believed that the psyllid had been present in South Florida 6 to 12 months prior to its discovery, so that eradication was no longer feasible at the time of detection. Today, the insect is throughout Florida, essentially as a result of distribution of *M. paniculata* plants, largely through discount stores (S. Blake, personal communication).

HLB was discovered in Miami-Dade County, Florida, on two samples of a pummelo (*Citrus grandis* L.) Osh tree in August 2005, seven years after detection of the vector in the same region (Halbert, 2005). By mid-October, as a result of additional surveys, the disease was found in many residential properties stretching northwards over 250 km from Miami-Dade County to St. Lucie County. Several commercial citrus farms were also affected in Palm Beach and Volusia Counties. In view of the large area already affected, it appeared that the disease had probably been present in Southern Florida since quite some years, a situation excluding its eradication.

PCR was used to confirm HLB and *Ca. L. asiaticus* was detected by PCR in pummelo, grapefruit, sour orange, sweet orange, Key lime, lemon, kumquat, and citrimonday, as well as in *D. citri* psyllids, but not in *M. paniculata* (Sprunt *et al.*, 2005).

WORLD DISTRIBUTION OF HLB, NATURE OF THE LIBERIBACTERS AND THE PSYLLID VECTORS INVOLVED

Asia, Southeast Asia, and Oceania. *D. citri* is the vector, *Ca. L. asiaticus* is the HLB agent, and both are heat-tolerant (Asian form of HLB, see above). As an region or countries where *Ca. L. asiaticus* has been detected by EM, DNA hybridisation and/or PCR include the Indian subcontinent, India, Pakistan, Nepal, Bhutan, Bangladesh, Sri Lanka, Indonesia (Myanmar, Thailand, Malaysia, Cambodia, Laos, Vietnam), South-eastern China, Taiwan, Southern Japan (Ryukyu islands, Okinawa) (Miyawaki and Tsunoi, 1989), Philippines, Indonesia (Java, Sumatra, Eastern Kalimantan, Sulawesi, Sulawesi, Bali), East Timor, Papua New Guinea.

Africa and Madagascar island. *D. citri* is the vector, *Ca. L. africanus* is the HLB agent, and both are heat-sensitive (African form of HLB). East and South African regions and countries where *Ca. L. africanus* has been detected by EM, DNA hybridisation and/or PCR include South Africa, Zimbabwe, Malawi, Burundi, Kenya, Senegal, Ethiopia. In West Africa only Cameroon is involved.

Arabian Peninsula. In Saudi Arabia, HLB is present




Fig. 5. Psyllid vectors of HLB. *Diaphorina citri* feeding on the lower side of a sour orange leaf in Peshawar, Pakistan (A), sweet orange leaf in Guangxi, China (B), and citrus shoot in Reunion island (C, courtesy of B. Aubert). Notice position of insect at a 45° angle. Heavily curled citrus leaves infested by *D. citri* nymphs and adults (D). Yellow eggs of *Trioza erytreae* along the midrib and on the edges of a lemon leaf (E). *T. erytreae* nymphs develop on the underside of the leaves, becoming embedded in pits or nests, which look like bumps on upper side of the leaves (G). When adult psyllids emerge from nymphs, the pits are left empty (F). Bumps on the upper side of citrange leaves (H). Nymphs on the underside of citrange leaves (I). *T. erytreae* nymphs parasitized by *Tamarixia dryi*, that emerges through a hole (J).

along the Red Sea region from Mecca to Najran, close to the Yemeni border. The vector is *D. citri*. HLB occurs in hot oases, and is, for that reason, of the heat-tolerant form. In Yemen, the vector is *T. erythrae*, and HLB is only seen in cool highlands. Therefore, Yemeni HLB is of the heat-sensitive form (Bové and Garnier, 1984).

North of the Saudi/Yemeni border, in the Abha-Khamis Mushayt area, both vectors have been found (Bové and Garnier, 1984), and the two forms of HLB are probably present. This situation can be explained in the following way. In Yemen, African HLB and *T. erythrae* have undoubtedly been introduced from nearby Ethiopia, across the Southern Red Sea, through the narrow "Bab al Mandab" strait, and have moved northwards. In Saudi Arabia, Asian HLB and *D. citri* have entered the Arabian Peninsula, probably with pilgrims to Mecca, and have moved southwards. Eventually, the two vectors, as well as the African and Asian HLB, have met in the Abha-Khamis Mushayt region. Further movement of *T. erythrae* up North will be hindered by hot climate, but movement of *D. citri* southwards towards Yemen would not be impaired by climatic conditions.

Indian Ocean islands: Reunion and Mauritius. The two vectors and the two forms of HLB occur in Reunion and Mauritius (Garnier *et al.*, 1996). Distribution of the two psyllid vectors according to temperature/altitude is remarkable. *D. citri* is present on the coastal areas up to an altitude of ~500 m., while *T. erythrae* is prevalent above this altitude. *D. citri* has also been detected in Rodrigues island, East of Mauritius (J. Bové and M. Garnier, unpublished). Both *Ca. L. africanus* and *Ca. L. asiaticus* are present. Some trees proved to be simultaneously infected with the two liberibacters, a situation also observed in São Paulo State with *Ca. L. americanus* and *Ca. L. asiaticus*.

These islands are crossroads, and have been populated by immigrants from both Africa and India. In Reunion, French, African, Malagasy, Chinese, Pakistani and Indian ethnic groups are present. In Mauritius, Indo-Mauritians represent 68%, Creoles, 27%, and Sino-Mauritians, 3%. African HLB has probably been introduced from Eastern Africa and/or Madagascar, and Asian HLB, from the Indian subcontinent.

South America. SPS in Brazil has been the first American region to report HLB, in 2004. The vector is *D. citri*, present in SPS for almost 70 years. Two liberibacters occur. The major HLB agent is the new liberibacter species, *Ca. L. americanus*, present in 92% of trees. *Ca. L. asiaticus* affects 6% of trees, and the two liberibacters occur in 2% of trees (see above).

North America. Florida, U.S.A., has been the second American region to report HLB, in 2005. *D. citri* was recorded 7 years earlier. The liberibacter is *Ca. L. asiaticus*.

CITRUS REGIONS WITH PSYLLID VECTORS BUT NO HLB

The examples of SPS and especially Florida show that once a psyllid vector is present, HLB is not very far. Therefore, the following regions and countries must be seriously on the alert, and scout for HLB, with the hope to catch it early enough to try eradication. In SPS and Florida, several years probably elapsed between the moment the disease was correctly recognized and properly diagnosed, and its first occurrence.

Atlantic ocean islands: Saint Helena, Madeira, and Tenerife. The three islands harbor *T. erythrae*. The insect was reported in 1994 from Madeira, and in 2002 from Tenerife (Canary islands). Madeira is located North of Tenerife, both islands being West of Morocco. It is believed that the African citrus psyllid was introduced from Madeira into the Canary island by the dominant North-South trade winds (Gonzales Hernandez, 2003).

Middle East: Iran. A very small number of *D. citri* psyllids were seen in December 1997 in Iran, close to the border with Pakistan (Bové *et al.*, 2000a). By 2005, large populations of the insect occurred in the major line [*Citrus aurantifolia* (Christm.) Swing] growing region, North of Bandar Abbas (Alimorad Sarafrazi, personal communication).

Americas. Halbert and Nuñez (2004) have described the distribution of *D. citri* in South, Central and North America, and the Caribbean basin.

South America: Brazil, Argentina, Venezuela. In Brazil, as of November 2005, 79 municipalities in SPS and 1 adjacent municipality in Minas Gerais are contaminated with HLB, the vector being *D. citri* for *Ca. L. asiaticus*, and probably also for *Ca. L. americanus*. Other Brazilian regions grow citrus, but they do not (yet) have HLB, even though *D. citri* is present. These regions are at great danger of becoming contaminated with HLB. In 1997, *D. citri* was found on citrus in Corrientes, Argentina. The infestation was minor, the populations being probably controlled by local natural enemies. *D. citri* was reported for the first time in Venezuela in 1999 in the "Península de Paraguana" on *C. aurantifolia*, *C. reticulata*, *C. latifolia*, and *M. paniculata*.

Central America, West Indies, Caribbeans. *D. citri* was found in Guadeloupe, Bahamas, Cayman islands, Cuba and Dominican Republic, and Puerto Rico in 1998, 1999, 2000, 2001, and 2002, respectively (Halbert and Nuñez, 2004). *D. citri* was intercepted on citrus from Belize in a baggage at Houston, Texas, in October 2002.

North America: Texas, Yucatan. *D. citri* is present in



Fig. 6 Sweet orange trees in young high density orchard (1500 trees per hectare) showing convergence symptoms of "yellow shoots" in the Guilin-Yangshuo region, Guangxi, China. Because of HLB, trees do not live longer than 10 years. High tree densities compensate for short orchard life.

the Rio Grande Valley of Texas, U.S.A. (French *et al.*, 2001). It was accidentally introduced in the spring of 2001 on potted *Moraya* plants from Florida. *D. citri* was reported in April 2002 in Cancun, Mexico, and samples were collected in November 2003 (Halbert and Nunez, 2004).

CITRUS REGIONS FREE OF PSYLLIDS AND HLB

The Mediterranean basin, most of Western Asia (Near and Middle East), Australia, and North and South Pacific islands are still free of HLB as well as of the psyllid vectors of the disease. However, for these regions, HLB is a serious threat. The Mediterranean basin could become contaminated with *T. erythrae*, well established in the Portuguese Madeira island and the Spanish Canary islands, not very far from the Moroccan coast. It must be remembered that the most efficient aphid vector of citrus tristena, *Toxoptera citricola*, reported in Madeira island together with *T. erythrae* in 1994, is now present in Northern Portugal and Spain. It is however not firmly established that Madeira was the source of contamination of the Iberian Peninsula with the tristena vector).

The Near and Middle East regions are endangered by *D. citri*, which is well established in South-eastern Iran and in southern Saudi Arabia along the Red Sea. *T. erythrae* is in Yemen. Both Saudi Arabia and Yemen, but not yet Iran, host HLB in addition to the psyllid vectors.

Australia has two northern neighbours contaminated with both *D. citri* and HLB: East Timor and Papua New Guinea.

The North and South Pacific islands have not reported HLB nor psyllid vectors. They must be aware of the situation in Papua New Guinea. During a recent survey for HLB in French Polynesia, no evidence of its presence was found (Bové and Dutac, *Vdo*, unpublished observations). Similarly, no evidence was found for the presence of HLB in four Pacific island countries (Cook islands, Fiji islands, Samoa and Tonga) (Davis *et al.*, 2005b).

ORIGIN OF LIBRIBACTER

Hitherto, *C. L. africanus* has only been found in Africa, and *C. L. asiaticus*, only in Asia (Garner and Bové, 1996). This is probably not due to the fact that the psyllid vector *T. erythrae* is only present in Africa, and the psyllid vector, *D. citri*, only in Asia, because both insects have the ability to transmit both liberibacters, at least under experimental greenhouse conditions (Massoné *et al.*, 1976; Lallemand *et al.*, 1986). Therefore, it seems as if *C. L. africanus* originated in Africa and *C. L. asiaticus* in Asia. After having spread initially

within these continents, disease and vector have spread later to neighbouring regions. In the SPS in Brazil, the situation is more complex. The Asian psyllid, *Diuraphis citri*, has been present, undisturbed, since the 1940s. In these early years, the insect was probably free of an HLB agent, as HLB occurs in SPS only since about 1990. The presence, in SPS, of *C. L. asiaticus*, a very minor component of Brazilian HLB, is probably the result of a relatively recent, uncontrolled introduction from Asia.

It is suggested above, *C. L. africanus* and *C. L. asiaticus* have originated in their respective continents; the same could apply to *C. L. americanus*, supposed to having arisen in SPS. *C. L. americanus* is not a recent "mutation" of *C. L. asiaticus*, because the sequence differences in the 16S rDNA and the RFL of the two liberibacters are far too great, and because there is very little sequence variation between isolates of *C. L. asiaticus*, including the SPS isolate. The sequence of the 16S rDNA of *C. L. americanus* seems to be quite stable too.

The hypothesis proposing that different liberibacter species have evolved in different continents implies that *C. L. americanus* is native to SPS, and has not been introduced from known HLB-affected regions. The countless PCR assays made in Africa and Asia for detection of *C. L. asiaticus* and *C. L. africanus*, have given no indications of the occurrence, in these regions, of a liberibacter different of *C. L. asiaticus* and *C. L. africanus*. Several hundreds of PCR reactions have been carried out in Africa and Asia, and have always detected either *C. L. asiaticus* or *C. L. africanus*. However, the number of these PCR assays might not have been sufficiently large to draw conclusions. Hence, in order to confirm or infirm the above hypothesis, search for *C. L. americanus*, in Asia in particular, should be continued and intensified, especially since PCR tools for specific detection of the various liberibacters are available.

It has recently been suggested that the liberibacters have Gondwanan origins (Deantje *et al.*, 2005). Indeed, until 160 million years ago, Australia, Africa, South America, Madagascar, India, and Antarctica were part of the single "super continent" Gondwana. Speciation of the putative Gondwanan liberibacter ancestor could have occurred after dislocation and fragmentation of the super continent, isolating the African liberibacter lineage within Africa, the American lineage within South America, and the Asian lineage within India, with subsequent eastward spread to China. A similar hypothesis might account for the distribution of the *Diuraphis*. However, if so, it is difficult to explain why *D. citri* is apparently not native to South America, and seems to have been introduced recently. In any case, the Gondwanan hypothesis opens an interesting avenue for academic and applied research.

RECOMMENDATIONS FOR HLB IDENTIFICATION

These recommendations apply to small HLB outbreaks, with eradication of the disease as the objective, as well as to situations where HLB is established, the goal being to live with the disease. They are meant to facilitate quick diagnosis and confirmation of HLB.

Search and monitoring for citrus psyllid vectors. McElhen and Manjunath (2004) have recently reviewed the literature on *D. citri* (Casting (1969a, b, c; 1970), b) having ruled out *protonotaria* wasps on *T. citrifolia*.

The psyllids can feed on many citrus species, but preferred hosts of *D. citri* are *M. pumila* (Orange, tangelo), *M. exoniata* (Citrus hock), and *C. aurantiifolia*, followed by *Bergenia koenigi* (Curry leaf), and *Clusia* spp. These plants should be favoured for monitoring the insects. Lemon (*Citrus limon*) is a good host of *T. erythrae*. Psyllids feed young, actively-growing foliage (fish) for development, and their populations peak at flush periods. In southern Florida, peaks of *D. citri* occur in May, August, and October through December, and this coincides with the new flush growth of *M. pumilata*. This citrus relative has a more continuous flushing pattern, and serves as an alternative host when citrus is not in flush. Psyllid numbers go down after heavy rains. In regions with alternating dry and humid seasons, psyllids will peak at the end of the first growth flush of the new wet season following dry weather (Dav's *et al.*, 2005a).

Eggs of *T. erythrae* are seen mostly along edges of young leaves (Fig. 5 I). Those of *D. citri* occur on tips of growing shoots, or just between unfolding leaves. Similarly, *T. citrifolia* nymphs develop on the underside of leaflets, while *D. citri* nymphs are found on young stems and petioles. The major difference between the two psyllids concerns precisely the nymphs. The *T. erythrae* nymphs are encased in pits or rows on the underside of the leaves (Fig. 5 F) which look like worms on the up-side (Fig. 5 G, H). After an adult has emerged from a nymph, the pest is empty, but the bump remains. Hence, the presence of even one single bump on one single leaf is proof of *T. erythrae* occurrence. No such bumps are produced by *D. citri*. Leaves with humps may be severely distorted. With *D. citri*, leaves are heavily curled (Fig. 5 D), and may be covered with honeydew. Characteristically, *D. citri* adults form a 45° angle with the petiole on which they feed (Fig. 5 A, B, C).

Symptoms of HLB. It is generally thought that it is difficult to diagnose HLB on the basis of symptoms, as none of them is specific, and citrus trees are often affected by different problems. However, some symptoms are highly characteristic, and these will be emphasized here. They apply to most species of citrus, since nearly all citrus seedling trees and scion rootstock combinations are sensitive.

Yellow shoots and blotchy mottled leaves. A tree, which begins to show symptoms in an orchard in early stages of infection, can be identified, among still many symptomless neighbouring trees, by the presence of one or several yellow shoots. These shoots stand out against the green canopy of the tree (Fig. 6). These conspicuous 'yellow shoots', or 'huang long' in Chinese, gave the disease its name. A 'yellow' shoot is seen as resulting from a shoot where psyllids have made a successful infection, as, indeed, HLB bacterium can easily be detected in the symptomatic leaves of these shoots. A yellow shoot is thus an early symptom of the disease. On some trees, only one yellow shoot is present, and, with time, this affected shoot grows into a larger yellow branch, or 'yellow dragon', since 'long' means also 'dragon'. This yellow shoot or yellow branch is a very characteristic aspect of HLB. Often, more than one yellow dragon 'wrap' themselves up around a tree. Eventually, in late stages of the disease, the yellow branches have taken over and canopy the whole tree, which has now become totally infested.

Under cool weather, new flushes may be pale green/yellowish, and zinc deficiency also induces yellow flushes. In these cases, practically all trees in an orchard show the problem. With HLB, however, only some trees carry symptomatic shoots.

The yellow shoots on HLB affected trees look yellow because their leaves are partly yellow, partly green, with several shades of yellow, pale green and dark green, blending into each other, with no sharp limits between the various shades of colour (Fig. 7, 8). Such leaves show what is called 'blotchy mottle'. This symptom was described in particular by Liu (1956) in China (Fig. 7 A), and McElhen and S. Swane (1970) in South Africa (Fig. 7 B, C). It is the most characteristic symptom of HLB wherever the disease occurs, be it in Asia, Africa or America, and wherever the citrus species affected. Also, as seen on Fig. 7 and 8, the blotchy mottle pattern on one side of the leaf midrib is not symmetrical to that on the other side. This asymmetry distinguishes blotchy mottle from zinc (Zn), manganese, magnesium, calcium, and iron deficiency symptoms. With time, the whole leaf blade may arbitrarily turn uniformly yellow (in addition to blotchy mottle and yellowing), the leaves may become thicker, and leathery. Midrib and lateral veins are sometimes enlarged, swollen, and curly. In early stages of infection, blotchy mottle might be the only leaf pattern to be seen. In later stages, foliar symptoms of zinc deficiency will eventually develop, and such leaves have in general an upright growth, making a close angle with the shoot on which they are attached. Eventually, defoliation and dieback occur. With trees in early stages of disease, as observed in SDS, blotchy mottle is seen to affect large, well developed leaves. The same trees, one year later, had produced new mottled leaves, but smaller in size than those from the previous year.




Fig. 7. Blotchy mottle-affected sweet orange leaves from China (A), South Africa (B, C), São Paulo State, Brazil (E), and Madagascar (F). Healthy sweet orange leaves in D.




Fig. 8. Blotchy mottle-affected leaves of sweet orange from São Paulo State (A), rough lemon (B) and grapefruit (C) from Poona India, clementine from South Africa (F), and mandarin from Coorg, India (G, H). Healthy clementine leaves (E). Sweet orange leaves showing zinc deficiency symptoms with increasing severity from right to left (D).

In certain situations, blotchy mottle is difficult to find. In 1996 in North Bali, Indonesia, HLB was severely affecting almost 100% of young, 4 years old Tejakula mandarin trees in many orchards of the coastal plain. The trees were uniformly yellow. Some leaves showed zinc deficiency symptoms, but leaves with clear cut blotchy mottle were rare. These trees had practically no new flushes. PCR tests did however confirm HLB as the cause of decline in all these orchards. At the same time in some orchards at higher altitudes, HLB was just starting, yellow shoots occurred, and leaves had strong blotchy mottle.

Blotchy mottle is probably best observed on sweet orange trees, but many citrus species and varieties show it, including some mandarin varieties such as Tejakula in Bali and Clementine in South Africa (Fig. 9 F). Other mandarin varieties, such as the Coorg mandarin in India, do not present good, classic blotchy mottle. The colour patches of yellow, stark green and pale green are not as large as on sweet orange leaves, and rather produce a mosaic pattern (Fig. 8 G, H). Also, in the orchard, affected mandarin leaves tend to be more curled than sweet orange leaves (Fig. 8 G). However, in the greenhouse, leaves of affected sweet orange trees are frequently curly.

Fruit symptoms. HLB induces also very characteristic fruit symptoms – seen especially well on sweet oranges and mandarins, and sometimes pumelos. Again, these symptoms are non-specific, as similar symptoms are associated with stubborn disease. Fruits with HLB symptoms should be looked for as yellow shades on branches. Symptomatic fruits are small, asymmetric, lopsided, with a bent fruit axis. At the time when the fruit changes colour, the top (green to yellow/orange). HLB-affected fruit shows colour inversion: the peduncular end of the fruit turns yellow/orange, while the styler end is still green, whereas on normal fruit the coloration starts first at the styler end, and moves only later to the peduncular end (Fig. 9 A, D, F to J). In addition, when the peduncle of a fruit with colour inversion is carefully removed, the resulting circular scar is stained orange, while on a normal fruit the scar is pale green. Colour inversion can be very pronounced in China, where it is called ‘red nose’. In South Africa, HLB affected fruit colours also unevenly (Fig. 9 C), and this symptom was responsible for calling the disease ‘greening’. Sometimes, when one prunes such fruit with the thumb, a silver ‘lieger mark’ results (Fig. 9 E).

To further observe the symptoms, fruits should be cut in half. When the sectioning is made perpendicular to the fruit axis, it is easy to see the small, brownish/black aborted seeds, highly characteristic of HLB, but also present in stubborn affected sweet oranges. Cutting a lopsided fruit through the fruit axis, reveals its asymmetry, and some aborted seeds can also be seen (Fig. 10

A, B, C). In addition, the vascular bundles within the fruit axis at the peduncular end have a strong brownish stain (Fig. 10 B). As in stubborn affected fruit, the albedo is sometimes thicker at the peduncular end than at the styler end (Fig. 10 B).

In summary, when a tree shows one or several yellow shoots with blotchy mottled leaves, and small, lopsided fruits with colour inversion and aborted seeds, it is highly likely that the tree is affected by HLB. To confirm the HLB diagnosis, PCR detection of the HLB liberibacters should be made.

PCR confirmation of HLB diagnosis. For PCR detection of the HLB liberibacters blotchy mottled leaves should be used. Such leaves almost never fail to give strong, positive PCR reactions when the symptoms are due to HLB. Therefore it is untrue to state, as it is sometimes written, that it is difficult to detect liberibacters. For detection of *Ca. L. africanus* and *Ca. L. asiaticus* two sets of primers can be used, primers f-OA1 and r-ODc for amplification of 16S rDNA, and primers f-rplA2 and r-rplJ5 for amplification of ribosomal protein genes (rpl). In the case that the presence of both African and Asian liberibacters is suspected, the primer pair f-(OA1-OH) / r-ODc is recommended. *Ca. L. americanus* is detected with PCR primers f-GB1 and r-GB3. Simultaneous detection of the three liberibacters by multiplex PCR has been developed with the following two primer pairs in the same reaction mixture: (i) f-rplA2 / r-rplJ5 and (ii) f-GB1 / r-GB3.

As mentioned above, initially in Brazil, most PCR reactions gave negative results, even though leaves with strong symptoms of blotchy mottle were used. This was later shown to be due to the fact that a new liberibacter species, *Ca. L. americanus*, was involved (Teixeira *et al.*, 2005a, b, c). If a similar situation is encountered elsewhere, search for a new liberibacter should be sought.

In some cases, when HLB symptoms are not immediately recognized, and are attributed to some other causes, such as mineral deficiencies, the relevant trees might be in relatively late stages of the disease, and mottled leaves may be lacking. In such cases, leaves with zinc-deficiency symptoms should be used.

For PCR detection, HLB-affected leaves, infected with either one of the three liberibacters, should be available as positive controls. If not, DNA preparations from such leaves, obtainable from several laboratories, can be used. Healthy citrus leaves are used as negative controls.

CONTROL OF HLB

There are no curative methods to control HLB. When HLB was found to be a bacterium, rather than a viral disease, antibiotic control by trunk injections of streptomycin

cyclines was tried in several countries, including South Africa, Indonesia, and Taiwan, but soon abandoned, because of ecological reasons, but essentially because tetracycline is bacteriostatic, rather than bactericidal, and the treatment had to be repeated each year. Thus, control is by preventing trees from becoming infected. If HLB is not yet present, quarantine measures should be enforced to keep it out. The most serious situation is when HLB has entered a region previously HLB-free. The first immediate measure is to survey and determine the extent of the outbreak. The next objective is to prevent as many trees as possible from becoming infected. This can only be done (i) by eliminating as much as possible liberibacter inoculum, i.e. by removing infected trees, and (ii) by keeping psyllid populations as low as possible. Without these control measures, HLB evolution in an affected orchard is fast. In China, it has been stated repeatedly that, without control, it takes only about five years for an orchard to reach 100% infection. Similar situations seem to occur in SPS. In the Reunion island, the time needed to reach 100% infection was estimated to be 15 years. Average time is eight years.

Quarantine measures. If HLB is still absent from a given region, do not introduce it. The biggest quarantine threat is transport, by people, of citrus and citrus relatives. These plants may be infected with liberibacters and may carry psyllid eggs and/or nymphs. *M. paniculata*, an ornamental citrus relative, is a preferred host of *D. citri*, and if further work confirms that it is also a host of the liberibacters, it deserves special quarantine attention. Also, it must be remembered that the liberibacters cannot be detected in symptomless leaves, even though their detection in mottled leaves is straightforward. Symptomless leaves are either healthy, and will never give positive PCR reactions, or they are already infected, but the liberibacter titer is still too low to be detected by the PCR methods available today. For these reasons it is meaningless to use conventional HLB PCR techniques with symptomless plant material.

Eliminating inoculum by removal of infected trees. If all infected trees were symptomatic, it would be easy to spot them, on the basis of their symptoms, and remove them. Unfortunately, there is a latency period during which recently infected trees do not show symptoms. This period extends from the moment the tree becomes infected by the psyllids up to the moment it expresses symptoms. The length of this period may vary from tree to tree, but is generally assumed to be 6 to 12 months long, if not longer. Therefore, removal of all symptomatic trees will not result in removal of all infected trees. In practice, several surveys are required to remove as many infected trees as possible. Some infected trees, which were still symptomless at survey # n , will show symptoms at survey # $n+1$, and can be removed;

other trees will only be removed at survey # $n+2$. Also, within the time period between surveys n and $n+1$, or $n+1$ and $n+2$, psyllids (if not well controlled), may infect additional trees, and so on. Obviously, the time period between successive surveys is important. In particular, if this period is too long, some symptomatic trees will have become severely affected before they are removed, and serve as new sources of inoculum. In SPS, experience seems to show that a period of about three months is a good compromise.

Surveys must be carried out carefully. All trees in an orchard, however large it is, should be examined one by one, and enough time, say a few minutes, must be spent at each tree. Experience gained in a large grove (one million trees) in southern China suggests that the scouts in charge of the surveys should work in pairs, so that each tree is examined by the two scouts of a pair, one scout on each side of the row. In the case of orchards with large, adult trees, it is essential to examine the top of the trees. In SPS, relatively high towers have been built onto tractors to permit efficient observation of tree tops. Finally, once affected trees have been identified, they should be removed as quickly as possible.

In SPS, recommendation is that all symptomatic trees be removed, whatever their age or the severity of their infection (Ayres *et al.*, 2005). Removing HLB-affected branches by pruning, instead of removing the whole tree, has been discarded because the many sprouts that develop at the site of pruning turn into 'yellow shoots' and favour psyllid development. In South Africa, it has been recommended (Butterdag and von Broembsen, 1993) that HLB-affected trees up to 5 years of age be removed. Trees from 6 to 10 year-old, when severely affected, are also removed. With less affected trees, and with trees older than 10 years, only the affected branches are removed.

In SPS, citrus canker is another disease of citrus, the control of which also involves removal of trees. In this case, the infected tree as well as all surrounding trees, within a distance of 15 m, are removed. From the epidemiology studies of HLB (Gerswald, 2005; Bassarezi, 2005), it seems that this strategy is probably inadequate for HLB. Indeed, when HLB vectors move, either naturally in search of new feeding opportunities or when disturbed, they not only move to neighbouring trees (primary foci), but also to trees located at distances of 25 to 50 m (secondary foci). In addition, psyllids can move a distance of at least 1.5 km with wind, and probably much longer distances with hurricanes or tropical storms, such as those in Florida.

When an orchard is too severely affected, with a high percentage of affected trees, it might be best to remove the whole stand. The following figures might help to make the decision. It can be roughly estimated that with Asian HLB, *D. citri* being the vector, an orchard with 10% symptomatic (Sy) trees, has a total of 20% infecti-




Fig. 9. The two most characteristic symptoms of HLB are leaves with blotchy mottle and fruits with small size and colour inversion (A, normal sweet orange on right; Beihai, Guangxi, China). Poorly coloured "greening"-affected oranges from South Africa (C), which, when pressed with a finger, develop a silvery "finger mark" (E). Normal oranges with coloration starting at the stylar end (B). Oranges and mandarins with coloration starting at the peduncular end (colour inversion) (i) sweet oranges from China, Guangxi (D) and Fujian (H, normal fruits with orange stylar ends on far right, and green peduncular ends on far left; fruits with colour inversion, in the middle), and São Paulo State, Brasil (G, I); (ii) mandarins from North Bali (F), and China (J, lower fruit: normal, upper fruit: with colour inversion).



Fig. 10. A. Far left and far right, normal sweet orange halves showing normal seeds; middle, two halves from a lopsided HLB-affected sweet orange from China. Notice asymmetry of the fruit section, and the brownish-black seeds. B. Sectioned lopsided sweet orange from São Paulo State, with brownish-black seeds, thick albedo at the peduncular end, and vascular bundles stained orange-brown at the peduncular end of fruit axis. C. Lopsided pummelo showing vascular bundles stained orange-brown.

ed trees, i.e., symptomatic trees plus asymptomatic trees (Sy+Asy). For 20%, 30% and 50% Sy trees, there will be 36%, 50%, and 70% (Sy+Asy) trees. In other words, in an orchard with 30% symptomatic trees, half of the trees are infected, and will have to be pulled out sooner or later. Is it worthwhile to keep such an orchard? Probably not!

When affected trees have been pulled out, they can be replaced by young, healthy trees. In SPS, by law, all nursery trees have to be produced in covered, insect proof screenhouses, and this obligation guarantees that they are HLB-free.

Even though in SPS removal of HLB-affected trees is compulsory by law, great efforts have been made through the media (articles in newspapers, programs on radio and TV, training courses for growers and technicians) to explain the situation, and convince the growers to remove the affected trees on a voluntary base (Ayres

et al., 2005). Most of the growers cooperate, and 350 thousand symptomatic trees have been voluntarily eliminated, as of October 2005. Cooperation of the growers is also required for vector control.

Psyllid vector control. Key features of the psyllids, which need to be taken into account in the formulation of control strategies have been reviewed (Buitendag and von Broembsen, 1993; Halbert and Manjunath, 2004; Hall, 2005).

Treatments with contact and systemic insecticides reduce psyllid populations, and thus HLB spread. The efficiency of the treatments is easier to estimate with *T. erytreae* than with *D. citri*, because of the foliar "bumps" produced by *T. erytreae* nymphs. In the large, South African citrus estates, control of *T. erytreae* can be very efficient, as "bumps" are hardly seen. Systemic insecticides, aimed specifically at psyllids (monocro-

tophos), have been applied directly to the tree trunks, and have been very effective. In Asia, a large range of insecticides, mostly organophosphates and pyrethroids, are used in very intensive spray programs to kill eggs and nymphs on flush growth. Horticultural and agricultural mineral oils are being developed as alternative treatments, and these are much less damaging to the environment and less disruptive to biocontrol of other pests (Davis *et al.*, 2005a).

Reunion Island is the only region where biological control of both *T. erythrae* and *D. citri* has been effective (Lacenne and Aubert, 1980; Aubert *et al.*, 1980). The success is due to the fact that, initially, the two psyllids occurred in the island in the absence of parasites, and thus, hyperparasites. Therefore parasites of the two psyllids were introduced into the island, care being taken not to introduce hyperparasites at the same time. *Tamarixia dryi*, a parasite of *T. erythrae*, and *Tamarixia radiata*, a parasite of *D. citri*, were collected, respectively in South Africa and India, reared in Reunion and released in abandoned citrus orchard having received no insecticide treatments for quite some time. A second reason for success is the fact that *T. dryi* multiplied not only on *T. erythrae*, but also on an additional psyllid, *Trioza castaneipes*, not a citrus psyllid, but multiplying on a widely distributed weed, *Litsea chinensis*. Releases of *Phyllaephagus pulvinatus*, introduced from South Africa, were also made, but the insect, much less vigorous than *T. dryi*, disappeared from the island.

In Florida, *T. radiata* has been introduced from Taiwan and Vietnam, and is being established in Florida. In China, studies are conducted on *Carapellum chinensis* Schmitz, as a potential biocontrol predator of *D. citri* (Wu, 2005).

The Brazilian experience. In SPS, encouraging results have been obtained in the control of HLB by tree removal and insecticide treatments against psyllids (Ayres *et al.*, 2005). For instance, in one large citrus orchard (71,000 trees), 10 surveys for identification of HLB-affected trees were carried out from September 2004 until September 2005. The first survey identified 7% affected trees. On the 2nd survey, in December 2004, the percentage went down to 0.01%, but increased again to 1.1% after the 6th survey in May 2005. From there on, the percentages decreased regularly to 0.5% (7th survey, June 2005), 0.15% (8th survey, July 2005), 0.05% (9th survey, August 2005), and 0.03% (10th survey, September 2005). It is hoped that the percentages of affected trees will remain at the level of 0.05%, even with less frequent surveys and tree removals. HLB will not be eradicated, but it will be possible to live with the disease.

To achieve this goal, SPS has a number of favourable factors. Brazil, contrary to Florida, is not affected, year after year, by hurricanes and tropical storms, the role of

which in spreading the disease and its vector, might be more significant than thought. A task force of 800 people from both the private sector (Fruitecitra) and the public sector (Government, public research organizations) is involved in HLB control and helps the growers. A good interface has been built between growers, Fruitecitra and the government. Sixty million trees have already been surveyed, 70 million more trees have still to be inspected. The first wide range survey will be completed in February 2006. The surveys for HLB benefit from the year long experience of Fruitecitra as the control of citrus canker. The nursery trees required to replace the removed HLB-affected trees are produced from healthy budwood in covered, insect proof nurseries, a procedure that guarantees that they are free of HLB and all other graft-transmissible diseases (except CTV strains used for cross protection). Even though HLB has been present for several years before it was recognized, most orchards in the affected municipalities have still a low disease incidence.

Very few countries have been able to control Asian HLB. SPS might be one of the first to be successful, and live with the disease until genetically engineered citrus cultivars, resistant to HLB, will eventually become available (Liu *et al.*, 2005).

ACKNOWLEDGMENTS

This review has benefited from my recent visits to Brazil, China, and Florida to discuss yellow fever disease and be involved in control strategies. I am very grateful to my colleagues from Fruitecitra and the many private citrus estates in São Paulo State for sharing their information with me, and making it possible to produce new results. China is afflicted with Huanglongbing since more than a century, and we have taken great advantage of the experience provided by our colleagues from Guangxi, Guangdong, Fujian, and Sichuan provinces. In Florida, the dragon is emerging, and a most interesting workshop was held recently to exchange information and put the beast on the leash. I am indebted to André Wertheim for having provided me with rare reprints and for interesting discussions over the net. Finally, I thank my laboratory colleagues for their years-long indulgence to me.

REFERENCES

- Ayres P. S. 2005. Estudos indicativos de controle de psilídeos e da doença amarelo-verde em laranjeiras. <http://www.fruitecitra.com.br>
- Aubert R. D. 1958. The citrus yellow-bottle fly, *D. citri*, and the collection of citrus fruits in India. *Indian Journal of Horticulture* 13: 285-286.

- Aubert B., Bové J.M., 1980. Effect of penicillin or tetracycline injections of citrus trees affected by greening disease under field conditions in Reunion island. *Proceedings of 8th Conference of IOCV, IOCV, Riverside* 1980, 103-108.
- Aubert B., Bové J.M., Etienne J., 1980. La lutte contre la maladie du «greening» des agrumes à l'île de la Réunion. Résultats et perspectives. *Fruits* 35, 605-624.
- Aubert B., Garnier M., Guillaumin D., Herbogardeno B., Scribani L., Nuzhadi E., 1985. Le greening, une sérieuse menace pour les productions agrumicoles de l'archipel indonésien. *Fruits* 40, 549-551.
- Aubert B., Garnier M., Cassin J.C., Bertia Y., 1986. Citrus greening disease (or-yu) in East and West African countries South of Sahara. *Proceedings of 10th Conference of IOCV, IOCV, Riverside* : 188, 211-237.
- Ayres A.J., Massad C.A., Lopes S.A., Bastaneni R.B., Belasque Junior J., Yamamoto P.T., Teixeira D.C., Wolff N.A., Guitierrez-Fernandes N., Bergamaschi O.A., 2005. *Proceedings of 2nd International Citrus Canker and Huanglongbing Research Workshop, Florida Citrus Mutual, Orlando* 2005, H-4.
- Bastaneni R.B., Busato L.A., Bergamin Filho A., Amorim L., Gotovald T.R., 2005. Spatial distribution of huanglongbing symptomatic trees in citrus groves in São Paulo, Brasil. *Proceeding, 2nd International Citrus Canker and Huanglongbing Research Workshop, Florida Citrus Mutual, Orlando* 2005, H-7.
- Bastaneni R.B., Garnier Semarick M., Renaudin J., Bové J.M., Everlard S., 2005. Diversity of 'Candidatus Liberibacter asiaticus' based on the *omp* gene sequence. *Applied Environmental Microbiology* 71: 6473-6478.
- Beattie G.A.C., Mabblerley D.J., Holford P., Broadbent P., De Bono P., 2005. Huanglongbing: its possible origins, collaborative research in Southeast Asia, and developing incursion management plans for Australia. *Proceedings of 2nd International Citrus Canker and Huanglongbing Research Workshop, Florida Citrus Mutual, Orlando* 2005, H-5.
- Bonavia E., 1888. The cultivated oranges and lemons of India and Ceylon. W.H. Allen & Co., London.
- Bové J.M., Garnier M., 1984. Citrus greening and psylla vectors of the disease in the Arabian Peninsula. *Proceedings of 8th Conference IOCV, IOCV, Riverside* 1984, 109-114.
- Bové J.M., Saglio P., 1964. Stubborn and Greening: a review, 1969-1972. *Proceedings of 6th Conference IOCV, IOCV, Riverside* 1974, 1-11.
- Bové J.M., Calavan E.C., Kapoor S.P., Coster R.F., Schwarz R.E., 1974. Influence of temperature on symptoms of California stubborn, South African greening, Indian citrus decline and Philippines leaf mottling diseases. *Proceedings of 6th Conference IOCV, IOCV, Riverside* 1974, 12-15.
- Bové J.M., Bonnet P., Garnier M., Aubert B., 1980. Penicillin and tetracycline treatments of greening disease affected citrus plants in the glasshouse, and the bacterial nature of the procytote associated with greening. *Proceedings of 8th Conference IOCV, IOCV, Riverside* 1980, 91-102.
- Bové J.M., Garnier M., Al-Jawar Y.S., Chakraborty N.K., Morita A., 1995. Detection of the Asian strains of the greening BLO by DNA-DNA hybridisation in Indian orchard trees and Malaysian *Diaphorina citri* psyllids. *Proceedings of 12th Conference IOCV, IOCV, Riverside* 1995, 258-263.
- Boer J.M., Chou N.M., Ha Minh Trung, Bourdeaut J., Garnier M., 1996. Huanglongbing (greening) in Vietnam: detection of *Liberibacter asiaticus* by DNA hybridisation with probe in 2.6 and PCR amplification of 16S ribosomal DNA. *Proceedings of 13th Conference IOCV, IOCV, Riverside* 1996, 258-266.
- Bové J.M., Datto J.L., Banarjee K., Hasanmajeed N., Salehi M., Garnier M., 2000a. Witches' brown disease of lime (WB-DL) in Iran. *Proceedings of 14th Conference IOCV, IOCV, Riverside* 2000, 207-212.
- Bové J.M., Fei Dwiastuti M., Trivisnanto A., Supriyanto A., Nasli H., Bocu P., Garnier M., 2000b. Incidence of huang longbing and citrus rehabilitation in North Bali, Indonesia. *Proceedings of 14th Conference IOCV, IOCV, Riverside* 2000, 209-206.
- Bruton R.D., Fletcher J., Parr S.D., Shaw M., Srinets-Bhatkar H., 1998. Association of a plasmid limited bacterium with yellow vein disease in Cucurbit. *Plant Disease* 82: 512-520.
- Buitendij C.H., von Broembsen A., 1993. Living with citrus greening in South Africa. *Proceedings of 12th Conference IOCV, IOCV, Riverside* 1993, 269-273.
- Capoor S.P., 1963. Decline of citrus trees in India. *Bulletin National Institute of Science India* 24: 48-64.
- Capoor S.P., Rao D.G., Viswanath S.M., 1961. *Diaphorina citri* Kowaly, a vector of the greening disease of citrus in India. *Indian Journal of Agricultural Science* 31: 572-576.
- Carling H.D., 1969a. The bionomics of the South African citrus psylla, *Trioxys erythrae* (Del Guercio) (Homoptera: Psyllidae). I. The influence of the feeding rhythm of citrus and factors which regulate feeding. *Journal of Entomology Society South Africa* 32: 191-208.
- Carling H.D., 1969b. The bionomics of the South African citrus psylla, *Trioxys erythrae* (Del Guercio) (Homoptera: Psyllidae). II. The influence of parasites and never on the species involved. *Journal of Entomology Society South Africa* 32: 209-223.
- Carling H.D., 1969c. The bionomics of the South African citrus psylla, *Trioxys erythrae* (Del Guercio) (Homoptera: Psyllidae). III. The influence of extremes of weather on survival. *Journal of Entomology Society South Africa* 32: 274-296.
- Carling H.D., 1970a. The bionomics of the South African citrus psylla, *Trioxys erythrae* (Del Guercio) (Homoptera: Psyllidae). IV. The influence of predators. *Journal of Entomology Society South Africa* 33: 341-348.
- Carling H.D., 1970b. Distribution of the psyllid vectors of citrus greening disease, with notes on the biology and bionomics of *Diaphorina citri*. *FAO Plant Protection Bulletin* 18: 8-15.
- Carling H.D., M. Garnier, J.M. Bové, 1978. Presence of citrus greening disease in Bangladesh and a new method for rapid diagnosis. *FAO Plant Protection Bulletin* 36: 14-18.
- Chou Chien-shan, 1943. A report of a stubborn disease of citrus in Cansuhar. *Quarterly Bulletin of New Formosa* 1: 142-152.

- Chen M.H., Miyakawa T., Marui C., 1991. Mycoplasma like bodies associated with 'Shuhin-disease' Ponkan (citrus) *Phytopathology* 81: 528.
- Chen M.H., Miyakawa T., Marui C., 1993. Citrus 'Shuhin' pathogens in the salivary glands of *Diosaphis citri* (Homoptera: Pemphigidae) *Entomol. exp. appl.* 74: 145-155.
- Chen Y.Y., Shen C.Y., Tai S.C., Kung T.H., Chen H.Y., Chiang Y.H., Chen C.S., 1979a. Mycoplasma like organisms associated with Huanglongbing of sweet orange (*Citrus sinensis*) *Biophytologia Sinica* 11: 389-391.
- Chen Y.Y., Shen C.Y., Tai S.C., Kung T.H., Chen H.Y., Tai Y.M., 1979b. Studies on the pathogen of Huanglongbing (citrus yellow shoot disease), a Mycoplasma like organism associated with Huanglongbing on Kwantung *Acta Biobotanica et Biophysica Sinica* 11: 191-197.
- Colleta Filho R.D., Targui M.L.P.N., Takai M.A., De Negrí J.D., Pompeu J.J., Machado M.A., 2004. First Report of the causal agent of Huanglongbing (*Candidatus Liberibacter asiaticus*) in Brazil *Plant Disease* 88: 1382.
- Da Graça J.V., 1991. Citrus greening disease *Ann. of Review of Phytopathology* 29: 111-116.
- Da Graça J.V., Kester L., 2001. Citrus Huanglongbing: recent progress status and future strategies. In: S.A.M.L. Nzuji (Ed.) *Diseases of Fruits and Vegetables*, pp. 229-245. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Davis M.J., Kramer J.H., Kowwda H., Brunner E.S., 1999. Association of a bacterium and not a phytoplasma with palm little-leafy top disease. *Phytopathology* 89: 11-19.
- Davis M., Gnanou Y., Bellis G., Grundler L., 2005a. The Huanglongbing disease of citrus trees. *Pest advisory bulletin* 45: 24-25. Plant Protection Services, Secretariat of the Pacific Community ISSN 1171-8236.
- Davis M., Gnanou Y., Easo M., Tejada D., Barber T., 2005b. Spread of citrus Huanglongbing (greening) disease to island ecosystems on into Papua New Guinea. *Australian Plant Pathology* 34: 517-524.
- Du J., Fan Y., Gao J., Wang Guoping, 2004. Research on the ELISA method for detection of citrus Huanglongbing pathogen. *Acta Horticulturae Sinica* 31: 807-811.
- Dong Fang, Yi Gejin, Wang Guoping, Yun Zhang, 2006. Huanglongbing (*Candidatus Liberibacter asiaticus*) infecting watermelon and lemon in China by microscopy, PCR and nested PCR analysis. *Plant Science* 171: 105-114.
- Die Dot, Nungay On, Camille Dot, Pierre Gnanou M., Jacques-Eveillard S., Bové J.M., 2003. First report of *Candidatus Liberibacter asiaticus*, the agent of citrus Huanglongbing (greening), in Réunion. *Plant Disease* 87: 218.
- Doi Y., Terashima M., Yone K., Asayama H., 1997. Mycoplasma-like bodies in citrus greening found in the field specimens of plants infected with naturally derived pathogen with electron microscope, in *Paulownia* infected broom. *Journal of Phytopathology Society Japan* 33: 259-266.
- Fraser J., Aubert B., 1980. Biological control of psyllid vectors of greening disease on Reunion Island. *Proceedings of 8th Conference IOCV, IOCV, Riverside* 1980, 10-121.
- Fraser L.R., Singh D., 1968. Citrus dieback in India - the isolation of greening virus. *Proceedings of 4th Conference IOCV, IOCV, Riverside* 1968, 141-144.
- Fraser L.R., Singh D., Capoor S.P., Nariani T.K., 1966. Greening virus, the likely cause of citrus dieback in India. *Indian Plant Protection Bulletin* 14: 127-130.
- French J.V., Kalike C., da Graça J.V., 2001. Asian psyllid found on Texas citrus. *Citrus Center* 19: 1.
- Gao S., Garnier M., Bové J.M., 1993. Production of monoclonal antibodies recognizing insect strains of the greening BLO by in vitro immunization with an antigenic protein purified from the BLO. *Proceedings of 12th Conference IOCV, IOCV, Riverside* 1993, 244-247.
- Garnier M., Bové J.M., 1997. Structure tridimensionnelle des deux membranes qui entourent les organismes procaryotes associés à la maladie du "greening" des agrumes. *Ineur* 32: 149-152.
- Garnier M., Bové J.M., 1981. Transmission of one organism associated with citrus greening disease from sweet orange to pomelo by dodder. *Phytopathology* 73: 1358-1361.
- Garnier M., Bové J.M., 1993. Citrus greening disease and the greening bacterium. *Proceedings of 12th Conference IOCV, IOCV, Riverside* 1993, 217-219.
- Garnier M., Bové J.M., 1996. Distribution of the Huanglongbing (greening) *Liberibacter* species in fifteen African and Asian countries. *Proceedings of 13th Conference IOCV, IOCV, Riverside* 1996, 388-391.
- Garnier M., Bové J.M., 2000. Huanglongbing in Cambodia, Laos, and Myanmar. *Proceedings of 14th Conference IOCV, IOCV, Riverside* 2000, 378-380.
- Garnier M., Lantier J., Bové J.M., 1976. *Spiroplasma citri* and the organism associated with little-leafy top: comparison of their envelope systems. *Proceedings of 7th Conference IOCV, IOCV, Riverside* 1976, 13-17.
- Garnier M., Darel N., Bové J.M., 1984a. Aetiology of Citrus Greening Disease. *Annals of Microbiology (Gustav Fischer)* 135A: 169-179.
- Garnier M., Darel N., Bové J.M., 1984b. The greening organism is a Gram negative bacterium. *Proceedings of 9th Conference IOCV, IOCV, Riverside* 1984, 113-124.
- Garnier M., Gao S.J., He Y., Villechanoux S., Garnier J., Bové J.M., 1991. Study of the greening organism (GO) with monoclonal antibodies: serological identification, morphology serotypes and purification of the GO. *Proceedings of 11th Conference IOCV, IOCV, Riverside* 1991, 428-435.
- Garnier M., Jaquiez-Eveillard S., Toorawa P., Grisoni M., Mallesard R., Desbren A., Saumtally S., Aubrey J.C., Bové J.M., 1996. Both Huanglongbing (greening) *Liberibacter* species are present in Mauritius and Reunion. *Proceedings of 13th Conference IOCV, IOCV, Riverside* 1996, 392-394.
- Garnier M., Bové J.M., Cronje G.P.R., Sanders G.M., Kovton L., Le Roux H.H., 2000a. Presence of "*Candidatus Liberibacter africanus*" in the Western Cape province of South Africa. *Proceedings of 14th Conference IOCV, IOCV, Riverside* 2000, 369-372.
- Garnier M., Jaquiez-Eveillard S., Cronje P., Le Roux H., Bové J.M., 2000b. Genomic characterization of a *Liberibacter* present in an ornamental Rutaceae tree, *Colodendrum tomentosum*.

- in the Western Cape province of South Africa. Proposal for a "*Candidatus Liberibacter africanus subsp. capensis*". *International Journal of Systematic Evolution Microbiology* 50: 2119-2125.
- Garcinea E., Jacob N., Vauvrie S., Larue J., Lherminier J., Richard-Misard M., Besson-Padon E., 2002. Association with the syndrome «flaves richness» of sugar beet of a phytoplasma and a bacterium-like organism transmitted by a *Pentatomidae* sp. *Phytopathology* 92: 184-192.
- González Hernández A., 2003. *Tristeza citrícola* (Cil Gulchicis 1918): nueva plaga de los cítricos en Canarias. *Phytoma* 153: 112-118.
- Gioach S.C., Giacomini J., Louis C., 1977. Multiplication intensive des prokaryotes associés aux maladies de type "greening" des agrumes dans les cellules tubées de cuscute. *Annals of Phytopathology* 9: 525-530.
- Goodall TR., 2005. Huanglongbing epidemiology: tracking the dragon through time and space. *Proceedings of 2nd International Citrus Canker And Huanglongbing Research Workshop, Florida Citrus Mutual, Orlando 2005*, 11-6.
- Halbert S.E., 1998. Entomology section. *Tristogy*, 37: 6-7.
- Halbert S.E., 2005. The discovery of huanglongbing in Florida. *Proceedings Of 2nd International Citrus Canker And Huanglongbing Research Workshop, Florida Citrus Mutual, Orlando 2005*, 11-3.
- Halbert S.E., Manjunath K.L., 2004. Asian citrus psyllid (*Sternorhyncha: Psyllidae*) and greening disease of citrus: a literature review and assessment of risk in Florida. *Florida Entomologist* 87: 330-353.
- Halbert S.E., Nunez C.A., 2004. Distribution of the Asian citrus psyllid, *Diuraphis citri* Kawayana (Rhopidota: Psyllidae) in the Caribbean basin. *Florida Entomologist* 87: 401-402.
- Halbert S.E., Nisbett C.L., Manjunath K.L., Lee R.F., Brown L.G., 2003. Establishment of two new vectors of citrus pathogens in Florida. In: *Proceedings of IX Congress International Society Citriculture, ASIIS Press, Alexandria*, 1016-1017.
- Hall D.G., 2005. Overview of the Asian citrus psyllid. *Proceedings of 2nd International citrus canker and huanglongbing research workshop, Florida Citrus Mutual, Orlando 2005*, 11-12.
- Hartung J.S., Li W., Levy L.E., 2005a. Comparison of methods for the detection of *Candidatus Liberibacter asiaticus* in plant samples. *Proceedings of 2nd International Citrus Canker And Huanglongbing Research Workshop, Florida Citrus Mutual, Orlando 2005*, 11-9.
- Hartung J.S., Folimonov A.S., Foliminov S.V., Dawson W.O., 2005b. Screening of anti-bacterial peptides in citrus trees for activity against *Candidatus Liberibacter asiaticus*. *Proceedings of 2nd International Citrus Canker And Huanglongbing Research Workshop, Florida Citrus Mutual, Orlando 2005*, 13-16.
- Hequellet A., Bové J.M., Garnier M., 1997. Production and evaluation of new radiolabeled probes for the detection of the two "*Candidatus Liberibacter*" species associated with citrus huanglongbing (greening). *Molecular Cellular Probes* 11: 433-435.
- Hequellet A., Tournan P., Bové J.M., Garnier M., 1999. Detection and identification of the two "*Candidatus Liberibacter* sp." associated with citrus huanglongbing by PCR amplification of ribosomal protein genes of the beta operon. *Molecular Cellular Probes* 13: 375-379.
- Hong T.H., Wu M.L., Su H.J., 2000. Identification of alternative hosts of the fastidious bacterium causing citrus greening disease. *Journal of Phytopathology*, 148: 251.
- Igwagbe F.C.K., Calvert E.C., 1970. Occurrence of mycoplasma-like bodies in phloem of scabion infected citrus seedlings. *Phytopathology* 60: 1525-1526.
- Jagoueix S., Bové J.M., Garnier M., 1994. The phloem limited bacterium of greening disease of citrus is a member of the re-subdivision of the *Proteobacteria*. *International Journal of Systematic Bacteriology* 44: 397-386.
- Jagoueix S., Bové J.M., Garnier M., 1996. PCR detection of the two "*Candidatus*" *Liberibacter* species associated with greening disease of citrus. *Molecular Cellular Probes* 10: 43-50.
- Jagoueix S., Bové J.M., Garnier M., 1997. Comparison of the 16S/23S Ribosomal Intergenic Regions of "*Candidatus Liberibacter asiaticus*" and "*Candidatus Liberibacter africanus*", the two species associated with Citrus Huanglongbing (Greening) Disease. *International Journal of Systematic Bacteriology* 47: 224-227.
- Ke C., Lin X.J., Chen H., Chen X.C., Chang J.J., 1979. Preliminary study on the relation between rickettsia like organism and a filamentous virus citrus yellow shoot disease. *Kuaxue Tongbao* 10: 463-466.
- Korsten L., Jagoueix S., Bové J.M., Garnier M., 1996. Huanglongbing (greening) detection in South Africa. *Proceedings of 13th Conference ICCV, IOCV, Riverside 1996*, 339-338.
- Lafrière D., Bové J.M., 1970a. Structures de type mycoplasme dans les feuilles d'orangers atteints de la maladie du greening. *Comptes Rendus de l'Académie des Sciences Paris*, 270: 1915-1917.
- Lafrière D., Bové J.M., 1970b. Mycoplasmes dans les agrumes atteints de "greening", de "scabion" ou de maladies similaires. *Fruits* 25: 439-440.
- Lafrière D., Fus A., Bové J.M., 1986. Transmission de la bactérie associée à la forme africaine de la maladie du greening par le psylle asiatique *Diuraphis citri* Kawayana. *Fruits* 41: 341-345.
- Lee H.A., 1921. The relation of stocks to mottled leaf of citrus leaves. *Philippine Journal of Science* 18: 85-95.
- Li T., Ke C., 2002. Detection of bearing rate of *Liberibacter asiaticus* in citrus psylla and its host plant *Murraya paniculata* by nested PCR. *Acta Phytopathologica Sinica* 29: 31-37.
- Li W., Hartung J.S., Levy L., 2005. Development of multiplex real-time PCR for detection and identification of *Candidatus Liberibacter* species associated with citrus huanglongbing. *Proceedings of 2nd International Citrus Canker And Huanglongbing Research Workshop, Florida Citrus Mutual, Orlando 2005*, 11-10.
- Lima A.C., 1942. Insetos do Brasil, Homopteros. *Scara didattica e Folia Brasiliensis de Agricultura* 3: 321.
- Lin Kung Hsiang, 1926. Yellow shoot of citrus. Symptomatology. Investigations in the cause of huanglongbing. *Natural*

- transmission and spread. General conclusions. *Acta Phytopathologica Sinica* 2: 1-42.
- Lin Kung Shun, Kossischer C.N., da Graça J.V., 1996. Dedicated to Lin Kungshun. *Proceedings of 13th Conference IOCV, IOC, Riverside* 1996.
- Lopes S.A., Martins E.C., Frate G.F., 2005. Detecção de *Candidatus Liberibacter asiaticus* em *Murrays paniculata*. *Anais Fitopatológicos* 11: 48-49.
- Martinez A.L., Wallace J.M., 1967. Citrus leaf mottle-yellow disease in the Philippines and transmission of the causal virus by a psyllid, *Diuraphis citri*. *Plant Disease Reporter* 51: 692-695.
- Martinez A.L., Wallace J.M., 1968. Studies on leaf mottle-yellow disease of citrus in the Philippines. In: *Proceedings of 4th Conference IOCV, University Florida Press, Gainesville* 1968, 167-176.
- Mausson G., Garnier M., Bove J.M., 1976. Transmission of Indian citrus decline by *Trialeurodes erythrae* (DeG. Guercio), the vector of South African greening. In: *Proceedings of 7th Conference IOCV, IOC, Riverside* 1976, 18-20.
- McClean A.P.D., Oberholzer P.C.J., 1965a. Greening disease of sweet orange: evidence that it is caused by a transmissible virus. *South Africa Journal of Agricultural Science* 8: 251-276.
- McClean A.P.D., Oberholzer P.C.J., 1965b. Citrus psylla, a vector of the greening disease of sweet orange. *South Africa Journal of Agricultural Science* 8: 297-298.
- McClean A.P.D., Schwarz R.E., 1970. Greening or Mottle-mottle disease of citrus. *Phytophylactica* 2: 177-194.
- Miyakawa T., Tsuno K., 1969. Occurrence of citrus greening in the southern islands of Japan. *Annot. Phytopathology Society Japan* 66: 667-670.
- Moll J.N., Martin M.M., 1973. Electron microscope evidence that citrus psylla (*Trioza erythrae*) is a vector of greening disease in South Africa. *Phytophylactica* 5: 41-46.
- Moll J.N., Martin M.M., 1974. Comparison of the organism causing greening disease with several plant pathogenic Gram negative bacteria, rickettsia-like organisms and mycoplasma-like organisms. *Proceedings of Conference Le Mycoplasmes/Mycoplasmas, INSTERM* 13: 89-96.
- Murray R.G.E., Schleifer K.P., 1994. Taxonomic notes: a proposal for recording the properties of primitive taxa of prokaryotes. *International Journal of Systematic Bacteriology* 44: 174-176.
- Oberholzer P.C.J., von Staden D.E.A., Basson W.J., 1965. Greening disease of sweet orange in South Africa. In: *Proceedings of 1st Conference IOCV, IOC, University Florida Press, Gainesville* 1965, 213-219.
- Ortiz P., Japónis S., Bove J.M., Garnier M., 1995. Detection and characterization of the African citrus greening liberibacter by amplification, cloning and sequencing of the rpl23L-rpl23B operon. *Current Microbiology* 30: 137-141.
- Pegoni C., Garnier M., Bove J.M., 1996. Detection of the Asian Huanglongbing (greening) liberibacter in Nepal by DNA-DNA hybridisation. In: *Proceedings of 13th Conference IOCV, IOC, Riverside* 1996, 257-270.
- Reinking O.A., 1919. Diseases of economic plants in southern China. *Philippine Agriculturalist* 8: 129-135.
- Reistacher C.N., 1996. The economics of living with citrus diseases - Huanglongbing (greening) in Thailand. In: *Proceedings of 13th Conference IOCV, IOC, Riverside* 1996, 279-285.
- Saglio P., Lafliche D., Boursol C., Bove J.M., 1971. Isolement, culture et observations au microscope électronique des structures de type cytoplasme associé à la maladie du traléorn des agrumes et leur comparaison avec les structures observées dans le cas de la maladie du greening des agrumes. *Physiologie Végétale* 9: 569-582.
- Saglio P., Hospital M., Lafliche D., Dupont G., Bove J.M., Tully J.G., French F.A., 1974. *Synsoplasma citri* gen. and sp. nov. a cytoplasm like organism associated with South born disease of citrus. *Intern Journal of Systematic Bacteriology* 23: 191-204.
- Sulbe A.A., Cortez R.E., 1968. Leaf mottling, a serious virus-disease of citrus in the Philippines. In: *University Florida 1ed1 Proceedings of 4th Conference IOCV, Gainesville* 1968, 111-136.
- Schneider H., 1968. Anatomy of greening-diseased sweet orange shoots. *Phytopathology* 58: 1155-1160.
- Schwartz R.E., Kraus L.C., Pruniera M., 1975. Progress of citrus greening and its psyllid vector in Thailand. *EPD Plant Protection Bulletin* 21: 132-138.
- Subandiyah S., Iwanami T., Taryansu S., Teiki H., 2000. Comparison of 16S rRNA and 16S23S intergenic region sequences among citrus greening organisms in Asia. *Plant Disease* 84: 15-18.
- Sutton B.D., Dixon Y.P., Halber S., Seo Kim-ari, Schulze T., Dixon W., 2005. *Proceedings of 7th International Citrus Canker And Huanglongbing Research Workshop*, Florida Citrus Manual, Orlando, 11-11.
- Tanaka S., Doi Y., 1973. Mycoplasma-like organisms found in citrus plants infected with ikusubi and leaf mottling. In: *International Citrus Congress, Murcia, Valencia*, pp. 353-355.
- Tanaka F.A.O., Kitajima E.W., de Jesus Junio W.C., Ayres A.J., Gimenes-Fernandes N., Bove J.M., 2004. First report on electron micrographs of liberibacter-like structures in citrus in Brazil. In: *16th Conference IOCV, Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias de México, Monterrey, Mexico* p. 177.
- Teixeira D.C., Ayres A.J., Kitajima E.W., Tanaka F.A.O., Danet J.J., Japónis-Silveira S., Sallard C., Bove J.M., 2005a. First Report of a Huanglongbing-like Disease of Citrus in São Paulo State, Brazil, and Association of a New Liberibacter Species, "*Candidatus Liberibacter asiaticus*", with the Disease. *Plant Disease* 89: 107.
- Teixeira D. do Carmo, Danet J.J., Foziliani S., Martins E.C., Jesus Junio W.C., Yamamoto R.T., Lopes S.A., Basson W.J., Ayres A.J., Sallard C., Bove J.M., 2005b. Citrus Huanglongbing in São Paulo State, Brazil. PCR detection of the "*Candidatus Liberibacter asiaticus*" species associated with the disease. *Molecular Cellular Probes* 19: 133-139.
- Teixeira D. do Carmo, Sallard C., Foziliani S., Danet J.J., Ayres A.J., Bove J.M., 2005c. "*Candidatus Liberibacter*

- americanus", associated with citrus huanglongbing (greening disease) in São Paulo State, Brazil. *International Journal of Systematic Evolution Microbiology* 53: 1857-1862.
- Tirtawidjaja S., Hadewidjaja T., Lasheen A.M., 1985. Citrus vein phloem degeneration virus, a possible cause of citrus chlorosis in Java. *Proceedings of the American Society of Horticultural Science* 86: 235-243.
- Van der Merwe A.J., Andersen P.G., 1937. Chromium and manganese toxicity. Is it important in Transvaal citrus greening? *Farming South Africa* 12: 433-440.
- Varma A., Ahlawat Y.S., Chakraborty N.K., Garnier M., Bové J.M., 1993. Detection of the greening BLO by electron microscope, DNA hybridization and ELISA in citrus leaves with and without mottle from various regions in India. In: *Proceedings of 12th Conference IOCV/ICCV Riverside 1993*, 250-286.
- Vasudeva B.S., Capoor S.P., 1958. Citrus decline in Bombay States. *FAO Plant Protection Bulletin* 6: 91.
- Villechanoux S., Garnier M., Bové J.M., 1990. Purification of the bacterium-like organism associated with greening disease of citrus by immunoaffinity chromatography and monoclonal antibodies. *Current Microbiology* 21: 177-180.
- Villechanoux S., Garnier M., Renaudin J., Bové J.M., 1992. Detection of several strains of the bacterium-like organism of citrus greening disease by DNA probes. *Current Microbiology* 24: 89-95.
- Villechanoux S., Garnier M., Lagret F., Renaudin J., Bové J.M., 1993. The genome of the non-cultured, bacterial-like organism associated with citrus greening disease contains the *mscG*, *rpmK*, *rpl*, *rpoB* gene cluster and the gene for a bacteriophage type DNA polymerase. *Current Microbiology* 26: 161-166.
- Weisburg W.G., Barns S.M., Pellmar D.A., Lane D.J., 1991. 16S ribosomal DNA amplification for phylogenetic study. *Journal of Bacteriology* 173: 697-703.
- Wu Z., 2003. Studies on *Campylobacterium chinensis* Scheib, a potential biocontrol predator of *Diuraphis citri* Kunoyama. *Proceedings of 2nd International Citrus Canker And Huanglongbing Research Workshop, Florida Citrus Mutual, Orlando 2003*, H-13.
- Zhao Xue-Yuan, 1981. Citrus yellow shoot disease (huanglongbing) in China - a review. *Proceedings of the International Society of Citriculture* 1: 456-459.
- Zrenik L., Bové J.M., Garnier M., 1998. Phylogenetic characterization of the bacterium-like organism associated with marginal chlorosis of scruberry and proposition of a *Candidatus* taxon for the organism: *Candidatus* Phloemobacter *legumum*. *International Journal of Systematic Bacteriology* 48: 257-261.

Received 7 December 2005

Accepted 31 January 2006