

The IPM Practitioner

Monitoring the Field of Pest Management

Volume XXXIII, Number 7/8, July/August 2011 (Published March 2013)

New Biopesticides for IPM and Organic Production

By William Quarles

Phaseout of the fumigant methyl bromide, restrictions on organophosphates, environmental problems with neonicotinoids and pyrethroids, pest resistance, and the exponential growth of organic agriculture has created a market for alternatives (Weston et al. 2004; Moran 2010; Quarles 2011ab; 1993a). Beneficial nematodes, *Bacillus thuringiensis* (BT), and a large array of fungi, viruses, and bacteria have been developed for greenhouse, turf, field crop, orchard and garden use (Grewal et al. 2005; EPA 2006; Butt et al. 2001; Hom 1996). Microbials are available for treatment of soil, foliar, and postharvest plant pathogens, pest nematodes, herbivorous insects, mosquitoes, structural pests, and weeds. Biocontrol microbials, their pesticidal metabolic products, and other pesticides based on living organisms are classified as biopesticides by the EPA. There are hundreds of registered products (EPA 2013).

Biopesticides do not pose the same regulatory problems seen with chemical pesticides. They are often target-specific, benign to beneficial insects, do not pose air or water quality problems, and crops can be reentered soon after treatment. Naturally occurring microbials can be used in organic production, and human health risks are low. As an added advantage, many pests are not resistant to their effects (EPA 2006; Goettel et al. 2001). Many of the products were developed by small companies, but large companies such as Bayer, Syngenta, and Novozymes have realized the commercial potential, and a flurry of



Photo courtesy of Marrone Bio Innovations

Dr. Pamela Marrone and a laboratory technician inspect biopesticide fermentation equipment. Biopesticides provide an alternative to conventional pesticides in IPM programs, and many formulations have been approved for organic production.

acquisitions has occurred within the last year.

To find a supplier of a product reviewed here, look in Table 1 or Resources. Hundreds of efficacy studies are available in the published literature, and a few of these are cited in the References. To keep this article to a manageable size, beneficial nematodes and BT products will not be discussed here. Microbial metabolites such as spinosad and the avermectins were reviewed earlier (Quarles 1991; 2005c).

Hot New Products

A powerhouse of product development is Marrone Bio Innovations (MBI) (see Resources). One of the

first products was Regalia™. The active ingredient of Regalia is an extract of giant knotweed, *Reynoutria sachalinensis*. Regalia is an effective organic treatment for a large number of plant diseases, and can lead to higher yields. It was

In This Issue

New Biopesticides	1
Book Review	10
EcoWise News	11
Calendar	11
ESA Report	12

The *IPM Practitioner* is published six times per year by the **Bio-Integral Resource Center (BIRC)**, a non-profit corporation undertaking research and education in integrated pest management.

Managing Editor William Quarles
Contributing Editors Sheila Daar
Tanya Drlik
Laurie Swiados
Editor-at-Large Joel Grossman
Business Manager Jennifer Bates
Artist Diane Kuhn

For media kits or other advertising information, contact Bill Quarles at 510/524-2567, birc@igc.org.

Advisory Board

George Bird, Michigan State Univ.; Sterling Bunnell, M.D., Berkeley, CA; Momei Chen, Jepson Herbarium, Univ. Calif., Berkeley; Sharon Collman, Coop Extn., Wash. State Univ.; Sheila Daar, Daar & Associates, Berkeley, CA; Steve Frantz, Global Environmental Options, Longmeadow, MA; Linda Gilkeson, Canadian Ministry of Envir., Victoria, BC; Joseph Hancock, Univ. Calif., Berkeley; William Olkowski, Birc Founder; George Poinar, Oregon State University, Corvallis, OR; Ramesh Chandra Saxena, ICIPE, Nairobi, Kenya; Ruth Troetschler, PTF Press, Los Altos, CA; J.C. van Lenteren, Agricultural University Wageningen, The Netherlands.

Manuscripts

The IPMP welcomes accounts of IPM for any pest situation. Write for details on format for manuscripts or email us, birc@igc.org.

Citations

The material here is protected by copyright, and may not be reproduced in any form, either written, electronic or otherwise without written permission from BIRC. Contact William Quarles at 510/524-2567 for proper publication credits and acknowledgement.

Subscriptions/Memberships

A subscription to the IPMP is one of the benefits of membership in BIRC. We also answer pest management questions for our members and help them search for information. Memberships are \$60/yr (institutions/libraries/businesses); \$35/yr (individuals). Canadian subscribers add \$15 postage. All other foreign subscribers add \$25 airmail postage. A Dual membership, which includes a combined subscription to both the *IPMP* and the *Common Sense Pest Control Quarterly*, costs \$85/yr (institutions); \$55/yr (individuals). Government purchase orders accepted. Donations to BIRC are tax-deductible.
FEI# 94-2554036.

Change of Address

When writing to request a change of address, please send a copy of a recent address label.

© 2013 BIRC, PO Box 7414, Berkeley, CA 94707; (510) 524-2567; FAX (510) 524-1758. All rights reserved. ISSN #0738-968X

Update




Photo courtesy of Michael E. Rogers

The invasive Asian citrus psyllid, *Diaphorina citri*, pictured here, is a vector of citrus greening disease that causes citrus trees to die.

reviewed earlier in the *IPM Practitioner* (see Quarles 2009).

The latest product from MBI is Grandevo™. The active ingredient of Grandevo is produced by the bacterium *Chromobacterium subsugae* (Strain PRAA4-1T). This naturally occurring bacterium was found by USDA researchers in soil underneath a hemlock tree in Maryland. Laboratory tests showed that ingestion of the bacterial suspension killed insects such as adult and nymphal whiteflies, adult stink bugs and cucumber beetles, larval Colorado potato beetles and diamondback moths. The bacterial suspension also had antifeedant properties (Martin et al. 2004; Martin et al. 2007a; Martin et al. 2007b).

According to the EPA, Grandevo is practically non-toxic to mammals, fish, and birds, and is not persistent in the environment. It has low toxicity to beneficials such as lacewings and parasitoids (EPA 2011).

Grandevo is toxic to bees, but the formulation can be applied when bees are not active to mitigate the problem. It is applied to foliage where bees do not normally forage, and biodegrades quickly. Since it is not a systemic, impacts on bees

should be minimal. It should not be applied to water, since it is toxic to some aquatic organisms (EPA 2011).

Grandevo is a dry formulation containing 30% fermentation broth solids from *C. subsugae*. It is labeled for a wide spectrum of pest insects and mites on agricultural and greenhouse crops, including vegetables, fruit, flowers, bedding plants, ornamentals, and turf. Grandevo is toxic by ingestion and has a complex mode of action. It is an antifeedant, and in some instances is repellent and affects pest reproduction. The product is exempt from tolerance, has a four hour re-entry interval, and it is OMRI certified for organic production (Grandevo 2012).

Field tests of Grandevo have found that it often has efficacy similar to some of the standard chemical pesticides. According to a company summary, for western flower thrips, *Frankliniella occidentalis*, on peppers, it had efficacy similar to or better than spinetoram (Radiant™) or spirotetramat (Movento™). It reduced potato psyllid, *Bactericera cockerelli*, on potato by 90%. Effectiveness was similar or better than neem (AZA-Direct™) or buprofezin (Applaud™) for vine mealybug,

Update

Planococcus ficus, on grape (Marrone 2012). It was more lethal than imidacloprid or esfenvalerate to brown marmorated stinkbug, *Halyomorpha halys* (Leskey 2012). When applied to turfgrass, it was more effective for grubs of the Southern masked chafer, *Cyclocephala lurida*, than applications of trichlorfon (Dylox™), which is the industry standard (Stamm et al. 2012). Field tests showed 7 day mortality of 75% to the pecan weevil, *Curculio caryae* (Shapiro-Ilan et al. 2013). It was effective for larvae, but not adults, of the yellowmargin-ed leaf beetle, *Microtheca ochrolema* (Balusu and Fadamiro 2012).

Grandevo may find application for management of Asian citrus psyllid, *Diaphorina citri*. The psyllid is a vector of citrus greening disease that has killed 60 million citrus trees worldwide (Quarles 2010b). Sprays of Grandevo are lethal to psyllid adults and nymphs. Field tests showed effectiveness similar to naled (Dibrom™) or fenproxiimate (Portal™). Treated trees were repellent to the psyllid, and Grandevo reduced psyllid reproduction (Marrone 2012).

Grandevo is likely to be a hit. It is effective for a number of key pests encountered in organic production, such as western flower thrips, two spotted mites, whiteflies, caterpillars, psyllids, and pest beetles (Grandevo 2012). Organic farmers cannot depend totally on BT and spinosad, and pyrethrins can destroy beneficial organisms. Conventional farmers may find Grandevo useful for resistance management and in a rotation to spare beneficial insects.


Other biopesticides in the Marrone pipeline include the bioinsecticide MBI-206 (Venerate™), and the bioherbicides MBI-005 (Opportune™), MBI-010, and MBI-011.

New Bioherbicides

There is a large commercial market for effective, non-polluting bioherbicides that can be used in organic production. There is also a large market for a low impact bioherbicide to manage turfgrass

weeds (Quarles 2010a). Marrone Bio Innovations has two new bioherbicides based on microbials. MBI-005 (Opportune™) is based on *Streptomyces* sp. It was registered with the EPA in 2012, but is not yet registered in California. MBI-010 is based on non-pathogenic *Burkholderia* A396, and it has both contact and systemic effects on pest weeds. EPA registration should be completed in 2013 (Marrone 2012). The company has applied for EPA registration for a bioherbicide MBI-011 based on an extract of long pepper, *Piper longum*, containing the active ingredient sarmentine (Huang et al. 2010; Marrone 2013).

Other bioherbicides include the fungus, *Phoma macrostoma*, which has been commercialized by Scotts Company and was recently registered in California. Phoma™ is expected to be a commercial success for turf weeds. A bioherbicide based on *Sclerotinia minor*



Ixodes* sp. tick infected with the fungus *Metarhizium anisopliae

Photo courtesy of RVAU Denmark

(Sarritor™) is commercially available in Canada, and may eventually be sold in the U.S. (Quarles 2010a).

Fungal Biopesticide Effective for Ticks

The fungus *Metarhizium anisopliae* has appeared sporadically in commercial formulations for structural pest control. A microbial bait station containing *Metarhizium anisopliae* (BioPath™) has been sold for cockroach control. *M. anisopliae* was chosen because it is effective, and because its safety has been


thoroughly tested over the last 100 years (Zimmerman 1993; Mueller-Koegler 1967; Hajek and St. Leger 1994). Another formulation of *M. anisopliae* called Bioblast™ was commercialized for drywood termite control (see Quarles 1997b; 1999c; Quarles and Bucks 1995).

A spray formulation of *M. anisopliae* called MET52™ is now commercially available from Novozymes to control insects on food crops (see Resources). It is also labeled for control of ticks, beetle grubs, and the black vine weevil, *Otiiorhynchus sulcatus*. *M. anisopliae* is a potential alternative to chemical perimeter sprays around structures for pests such as ticks. Field tests show that aqueous formulations of *M. anisopliae* can kill about 50% of adult ticks in an area in about 5 weeks (Zhioua et al. 1997; Benjamin et al. 2002; Quarles 2003). Tests in potting soil show 56-74% mortality of nymphal *I. scapularis* and 8 week persistence of the formulation (Behle et al. 2013). Other insecticidal biopesticides are discussed below.

New Biopesticide for Pest Nematodes

Another hot new product is a biopesticide for pest nematodes. Pest nematodes can be managed with cover crops, crop rotation, and incorporation of organic material into the soil (McSorely 1999). But in some instances, a pest suppression product may be needed. Conventional producers are looking for alternatives to 1,3-dichloropropene, a toxic and carcinogenic material. Organic farmers sometimes need a non-synthetic material compatible with organic methods. Pasteuria BioScience (see Re-sources) has found a practical way to mass produce the bacterium *Pasteuria* spp. The product Econem™ has been registered with the EPA (EPA 2010). Other products are in development, and Pasteuria BioScience was purchased by Syngenta in 2012.

Other biopesticides available for nematodes include formulations of fungi or fungal metabolites. DiTera™ is a formulation of



Capsule shaped *Bacillus subtilis* bacteria


Photo courtesy of AgraQuest Inc.

Bacterial Biopesticides for Plant Disease Protection

The workhorse of commercial microbials is *Bacillus subtilis*. This spore forming bacterium is used primarily as a biofungicide. *B. subtilis* strain QST713 is the most popular and is sold under brandnames such as Jazz™, Cease™, Optiva™, Plant Guardian™, Rhapsody™, Serenade™, and Bayer Advanced™ (CA DPR 2013). *B. subtilis* strain GB03 is sold by Bayer under the brandname Kodiak™. *B. subtilis* MB1600 (Subtilex™) is marketed by Becker Underwood. Many of these formulations became Bayer property when Bayer purchased AgraQuest in 2012.

B. subtilis and other bacterial species have been shown to work as antagonists to powdery mildew and other diseases on foliage. *Bacillus subtilis* is sold by AgraQuest in several formulations, including Serenade™, Serenade Garden™, and Rhapsody™. Serenade is registered for the agricultural market, Serenade Garden for the home market, and Rhapsody is for control of foliar diseases on landscape ornamentals (Marrone 2002; Quarles 2005a).

Serenade has controlled a variety of blights, wilts, rusts and mildews on tomatoes, lettuce, blueberry, plum, cherry, grape, and citrus (Stephan et al. 2005; Isebaert et al. 2002; Agostini et al. 2003). It can be more effective than streptomycin in controlling fireblight caused by *Erwinia* sp. on apples (Aldwinckle et al. 2002; Holtz et al. 2002; Momol et al. 1999). *Pseudomonas fluorescens* (Blight Ban™) is also an



***Bacillus subtilis* attack on powdery mildew**

Photo courtesy of AgraQuest Inc.

effective biopesticide against fire blight (see Nufarm, Resources).

The Rhapsody formulation is an aqueous suspension of a strain of *Bacillus subtilis* known to produce more than 30 lipopeptides that work in concert to destroy disease pathogens. According to AgraQuest, "Rhapsody can be applied in field, greenhouse, interiorscape, residential and commercial landscapes and shadehouse environments. It controls bacterial diseases such as *Pseudomonas*, *Xanthomonas* and *Erwinia* spp. while also controlling common fungal diseases such as powdery mildew, *Botrytis*, Anthracnose and several leaf spot diseases caused by *Alternaria* and *Entomosporium*."

AgraQuest also sells *Bacillus pumilus* (Sonata™) for control of powdery mildew, downy mildew, and rust in agricultural situations. Sonata is very effective for management of early blight caused by *Alternaria solani* in organic tomato production (Wszelaki and Miller 2005).

Bacterial Biopesticides to Control Soil Pathogens

Biocontrol bacteria being sold for soil pathogen management include various species of *Pseudomonas*, *Bacillus* and *Streptomyces*. *Bacillus subtilis* (Kodiak™) manufactured by Bayer is available as a seed or soil treatment, especially for agricultural crops such as soybeans or cotton (Estevez-Jensen et al. 2002; Brannen and Kenny 1997). *Streptomyces griseoviridis* (Mycostop™) controls a number of soil-borne pathogens, including *Fusarium*. It has been used in greenhouse production to protect flowers such as carnations from pathogens (Surviliene 2002; White et al. 1990). This product is available from AgBio in the U.S. (see Resources).

Other biocontrol bacteria for soil pathogens include *Streptomyces lydicus* (Actinovate™) sold by Natural Industries, and *B. subtilis* var *amyloliquefaciens* strain FZB24 (Taegro™) sold by Novozymes (see Resources).

Myrothecium verrucaria metabolites. The fungus is grown in fermentation tanks, then it is killed, and the metabolites are harvested. Since *M. verrucaria* can be herbicidal, this approach reduces the risk of crop damage from a living agent (Anderson and Hallett 2004). DiTera is sold by Valent BioSciences (see Resources). DiTera kills nematodes on contact and may encourage development of microbial populations antagonistic to nematodes (Fernandez et al. 2001; Warrior et al. 1999).

Paecilomyces lilacinus formulations have been developed for nematode suppression. This fungus has been used in India and other countries to manage pest nematodes. It is often applied to the soil along with organic material such as neem cake, and can be as effective as chemical nematicides such as carbofuran (Furadan™) (El-Shan-shoury et al. 2005; Kiewnick 2004; Arif and Parveen 2003).

According to the EPA, the fungus is not likely to cause harm to humans because field application rates are low; it is applied directly to soil; and personal protection equipment is required (EPA 2006). It is sold in the U.S. under the brandname Melocon™ (see Certis, Resources). The living organism does not persist under field conditions, and decays by 90% in 90 days (Kiewnick et al. 2005).




Photo courtesy of AgraQuest Inc.

***Bacillus pumilus* attacks a mildew spore, stopping germination.**

Fungal Biopesticides to Control Soil Pathogens

Beneficial fungi are also sold to suppress soilborne pathogens that cause plant disease (Quarles 1993a; Quarles and Grossman 1995; Quarles 1997a). These beneficial organisms work by competing with pathogens for nutrients and space, by producing antibiotics, by preying on pathogens, or by inducing resistance in host plants (Cook and Baker 1983; Quarles 2002a; Hyakumachi 1998; Isebaert et al. 2002). Commercially available fungi that suppress a broad range of soilborne pathogens include *Trichoderma harzianum* (Rootshield™, Plantshield™), *Gliricladium virens* (SoilGard™) and *G. catenulatum* (PrimaStop™ or PreStop™). *Coniothyrium minitans* (Contans™) is a more specific, and protects against *Sclerotinia* sp. (Jones et al. 2004) (see Resources).

Gliricladium products are sold as a powder that can be mixed into soil, mixed with water, sprayed on foliage, or used as a root dip. *Gliricladium* can be used indoors and outdoors, with vegetables, ornamentals, turf, trees and shrubs. It is applied as a soil treatment to protect crops from *Pythium*, *Rhizoctonia*, *Sclerotium* and other soil pathogens. It is also effective in greenhouse growth media (EPA 2006; Rose et al. 2004; Punja and Yip 2003; Rose et al. 2003; Burns and Benson 2000).

Trichoderma colonizes seeds and protects beans, cotton, peas,

Table 1. Biopesticides Registered in California

Active Ingredient	Name	Class	Company
<i>Bacillus amyloliquefaciens</i> D747, 98.85%	Double Nickel	Fungicide, soil, foliage	Certis
<i>Bacillus pumilus</i> QST2808, 1.38%	Sonata	Fungicide, soil, foliage	AgraQuest
<i>Bacillus subtilis</i> MB1600, 9.9%	Subtilex	Fungicide, soil, turf, ornamentals	Becker Underwood
<i>Bacillus subtilis</i> QST713, 1.34%	Rhapsody	Fungicide, soil, foliage	AgraQuest
<i>Bacillus subtilis</i> QST713, 1.34%	Serenade Soil	Fungicide, soil	AgraQuest
<i>Bacillus subtilis</i> QST713, 1.34%	Cease	Fungicide, soil, foliage	BioWorks
<i>Bacillus subtilis</i> QST713, 14.6%	Serenade Max	Fungicide, soil, foliage	AgraQuest
<i>Bacillus subtilis</i> var <i>amyloliquefaciens</i> FZB24, 24.5%	Taegro	Fungicide, soil, foliage	Novozymes
<i>Bacillus subtilis</i> , GB03, 2.75%	Kodiak	Fungicide, soil	Bayer
<i>Beauveria bassiana</i> GHA, 10.9%	Mycotrol-O	Insecticide	Laverlam, BioWorks
<i>Beauveria bassiana</i> , GHA, WP, 22%	Botanigard	Insecticide	Laverlam, Bioworks
<i>Chenopodium ambrosioides</i> 16.75%	Requiem EC	Insecticide	AgraQuest
<i>Chromobacterium subtsugae</i> 30%	MBI-203 (Grandevo)	Insecticide	Marrone
Codling moth <i>granulosis virus</i>	Madex HP	Insecticide	Certis
<i>Coniothyrium minitans</i> , 5.3%	Contans	Fungicide, soil	Prophyta
<i>Gliricladium virens</i> , GL-21, 12%	SoilGard	Fungicide, soil	Certis
<i>Metarhizium anisopliae</i> , F52, 11%	MET52 EC	Insecticide	Novozymes
<i>Myrothecium verrucaria</i> , 90%	DiTera	Nematicide	Valent Biosciences
<i>P. fumosoroseus</i> strain FE9901	NoFly (EPA registration only)	Insecticide	Natural Industries
<i>P. fumosoroseus</i> 20%	PFR-97	Insecticide	Certis
<i>Paecilomyces lilacinus</i> 6%	Melocon	Nematicide	Certis
<i>Pasteuria usage</i> , .002%	Econem (EPA registration only)	Nematicide	Pasteuria BioScience
<i>Phoma macrostoma</i> , 92%	Phoma Tech	Herbicide	Scotts Company
<i>Pseudomonas fluorescens</i> , 71%	Blightban A506	Fungicide, fireblight	Nufarm
<i>Pseudomonas syringae</i> ESC-10, 29.8%	Bio-Save	Fungicide, postharvest	Jet Harvest
<i>Pythium oligandrum</i>	Polyversum	Fungicide, soil	Gowan
<i>Reynoutria</i> , 20%	Regalia Maxx	Fungicide, foliage	Marrone
<i>Reynoutria</i> , 5%	Regalia	Fungicide, foliage	Marrone
<i>Streptomyces griesoviridis</i> , 35%	Mycostop	Fungicide, soil	Ag Bio
<i>Streptomyces lydicus</i> , 0.0371%	Actinovate	Fungicide, soil	Natural Industries
<i>Streptomyces acidiscabies</i> , 17%	MBI-005 (EPA registration only)	Herbicide	Marrone
<i>T. gamsii</i> , 2% and <i>T. asperellum</i> , 2%	Tenet	Fungicide, soil	Isagrow
<i>T. harzianum</i> , 1.15%	Rootshield	Fungicide, soil	BioWorks
<i>Trichoderma gamsi</i> 2% and 2% <i>T. asperellum</i>	Bio-Tam	Fungicide, soil	AgraQuest
<i>Trichoderma harzianum</i> , 1.15%	Plantshield	Fungicide, soil	BioWorks

cucumbers, tomatoes, radishes, sugar beets, and other crops from *Pythium*, *Rhizoctonia solani*, and *Sclerotium rolfsii*. *Trichoderma* soil treatments may control *Sclerotium cepivorum*, *Verticillium dahliae* and other pathogens (see Quarles 1993a).

BioWorks manufactures formulations of *Trichoderma harzianum* for plant disease suppression (see Resources). Soil mixes, soil drenches and seed treatments for the greenhouse market include powders and granular formulations (Quarles 2006). Yield increases have been seen with corn, beans and cotton (Quarles 1993a; BioWorks 2013). Higher yields and disease biocontrol have also been seen for greenhouse tomatoes (Larkin and Fravel 1998; Utkhede et al. 2001).

A mixture of *Trichoderma* species (Bio-Tam™) is sold by AgraQuest (see Resources) to provide a wider effective temperature range for pathogen protection (AgraQuest 2013). Mycorrhizae and compost teas may also help with plant disease suppression. These subjects are discussed elsewhere (Quarles 1999ab; 2001ab; Ingham 2005ab; Zak 1964).

Insecticidal Biopesticides

A lot of published research has focused on development of microbial insecticides. Two species, *Beauveria bassiana* and *Metarhizium anisopliae*, have been intensely studied and both are produced commercially. There are over 3500 published papers on *Beauveria bassiana* alone. The research shows that *B. bassiana* will infect a wide range of arthropod pests, such as beetles, bugs, mosquitoes, termites, thrips, whiteflies, house flies, grasshoppers, aphids, mites, and ticks (Ugine et al. 2005; Maranga et al. 2005; Lecuona et al. 2005; Gouli et al. 2012).

Beauveria bassiana (Naturalis™) has been used successfully to control whiteflies on cotton (Inglis et al. 2001). Naturalis is sold by Troy BioSciences (see Resources). Another formulation of *B. bassiana* called Botanigard™ is being sold by Laverlam and BioWorks to control

soft bodied insects, especially in greenhouses. It is also available for the home and garden market. Good results have been seen for pest mites, thrips, and whiteflies, but control is best when conditions are humid. The fungus can be used in mite IPM programs, since it is compatible with releases of predatory mites (Jacobson et al. 2001; Gill et al. 1998; Murphy et al. 1998ab; Shipp et al. 2003). It has potential in mole cricket IPM programs (Thompson and Brandenburg 2005), and in management of the emerald ash borer, *Agrilus planipennis* (Liu and Bauer 2008). The 12-day mortality to brown marmorated stinkbug is 100% (Gouli et al. 2012). An organic formulation called Mycotrol O™ is sold by BioWorks (see Resources).

Other fungal products are available especially for control of aphids and whiteflies. *Paecilomyces fumosoroseus* (= *Isaria fumosorosea*) is being marketed in the U.S. by Certis as PFR-97™, and by SePro as Preferal™. The product is targeted mainly for whitefly infestations in greenhouses (Feng et al. 2004). *P. fumosoroseus* Strain FE9901 (No Fly™) is OMRI approved for organic production, but is not registered in California. *Verticillium lecanii* (Vertalec™) is sold in Europe for whitefly and aphid control in greenhouses by Koppert (Fournier and Brodeur 2000).

Milky Spore Disease

A formulation of *Bacillus popilliae* that causes milky spore disease in grubs of Japanese beetles and related species is commercially available (see St. Gabriel, Resources). *B. popilliae* was one of the first microbials produced for the commercial market. It was used extensively in the 1940s over large areas (Lord 2005). Milky spore disease is extremely persistent and may help with longterm control of beetle grubs (Hutton and Burbutis 1974; Ladd and McCabe 1967; Easter 1947). It is difficult to measure field efficacy because longterm effects on beetle populations are complicated by climate, pesticides, patchy distributions, natural yearly

fluctuations, and microbial growth beyond the area of introduction (Lord 2005; Klein et al. 1976; Polivka 1956; Chada et al. 1943).

Viruses and Phages

Finally, a number of virus formulations are available mainly for control of pest caterpillars. Certis has recently registered Madex™, a high potency codling moth granulosis virus (GV) that also affects oriental fruit moth (OFM). Certis also sells the codling moth GV (Cyd-X™) that can be an effective tool in a codling moth IPM program (Arthurs et al. 2004; 2005). A codling moth nuclear polyhedrosis virus (NPV) is sold by Agricola (VPN-80) in the Central and South American markets. Andermatt sells a leafroller GV called Capex™ throughout Europe.

In addition to Madex and Cyd-X, Certis markets a *Heliothis zea* NPV called Gemstar™ and a beet armyworm NPV called Spod-X™. Gemstar is registered for control of pest Lepidoptera, such as the cotton bollworm and cotton budworm. These caterpillars are also pests of corn, soybean and other vegetables. Spod-X is registered for control of the beet armyworm (Kolodny-Hirsch et al. 1997). Certis has also registered a celery looper (*Syngrapha falcifera*) NPV and an alfalfa looper (*Autographa californica*) NPV with the EPA (EPA 2006).

Relatively new virus products are phages for control of bacterial spot diseases caused by *Xanthomonas* spp. and *Pseudomonas* spp. on peppers and tomatoes. These phages are extremely specific for the pathogens and do not persist in the environment (EPA 2006) (see Omnilytics, Resources).

Safety

The products mentioned here are pesticides, and should not be applied carelessly. A review of the literature shows that the commercial products have generally not been implicated in human health problems. Microbials such as *Metarhizium anisopliae* will not grow at temperatures greater than 35°C (95°F), and thus does not gen-

erally infect mammals (Zimmerman 1993).

The active ingredients of these formulations are widespread, naturally occurring microbes, and large numbers of people have been exposed on a regular basis. Though problems rarely occur, there have been published incidents of human infection. Problems have often centered on people with compromised immune systems, sometimes in hospital situations (Tucker et al. 2004). Or, there have been instances of food poisoning when bacteria were allowed to incubate in food (Pavic et al. 2005). *P. lilacinus* has caused fungal skin infections in humans (Hall et al. 2004). Ongoing exposures to microbes and spores could trigger allergies (Goettel et al. 2001). Generally, few problems with these microbes are expected if personal protection is used, and

application is according to label directions (EPA 2006).

Conclusion

Increased regulation of organophosphates, problems with neonicotinoids, phaseout of the fumigant methyl bromide, and newly discovered water quality problems associated with synthetic pyrethroids has created a market for alternatives. Biopesticides are available for treatment of soil, foliar, and postharvest pathogens, pest nematodes, herbivorous insects, structural pests, and weeds. They are generally less destructive to beneficials, cause less environmental pollution, and are less acutely toxic to mammals than conventional pesticides. Biopesticides may provide a satisfactory alternative to chemical pesticides when used as part of an overall IPM plan.

Resources

AgBio Inc., 9915 Raleigh St., Westminster, CO 80031; 877/268-2020, 303/469-9221, Fax 303/469-9598; agbio-inc.com

AgraQuest (see Bayer), 1540 Drew Avenue, Davis, CA 95618; 530/750-0150, Fax 530/750-0153; www.agraquest.com

Bayer Crop Science, PO Box 4913; 8400 Hawthorn Rd., Kansas City, MO 64120; 816/242-2000, Fax 816/242-2659

Becker Underwood, 801 Dayton Ave. PO Box 667, Ames, IA 50010; 800/232-5907, 515/232-5907, Fax 515/232-5961; www.beckerunderwood.com

BioWorks Inc., 100 Rawson Road, Suite 205, Victor, NY 14564; 800/877-9443, 585/924-4362, Fax 800/903-2377; www.bioworksinc.com

Certis, 9145 Guilford Rd. Suite 175, Columbia, MD 21046; 800/847-5620, 301/604-7340, Fax 301/604-7015; www.certisusa.com

Gowan Company, PO Box 5569, Yuma, AZ 85366; 800/883-1844, 928/783-8844, Fax 928/343-9255; www.gowanco.com

Isagro USA, 430 Davis Drive, Suite 240, Morrisville, NC 27560; www.isagro-usa.com

Jet Harvest, PO Box 915139, Longwood, FL 32791; 877/866-5773, 407/523-7842; www.jetharvest.com

Laverlam International (see BioWorks), 117 South Parkmont, PO Box 4109, Butte, MT 59702; 406/782-2386, Fax 406/782-9912

Marrone Bio Innovations, 2121 Second St., Suite B-107, Davis, CA 95618;

530/750-2800; www.marronebioinnovations.com

Natural Industries (see Novozymes), 1230 Cutten Road, Houston, TX 77066; www.naturalindustries.com

Novozymes Biologicals, 5400 Corporate Circle, Salem, VA 32216; 800/788-9886, Fax 540/389-2688; www.novozymes.com

Nufarm USA, 150 Harvester Drive, Suite 200, Burr Ridge, IL 60527; 866/241-0611, 630/455-2000, Fax 630/455-2001; www.nufarm.com

Omnilytics, 9100 South 500 West, Sandy, UT 84070; 866/285-2644, 801/746-3600, Fax 801/746-3461; www.omnilytics.com

Pasteuria BioScience, 12085 Research Drive, Alachua, FL 32615; www.pasteuriabio.com

Prophyta, Inselstrasse 12, D-23999, Malchow, Poel, Germany; www.prophyta.de

St. Gabriel Laboratories, 14044 Litchfield Rd., Orange, VA 22960; 800/801-0061, 540/672-0866, Fax 540/672-0052; www.milkyspore.com

Troy Biosciences, Inc., 113 S. 47th Avenue, Phoenix, AZ 85043; 800/448-2843, 602/233-9047, Fax 602/272-4155; www.troybiosciences.com

Valent BioSciences, 870 Technology Way, Liberty, IL 60048; 800/323-9597, 847/968-4700, Fax 847/968-4780; www.valentbiosciences.com

*for a more complete list, see the *IPM Practitioner's 2012 Directory of Least-toxic Pest Control Products*, www.birc.org

William Quarles, Ph.D., is an IPM Specialist, Executive Director of the Bio-Integral Resource Center (BIRC), and Managing Editor of the IPM Practitioner. He can be reached by email, birc@igc.org.

References

- Agostini, J.P., P.M. Bushong and L.W. Timmer. 2003. Greenhouse evaluation of products that induce host resistance for control of scab, melanose, and Alternaria brown spot of citrus. *Plant Dis.* 87(1):69-74.
- AgraQuest. 2013. Effective natural product solutions for pest management. www.agraquest.com/products/
- Aldwinckle, H.S., M.V.B. Reddy and J.L. Norelli. 2002. Evaluation of control of fire blight infection of apple blossoms and shoots with SAR inducers, biological agents, a growth regulator, copper compounds, and other materials. *Acta Hort.* 590:325-334.
- Anderson, K.I. and S.G. Hallett. 2004. Herbicidal spectrum and activity of *Myrothecium verrucaria*. *Weed Sci.* 52(4):623-627
- Arif, M. and G. Parveen. 2003. Nematode biocontrol, a review of *Paecilomyces lilacinus*. In: P.C. Trivedi, ed. *Advances in Nematology*, Scientific Publishers, Jodhpur, India, pp. 191-204.
- Arthurs, S.P. and L.A. Lacey. 2004. Field evaluation of commercial formulations of the codling moth granulosis virus: persistence of activity and success of seasonal applications against natural infestations of codling moth in Pacific Northwest apple orchards. *Biol. Control* 31(3):388-397.
- Arthurs, S.P., L.A. Lacey and R. Fritts, Jr. 2005. Optimizing use of codling moth granulosis virus: effects of application rate and spraying frequency on control of codling moth larvae in Pacific Northwest apple orchards. *J. Econ. Entomol.* 98(5):1459-1468.
- Balusu, R.R. and H.Y. Fadamiro. 2012. Evaluation of organically acceptable insecticides as standalone treatments and in rotation for managing yellowmargined leaf beetle, *Microtheca ochroloma* in organic crucifer production. *Pest Manag. Sci.* 68:573-579.
- Behle, R.W., M.A. Jackson, and L.B. Flor-Weiler. 2013. Efficacy of a granular formulation containing *Metarhizium brunneum* F52 microsclerotia against nymphs of *Ixodes scapularis*. *J. Econ. Entomol.* 106(1):57-63.
- Benjamin, M.A., E. Zhioua and R.S. Ostfeld. 2002. Laboratory and field evaluation of the entomopathogenic fungus *Metarhizium anisopliae* for controlling questing adult *Ixodes scapularis* (Acari: Ixodidae). *J. Med. Entomol.* 39(5):723-728.
- BioWorks. 2013. Bioworks products. www.bioworksinc.com
- Brannen, P.M. and D.S. Kenney. 1997. Kodiak, a successful biological control product for suppression of soilborne pathogens of cotton. *J. Ind. Microbiol. Biotech.* 19(3):169-171.
- Burns, J.R. and D.M. Benson. 2000. Biocontrol of damping-off of *Catharanthus roseus* caused by *Pythium ultimum* with *Trichoderma virens* and binucleate *Rhizoctonia* fungi. *Plant Dis.* 84(6):644-648.
- Butt, T.M., C. Jackson and N. Magan, eds. 2001. *Fungi as Biocontrol Agents*. CABI Publishing, Wallingford, Oxon, UK. 390 pp.

- CA DPR (California Department of Pesticide Regulation). 2013. Product registration database. www.cdpr.ca.gov
- Chada, H.L., J.A. Dittman and F.C. Daigh. 1943. Progress during 1942 of the program for the colonization of the milky disease of Japanese beetle larvae in Delaware. *Trans. Peninsula Hort. Soc.* 32(5):86-92 [CAB Abstracts]
- Cook, R.J. and K.F. Baker. 1983. *The Nature and Practice of Biological Control of Plant Pathogens*. American Phytopathological Society, St. Paul, MN. 539 pp.
- Easter, S.S. 1947. Control of Japanese beetles at Second Army Posts. *J. Econ. Entomol.* 40(5):632-634.
- El-Shanshoury, A.E.R., S.A. El-Sayed, Y.A.G. Mahmoud and D.M. Khalefa. 2005. Evaluation of *Pochonia chlamydosporia*, *Paecilomyces lilacinus* and *Arthrobotrys dactyloides* as biocontrol agents for *Meloidogyne incognita* under greenhouse conditions. *Pakistan J. Biological Sci.* 8(11):1511-1516. [CAB Abstracts]
- EPA (Environmental Protection Agency). 2006. *New Biopesticide Active Ingredients*. www.epa.gov/pesticides/biopesticides/product-list/
- EPA (Environmental Protection Agency). 2010. *Pasteuria usgae* (006545) fact sheet. 3 pp.
- EPA (Environmental Protection Agency). 2011. Biopesticides registration document, *Chromobacterium subsugae* strain PRAA4-1T. September 27, 2011. 36 pp.
- EPA (Environmental Protection Agency). 2013. Regulating Biopesticides. www.epa.gov/opp00001/biopesticides
- Estevez-Jensen, C., J.A. Percich and P.H. Graham. 2002. Integrated management strategies of bean root rot with *Bacillus subtilis* and *Rhizobium* in Minnesota. *Field Crops Res.* 74(2/3):107-115. [CAB Abstracts]
- Feng, M.G., B. Chen and S.H. Ying. 2004. Trials of *Beauveria bassiana*, *Paecilomyces fumosoroseus* and imidacloprid for management of *Trialeurodes vaporariorum* (Homoptera: Aleyrodidae) on greenhouse grown lettuce. *Biocontrol Sci. Technol.* 14(6):531-544.
- Fernandez, C., R. R.-Kabana, P. Warrior and J.W. Kloepper. 2001. Induced soil suppressiveness to a root-knot nematode species by a nematocide. *Biol. Control* 22(2):103-114.
- Fournier, V. and J. Brodeur. 2000. Dose-response susceptibility of pest aphids (Homoptera: Aphididae) and their control on hydroponically grown lettuce with the entomopathogenic fungus *Verticillium lecanii*, azadirachtin, and insecticidal soap. *Environ. Entomol.* 29(3):568-578.
- Gill, S.A., R. Reeser and M.J. Raupp. 1998. Battling thrips: five pesticides put to the test. *Grower Talks* 62(8):46-48.
- Goettel, M.S., A.E. Hajek, J.P. Siegel and H.C. Evans. 2001. Safety of fungal biocontrol agents. In: Butt et al., pp. 347-375 of 390 pp.
- Gouli, V., S. Gouli, M. Skinner et al. 2012. Virulence of select entomopathogenic fungi to the brown marmorated stinkbug. *Pest Manag. Sci.* 68(2):155-157.
- Grandevo. 2012. Grandevo Label. www.mar-ronebio.com
- Grewal, P.S., R-Udo Ehlers and D.I. S-Ilan, eds. 2005. *Nematodes as Biocontrol Agents*. CABI Publishing, Wallingford, Oxfordshire, UK. 505 pp.
- Hajek, A.E. and R.J. St. Leger. 1994. Interactions between fungal pathogens and insect hosts. *Ann. Rev. Entomol.* 39:293-322.
- Hall, V.C., S. Goyal, M.D.P. Davis and J.S. Walsh. 2004. Cutaneous hyalohyphomycosis caused by *Paecilomyces lilacinus*: report of three cases and review of the literature. *Intl. J. Dermatol.* 43(9):648-653. [CAB Abstracts]
- Holtz, B.A., E.W. Hoffman, S.E. Lindow and B.L. Tevotdale. 2002. Enhancing flower colonization of *Pseudomonas fluorescens* strain A506, and the efficacy of Apogee and Serenade for fire blight control in the San Joaquin Valley of California. *Acta Hort.* 590:319-324.
- Hom, A. 1996. Microbials, IPM and the consumer. *IPM Practitioner* 18(3):1-11.
- Hutton, P.O., Jr. and P.P. Burbulis. 1974. Milky disease and Japanese beetle in Delaware. *J. Econ. Entomol.* 67(2): 247-248.
- Huang, H., C.M. Morgan, R.N. Asolkar, M.E. Koivunen, and P.G. Marrone. 2010. Phytotoxicity of sarmentine isolated from long pepper (*Piper longum*) fruit. *J. Agric. Food Chem.* 58(18):9994-1000.
- Hyakumachi, M. 1998. Systemic resistance in plants induced by beneficial rhizosphere microorganisms. *J. Pesticide Sci.* 23(4):422-426.
- Ingham, E. 2005a. *Compost Tea Brewing Manual*, 5th ed. Soil Foodweb, 728 SW Wake Robin, Corvallis, OR 97330; www.soilfoodweb.com
- Ingham, E. 2005b. *The Field Guide for Actively Aerated Compost Tea*. Soil Foodweb, 728 SW Wake Robin, Corvallis, OR 97330; www.soil-foodweb.com
- Inglis, G.D., M.S. Goettel, T.M. Butt and H. Strasser. 2001. Use of hyphomycetous fungi for managing insect pests. In: Butt et al., pp. 23-69 of 390 pp.
- Isebaert, S., R. Verhoeven and G. Haesaert. 2002. Disease control by means of induced resistance. *Med. Fac. Landbou. Univ. Gent.* 67(2):159-164.
- Jacobson, R.J., D. Chandler, J. Fenlon and K.M. Russell. 2001. Compatibility of *Beauveria bassiana* with *Amblyseius cucumeris* to control *Frankliniella occidentalis* (Thysanoptera: Thripidae) on cucumber plants. *Biocontrol Sci. Technol.* 11(3):391-400.
- Jones, E.E., A. Mead and J.M. Whipps. 2004. Effect of inoculum type and timing of application of *Coniothyrium minitans* on *Sclerotinia sclerotiorum*: control of sclerotinia disease in glasshouse lettuce. *Plant Pathol.* 53(5): 611-620.
- Kiewnick, S. 2004. Biological control of plant parasitic nematodes with *Paecilomyces lilacinus*, strain 251. *Bull. OILB/SROP* 27(1):133-136. [CAB Abstracts]
- Kiewnick, S., C. Rumbos and R.A. Sikora. 2005. [Risk assessment of biological control products]. *Gesunde Pflanzen* 57(6):163-166. [CAB Abstracts]
- Klein, M.G., C.H. Johnson and T.L. Ladd, Jr. 1976. A bibliography of the milky disease bacteria (*Bacillus* spp.) associated with the Japanese beetle, *Popillia japonica* and closely related Scarabaeidae. *Bull. ESA* 22(3): 305-310.
- Kolodny-Hirsch, D.M., T. Sitchawat, T. Jansiri, A. Chenrachaivachirakul and U. Ketunuti. 1997. Field evaluation of a commercial formulation of the *Spodoptera exigua* (Lepidoptera: Noctuidae) nuclear polyhedrosis virus for control of beet armyworm on vegetable crops in Thailand. *Biocontrol Sci. Tech.* 7(4):475-488.
- Ladd, T.L., Jr. and P.J. McCabe. 1967. Persistence of spores of *Bacillus popilliae*, the causal organisms of type A milky disease of Japanese beetle larvae, in New Jersey soils. *J. Econ. Entomol.* 60: 493-495
- Larkin, R.P. and D.R. Fravel. 1998. Efficacy of various fungal and bacterial biocontrol organisms for control of Fusarium wilt of tomato. *Plant Dis.* 82(9):1022-1028.
- Lecuona, R.E., M. Turica, F. Tarocco and D.C. Crespo. 2005. Microbial control of *Musca domestica* (Diptera: Muscidae) with selected strains of *Beauveria bassiana*. *J. Med. Entomol.* 42(3):332-336.
- Leskey, T.C. 2012. Emergence of the brown marmorated stinkbug, *Halyomorpha halys* as a serious pest of agriculture. USDA Appalachian Fruit Research Station, Kearneysville, WV. 75 pp.
- Liu, H.P. and L.S. Bauer. 2008. Microbial control of *Agrilus planipennis* with *Beauveria bassiana* strain GHA: field applications. *Biocontrol Sci. Technol.* 18(6):557-571.
- Lord, J.C. 2005. From Metchnikoff to Monsanto and beyond: the path of microbial control. *J. Invert. Pathol.* 89(1):19-29.
- Maranga, R.O. G.P. Daaya, J.M. Mueke and A.Hassanali. 2005. Effects of combining the fungi *Beauveria bassiana* and *Metarhizium anisopliae* on the mortality of the tick *Amblyomma variegatum* in relation to seasonal changes. *Mycopathologia* 159(4):527-532.
- Marrone, P.G. 2002. An effective biofungicide with novel modes of action. *Pesticide Outlook* 13(5):193-194.
- Marrone Bio Innovations. 2012. Two new species of bacteria: MBI-203 (Grandevo) and MBI-206 insecticides. 28 pp. www.marronebioinnovations.com
- Marrone Bio Innovations. 2013. Press release for MBI-011. February 25, 2013. www.mar-ronebioinnovations.com
- Martin, P.A.W., M. Blackburn and A.D.S. Shromshire. 2004. Two new bacterial pathogens of Colorado potato beetle (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* 97(3):774-780.
- Martin, P.A.W., E. Hirose and J.R. Aldrich. 2007a. Toxicity of *Chromobacterium subsugae* to southern green stink bug (Heteroptera: Pentatomidae) and corn root-worm (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* 100(3):680-684.
- Martin, P.A.W., D. Gundersen-Rindal, M. Blackburn and J. Buyer. 2007b. *Chromobacterium subsugae* sp. nov., a betaproteobacterium toxic to Colorado potato beetle and other insect pests. *Int. J. Systematic Evol. Microbiol.* 57:993-999.
- McSorely, R. 1999. Nonchemical management of plant-parasitic nematodes. *IPM Practitioner* 21(2):1-7.
- Momol, M.T., J.L. Norelli and H.S. Aldwinckle. 1999. Evaluation of biological control agents, systemic resistance inducers, and bactericides for the control of fire blight on apple blossom. *Acta Hort.* 489:553-557.
- Moran, K. 2010. Pesticides in urban runoff, wastewater, and surface water. San Francisco Estuary Partnership. 38 pp. www.up3project.org/documents/up3use2010_Final.pdf
- Mueller-Koegel, E. 1967. Nebenwirkungen Insektenpathogener Pilze auf Mensch und Wirbeltier: Aktuelle Fragen. *Entomophaga* 15(5):429-441.
- Murphy, B.C., T.A. Morisawa and M.P. Parrella. 1998a. Insect-killing fungi: floriculture's IPM future? *Grower Talks* 61(10):60,62,64,66,68.
- Murphy, B.C., T.A. Morisawa, J.P. Newman, S.A. Tjosvold and M.P. Parrella. 1998b. Fungal

Update

- pathogen controls thrips in greenhouse flowers. *Calif. Agric.* 52(3):32-26.
- Pavic, S. et al. 2005. An outbreak of food poisoning in a kindergarten caused by milk powder containing toxigenic *Bacillus subtilis* and *Bacillus licheniformis*. *Archiv. Lebensmittelhygiene* 56(1):20-22. [CAB Abstracts]
- Polivka, J.B. 1956. Effectiveness of milky disease in controlling Japanese beetle in Ohio. *J. Econ. Entomol.* 49(1): 4-6
- Punja, Z.K. and R. Yip. 2003. Biological control of damping off and root rot caused by *Pythium aphanidermatum* on greenhouse cucumbers. *Can. J. Plant Pathol.* 25(4):411-417.
- Quarles, W. 1991. The avermectins: successful biopesticides. *IPM Practitioner* 13(5/6):6-17.
- Quarles, W. 1993a. Alternatives to methyl bromide: *Trichoderma* seed treatments. *IPM Practitioner* 15(9):1-7.
- Quarles, W. and C. Bucks. 1995. Non-toxic termite treatments. *Organic Gardening*:December 44-50.
- Quarles, W. and J. Grossman. 1995. Alternatives to methyl bromide in nurseries: disease suppressive media. *IPM Practitioner* 17(8):1-11.
- Quarles, W. 1997a. Alternatives to methyl bromide in forest nurseries. *IPM Practitioner* 19(3):1-14.
- Quarles, W. 1997b. Bio-Blast. *Pest Control Technology* October: 121,124,126,130.
- Quarles, W. 1999a. Plant disease biocontrol and ectomycorrhizae. *IPM Practitioner* 21(9):1-10.
- Quarles, W. 1999b. Plant disease biocontrol and VAM fungi. *IPM Practitioner* 21(4):1-9.
- Quarles, W. 1999c. Commercial biocontrol for termites. *IPM Practitioner* 21(1):1-6.
- Quarles, W. 2001a. Can composts suppress plant disease? *Common Sense Pest Control Quarterly* 17(3):12-22.
- Quarles, W. 2001b. Compost tea for organic farming and gardening. *IPM Practitioner* 23(9):1-9.
- Quarles, W. 2002a. Aspirin, composts, talking plants, and induced systemic resistance. *IPM Practitioner* 24(5/6):1-9.
- Quarles, W. 2003. Insect biocontrol with fungi. *IPM Practitioner* 25(9/10):1-6.
- Quarles, W. 2005a. Landscape IPM strategies for diseases of ornamentals. *IPM Practitioner* 27(3/4):1-8
- Quarles, W. 2005c. Spinosad finds a home in organic agriculture. *IPM Practitioner* 27(7/8):1-9.
- Quarles, W. 2006. IPM for turf diseases. *Common Sense Pest Control Quarterly* 22(1):12-15.
- Quarles, W. 2009. Giant knotweed, plant disease protection, and immortality. *IPM Practitioner* 31(3/4):1-6.
- Quarles, W. 2010a. Alternative herbicides in turf-grass and organic agriculture. *IPM Practitioner* 32(5/6):1-8.
- Quarles, W. 2010b. New invasives threaten California crops and ornamentals. *IPM Practitioner* 32(7/8):1-7.
- Quarles, W. 2011a. Pesticides and honey bee death and decline. *IPM Practitioner* 33(1/2):1-8.
- Quarles, W. 2011b. Pyrethroid perimeter sprays in structural pest control. *IPM Practitioner* 33(5/6):1-6.
- Rose, S., M. Parker and Z.K. Punja. 2003. Efficacy of biological and chemical treatments for control of Fusarium root and stem rot on greenhouse cucumber. *Plant Dis.* 87(12):1462-1470.
- Rose, S., R. Yip and Z.K. Punja. 2004. Biological control of Fusarium and Pythium root rots on greenhouse cucumbers grown in rock-wool. *Acta Hort.* 635:73-78.
- Shapiro-Ilan, D.I., T.E. Cottrell, M.A Jackson, B.W. Wood. 2013. Control of key pecan insect pests using biorational pesticides. *J. Econ. Entomol.* 106(1):257-266.
- Shipp, J.L., Y. Zhang, D.W.A. Hunt and G. Ferguson. 2003. Influence of humidity and greenhouse microclimate on the efficacy of *Beauveria bassiana* for control of greenhouse arthropod pests. *Environ. Entomol.* 32(5):1154-1163.
- Stamm, M.D., R.W. Baxendale, K.G. Koch and T.J. Porchaska. 2012. *Chromobacterium subtsugae* (MBI-203) for control of southern masked chafers. *Arthropod Management Tests* 37:1-2.
- Stephan, D., A. Schmitt, S.M. Carvalho, B. Seddon and E. Koch. 2005. Evaluation of biocontrol preparations and plant extracts for the control of *Phytophthora infestans* on potato leaves. *Eur. J. Plant Pathol.* 112(3):235-246.
- Survilience, E. 2002. Antagonistic properties of Mycostop (*Streptomyces griseoviridis*) to disease agents in greenhouse plants. *Bull. OILB/SROP* 25(1):257-260.[CAB Abstracts]
- Thompson, S.R. and R.L. Brandenburg. 2005. Tunneling responses of mole crickets (Orthoptera: Gryllotalpidae) to the entomopathogenic fungus *Beauveria bassiana*. *Environ. Entomol.* 34(1):140-147.
- Tucker, D.L., C.H. Beresford, L. Sigler and K. Rogers. 2004. Disseminated *Beauveria bassiana* infection in a patient with acute lymphoblastic anemia. *J. Clin. Microbiol.* 42(11):5412-5414. [CAB Abstracts]
- Ugine, T.A., S.P. Wraight and J.P. Sanderson. 2005. Acquisition of lethal doses of *Beauveria bassiana* conidia by western flower thrips, *Frankliniella occidentalis*, exposed to foliar spray residues of formulated and unformulated conidia. *J. Invert. Pathol.* 90(1):10-23.
- Utkhede, R., C. Bogdanoff and J. McNevin. 2001. Effects of biological and chemical treatments on Botrytis stem canker and fruit yield of tomato under greenhouse conditions. *Can. J. Plant Pathol.* 23(3):253-259.
- Warrior, P., L.A. Rehberger, M. Beach, P.A. Grau, G.W. Kirfman and J.M. Conley. 1999. Commercial development and introduction of DiTera™, a new nematocide. *Pesticide Sci.* 55(3):376-379.
- Weston, D.P., J. You and M.J. Lydy. 2004. Distribution and toxicity of sediment-associated pesticides in agriculture-dominated watersheds of California's Central Valley. *Environ. Sci. Technol.* 38(10):2752-2759.
- White, J.G., C.A. Linfield, M.L. Lahdenpera and J. Uoti. 1990. Mycostop, a novel biofungicide based on *Streptomyces griseoviridis*. *Brighton Crop Protection Council, Pests and Diseases*, Vol 1. pp 221-226.
- Wszelaki, A.L. and S.A. Miller. 2005. Determining the efficacy of disease management products in organically produced tomatoes. *Plant Health Progress* July:1-7. [CAB Abstracts]
- Zak, B. 1964. Role of mycorrhizae in root disease. *Ann. Rev. Phytopathol.* 2:377-392.
- Zhioua, E., M. Browning, P.W. Johnson, H.S. Ginsberg and R.A. LeBrun. 1997. Pathogenicity of the entomopathogenic fungus *Metarhizium anisopliae* to *Ixodes scapularis* (Acari: Ixodidae). *J. Parasitol.* 83(5):815-818.
- Zimmerman, G. 1993. The entomopathogenic fungus *Metarhizium anisopliae* and its potential as a biocontrol agent. *Pesticide Science* 37:375-379.



Subscribe!

Yes! I want to become a member of the Bio-Integral Resource Center and receive a free subscription to:

The IPM Practitioner

Enclosed is my check for:

- ☐ \$60/year, Institutions/
Businesses/Libraries
- ☐ \$35/year, Individual

*** SPECIAL DISCOUNT OFFER**
Receive subscriptions to both *The IPM Practitioner* and the *Common Sense Pest Control Quarterly* for:

- ☐ \$85/year, Institutions/
Businesses/Libraries
- ☐ \$55/year, Individual

Name _____

Address _____

City _____

State _____ Zip _____

Canadian members, add \$10 postage;
Other foreign, add \$20/air. Foreign orders must be paid in U.S. \$ or on an international money order.

Enclose your check
and mail to:

BIRC

PO Box 7414
Berkeley, CA 94707

Planning to change your address?

If so, please notify us six weeks in advance in order not to miss any issues of *The IPM Practitioner*. Just send a label with a copy of your new address, and we'll do the rest! Thanks.

Book Review

The Gardener's Guide to Common-Sense Pest Control

Revised and Updated. 2013.
William Olkowski, Sheila Daar and Helga Olkowski, coauthored and edited by Steven Ash. Taunton Press, Newtown CT. 391 pp.
Paperback. www.tauntonstore.com/pest


The *Gardener's Guide to Common Sense Pest Control* by BIRC founders William Olkowski, Sheila Daar, and Helga Olkowski is back in print at last. For those not familiar with the book, it is not a how-to book about gardening. So you will not find detailed instructions on how to grow geraniums, or a pest by pest analysis of what is eating your cabages.

The *Gardener's Guide* is really a book about ecological gardening with minimal pesticides and low environmental impact. The authors were influenced by Rachel Carson, Robert van den Bosch and others. Conventional gardening leads to overuse of pesticides, producing a "pesticide treadmill" and destructive results on the environment. The authors promote the integrated pest management (IPM) method that combines physical, biological, cultural, ecological, and reduced risk chemical approaches to manage pest populations. Implementation of IPM leads to pesticide reduction.

About half of the book discusses IPM alternatives to conventional pest control, along with principles of sustainable gardening, and best management practices for landscapes. Unfortunately, there is not even one mention of organic gardening. There should have been something on how IPM, sustainable, or reduced risk gardening relates to organic gardening, which for many people is a more familiar concept.

There is a well written section on biological controls that draws on the excellent *Natural Enemies Handbook* produced by the University of California. A list of

insectary plants to encourage beneficial insects is provided. There are discussions on reduced risk or least toxic pesticides such as soaps, oils, botanicals, microbials, and pheromones, which were mostly taken from the first edition of the *Gardeners Guide*. Part of the book is devoted to biopesticides, but many new commercial products are omitted. This is unavoidable, since biopesticides are a major focus of research activity, and several new products have been introduced within the last year.



The overview of pesticide toxicity may be the weakest part of the book. The authors may have been dealing with limited space. There is a good development of acute toxicity, but chronic exposure problems are not examined in enough detail. Endocrine disruption and the special toxicity of pesticides to children are not mentioned. Water quality issues are not addressed. Though there is a sidebar on bees, special problems with systemics and neonicotinoids are not mentioned.

Some might object that lawn maintenance is overemphasized, since it takes up 20% of the book.

Certainly, non-polluting management of turfgrass is a worthwhile subject. Turfgrass is present on playing fields at schools, golf courses and other areas where exposure to pesticides might be a problem. But turfgrass requires large quantities of water and energy, and the prevailing green philosophy is to reduce or eliminate lawns in a home gardening situation. So lawn maintenance is somewhat questionable in an ecologically friendly *Gardener's Guide*.

But the good news is that the sections on turf management and weed management are well written and provide practical tips on low impact turf maintenance. They will be welcomed by professionals trying to implement school IPM programs and others who are stuck with the difficult prospects of environmentally friendly turfgrass maintenance.

Only about 13% of the book is actually concerned with control of specific garden pests. This is not a weakness, since most of the important pests are covered, and detailed information on specific pests can be found online in University of California Pest Notes and in a number of BIRC publications. There are also a number of good gardening books published by Rodale or Taunton Press that fulfill this need.

The design and layout are a vast improvement over the first edition. A major upgrade is the inclusion of 200 striking color photos.

There is no mention of specific products and suppliers, which is probably wise. Each year BIRC spends at least four months upgrading product information for the *IPM Practitioner's Directory of Least-Toxic Pest Control Products*. Any product list for the book would have been out of date within a year.

Congratulations to Steve Ash, Pam Weatherford, and Taunton Press for producing an excellent resource!—*Bill Quarles*

Calendar

January 7-12, 2013. 25th Advanced Landscape IPM Short Course. University of Maryland, College Park. Contact: Avis Koeiman, Dept. Entomology, 301/405-3913. email akoeman@umd.edu

January 23-26, 2013. 33rd Annual EcoFarm Conference. Asilomar, Pacific Grove, CA. Contact: Ecological Farming Association, 831/763-2111; info@eco-farm.org

February 4-7, 2013. Annual Meeting Weed Science Society of America. Baltimore, MD. Contact: www.wssa.net

February 14-17, 2013. Annual Meeting Association Applied IPM Ecologists. Hyatt Regency, San Francisco, CA. Contact: www.aaie.org

February 21-23, 2013. 24th Annual Moses Organic Farm Conference. La Crosse, WI. Contact: Moses, PO Box 339, Spring Valley, WI 54767; 715/778-5775; www.mosesorganic.org

March 4-6, 2013. California Small Farm Conference. Fresno, CA. Contact: www.californiafarmconference.com

March 26, 2013. 22nd Annual UCR Urban Pest Management Conference. UCR Extension, University of California, Riverside. Contact: Kathleen Campbell 951-827-5729.

April 4, 2013. 12th San Francisco Urban IPM Conference. Golden Gate Club, Presidio, San Francisco, CA. Contact: www.sfenvironment.org.

April 5-6, 2013. 31st Annual Beyond Pesticides Conference. Sustainable Families, Food, and Farms. U. New Mexico, Albuquerque, NM. Contact: www.beyondpesticides.org

April 27, 2013. Reduced-Risk Pest Management. El Cerrito City Hall, 10890 San Pablo, El Cerrito, CA. Contact: Melanie Mintz, El Cerrito Environmental Services, 510-215-4350, green@ci.el-cerrito.ca.us

June 18-23, 2013. 70th Annual Convention, Pest Control Operators of CA. Harrah's, Las Vegas, NV. Contact: www.pcoc.org

August 4-9, 2013. 98th Annual Conference Ecological Society of America. Minneapolis, MN. Contact: www.esa.org

August 10-13, 2013. Annual Conference American Phytopathological Society (APS). Austin, TX. Contact: Betty Ford, bford@scisoc.org or www.apsnet.org

October 23-26, 2013. Pestworld, Annual Meeting National Pest Management Association (NPMA), Phoenix, AZ. Contact: NPMA, 10460 North St., Fairfax, VA 22031; 800/678-6722; 703/352-6762 www.npmapestworld.org

November 17-20, 2013. Annual ESA Meeting. Austin, TX. Contact: ESA, 10001 Derekwood Lane, Suite 100, Lanham, MD 20706; 301/731-4535; <http://www.entsoc.org>

December 12-14, 2013. Acres USA Conference. Springfield, IL. Contact: www.acresusa.com

EcoWise News

Hearts Pest Management Wins IPM Innovator Award



Brian Leahy, on the left, Director of the California Department of Pesticide Regulation, presents an IPM Innovator Award to Gerry Weitz, President of Hearts Pest Management of San Diego.

In Sacramento, CA on March 14, Brian Leahy, Director of the California Department of Pesticide Regulation, presented a 2012 IPM Innovator Award to Hearts Pest Management of San Diego. This prestigious award went to Hearts because of its dedication to reduced risk pest management, and its implementation of EcoWise Service throughout San Diego, Los Angeles, Orange, Riverside, and San Bernadino Counties.

Most of the IPM Innovator awards go to organizations that are associated with agriculture, or to non-profits and government agencies associated with urban pest management. Structural pest management companies that receive this award are an elite group. The EcoWise Certified Program received an IPM Innovator Award in 2007. Later awards went to our EcoWise Certified companies. Pestec IPM of San Francisco, CA received the award in 2008, ATCO Pest

Management of Novato, CA received it in 2009, and now Hearts Pest Management in 2012.

EcoWise Certification makes good business sense. At the ceremony, Hearts President Gerry Weitz reported that his EcoWise Certified "green thumb" business increased by 30% during 2012. There is clearly a market for reduced risk pest management, and EcoWise Certification helps government agencies and other customers identify the companies that provide these quality services.

A list of EcoWise Certified Service Providers and Practitioners can be found at www.ecowisecertified.org. An online EcoWise Certification course can be found at www.birc.org. To find out more about Hearts Pest Management, please see their website at www.heartspm.com.

Congratulations, Gerry Weitz and Hearts Pest Management!

Conference Notes

ESA 2012 Annual Meeting Highlights

By Joel Grossman

These Conference Highlights were selected from about 1,800 talks and over 600 poster displays at the Nov. 11-14, 2012, Entomological Society of America (ESA) annual meeting in Knoxville, Tennessee. ESA's next annual meeting is November 10-13, 2013, in Austin, Texas. For more information contact the ESA (10001 Derekwood Lane, Suite 100, Lanham, MD 20706; 301/731-4535; www.entsoc.org)

Pesticide Resistance in Bed Bugs

"If there is widespread resistance to pyrethroids in the field, then we have a problem," as over 90% of the 318 bed bug products in the EPA database are pyrethroids, said Mark Feldlaufer (USDA-ARS, 10300 Baltimore Ave, Beltsville, MD, 20705; mark.feldlaufer@ars.usda.gov). Among the "new" chemistries being tested is deltamethrin, a pyrethroid used since the 1970s. At low doses, deltamethrin provides 13-36% bed bug mortality, but at higher concentrations, it can be effective.

Pyrethroids are the most widely used pesticides against bed bugs, though it often takes more than three treatments to curb infestations, said Alvaro Romero (New Mexico State Univ, Skeen Hall N256, Las Cruces, NM, 88011; aromero2@nmsu.edu). Pesticide resistance was monitored in field populations, and compared to a pesticide-susceptible laboratory bed bug strain, Harlan. Against bed bugs in the field, imidacloprid had a resistance ratio of 286, which was 30 times greater than Harlan. Acetamiprid could not kill bed bugs in the field; and had a resistance ratio of 300,000, which was 286-fold greater than Harlan.

Surveys indicate that bed bugs are the hardest pest to control, said

Kenneth Haynes (Univ of Kentucky, S-225 Ag Sci Center N, Lexington, KY 40546; khaynes@uky.edu). LA-1 is the only field-collected strain still susceptible to pesticides. Field-collected strains of CIN-1 have a 10,000-fold increase in pesticide resistance; though lab colonies of CIN-1 lose some of their resistance. In New Jersey diagnostic assays, pesticide mortality is 50%.

In the USA, 87% of bed bugs show pesticide resistance and have one or more kdr (knockdown resistance) mutations, which are associated with resistance in field-collected bed bugs. However, LA-1 has kdr and no resistance. Thus, kdr mutations are not synonymous with resistance. In lab experiments with Temprid™, a product combining neonicotinoid (imidacloprid) and pyrethroid (cyfluthrin) pesticides, NY-1 bed bugs were killed in 19.2 hours, an indication of relative resistance; versus 2 hours for CIN-1 and an immediate 80% kill of LA-1.

Noting the high pesticide resistance of NY-1 in comparison to LA-1 (susceptible) and CIN-1 (intermediate), Jennifer Gordon (Univ of Kentucky, S-225 Ag Sci Center N, Lexington, KY 40546; jennifer.gordon2@uky.edu) is looking at pesticide resistance as one possible reason for the global resurgence of bed bugs.

Using topical bioassays, Gordon found that resistance evolved in LA-1 after Temprid™ treatments. Indeed, in one generation bed bug control declined from 80% to 40%. In CIN-1, mortality dropped from 70% to 3% after one generation. Along with cross resistance, there is the strong possibility that other products will also be less effective going forward.

Heritable pesticide resistance is also found with Transport®. Indeed, the LA-1 bed bug strain can be selected to go from 100% mortality in 5 minutes to only 30%. Similarly,

CIN-1 can be selected to go from 80% to 10% mortality. There may be a synergy when neonicotinoid and pyrethroid insecticides are applied together; and this could facilitate the evolution of pesticide resistance.

Neem Seed Oil and Bed Bug Traps

Because of resistance, pest management companies are turning to new pesticides and non-chemical methods of treatment. A 22% cold-pressed neem seed oil emulsifiable concentrate (EC), Cirkil™ CX (Terramera; Vancouver, BC, Canada), received EPA registration in May 2012, said Susan Jones (Ohio State Univ, 2501 W. Carmack Rd, Columbus, OH 43210; jones.1800@osu.edu). Neem seed oil is a very fragrant botanical product, though it can be used with odor-neutralizers. Neem oil has been used for many centuries in India in medicines, shampoos, toothpastes, cosmetics and other items. In California, neem seed oil is used to treat bed bugs and their eggs and cellar spiders.

Cirkil™ CX is diluted for application; it is reapplied against bed bugs every 1-3 weeks. A related neem seed oil product, Cirkil RTU (Ready-To-Use), is registered for commerce and consumer spot-treatments; it is available in trigger spray bottles for commercial use, and finger pump spray bottles for consumers.

In October 2012, an empty house with bed bugs was treated. The empty house was monitored with: Verifi™ CO₂ traps in each room at varied locations; and Climbug® Interceptor traps to assess bed bug populations beneath the bed. Four of each trap were installed per room.

Visual inspections revealed few bed bugs. Prior to neem treatments, 38 bed bugs were captured in

Conference Notes

Climbup traps, indicating bed bug infestations only in the master bedroom and bed of the empty house. Eight Verifi traps captured 48 bed bugs in the dining room, guest room and master bedroom. As part of the IPM approach, electrical sockets were treated with MotherEarth® D diatomaceous earth. Gorilla Tape® was used to seal around the doors and exclude bed bug movement from other rooms. Cirkil RTU was sprayed in various places, including on books, backs of picture frames and cardboard boxes. Vials of insecticide-susceptible Harlan bed bugs were placed around the house for on-site neem seed oil vapor toxicity assays.

Two days after spraying, bed bug mortality from neem seed oil vapors was highest in confined spaces; with 48% mortality in vials placed between the mattress and box spring, versus 28% mortality in open spaces. Two weeks post-treatment, 123 dead bed bugs were vacuumed up and live bed bugs were detected in a second bedroom. Bed bug numbers were low because the monitoring traps were doing double duty, also providing population suppression by removing many bed bugs.

Essential Oils Fight Bed Bugs


Heat chambers at 50°C (122°F) are effective against bed bugs, though expensive. Carbon dioxide (CO₂) fumigation chambers have dry ice handling and room air circulation issues. Organophosphate and sulfuryl fluoride fumigants have toxicity issues. Essential oils, many of which are GRAS (Generally Recognized as Safe) compounds and would not need such a long registration process, have been used against head lice and other pests, and are another alternative for bed bug IPM programs, said Dong-Hwan Choe (Univ of California, Entomol 382, Riverside, CA 92521; donghwan.choe@ucr.edu).

Clove essential oils, found in the leaves and flower buds of clove plants, *Syzygium aromaticum*, include GRAS compounds such as

eugenol, beta-caryophyllene and methyl salicylate, which are widely used as flavorings and fragrances in cooking, candy, soaps, chewing gum and medicines.

In experiments at different temperatures, 10 bed bugs were placed in plastic vials with mesh tops. The vials were placed inside 900 ml (1.9 pint) Mason jars; filter paper treated with essential oils was placed on the underside of the Mason jar tops. Methyl salicylate fumigant vapors provided 100% bed bug mortality in 30 hours at 26°C (79°F); 10 hours at 35°C (95°F) and 8 hours at 40°C (104°F) also provided 100% bed bug kill. Eugenol vapors produced similar results. However, there were no synergistic or additive effects from combining eugenol and methyl salicylate.

Mortality takes longer when bed bugs are orally fed essential oils. Choe's future trials will include:



botanical oil granules; exposing bed bug-infested items to essential oil vapors; and checking for sublethal essential oil effects on parameters such as female bed bug reproduction.

Bed Bug Repellents

The idea behind bed bug repellents such as DEET, permethrin, picaridin and essential oils is allowing people to work and travel while having fewer bed bug bites. Plus bringing home fewer bed bugs from infested areas, said Changlu Wang (Rutgers, 93 Lipman Dr, New Brunswick, NJ 08901; cwang@AESOP.Rutgers.edu). Besides repellents, there are also non-chemical alternatives such as: sleeping under bed bug tents, and bandaging yourself in a protective suit (inconve-

nient; expensive; and exposed areas still get bites).

DEET as a bed bug repellent is applied to the feet, the body, or clothing. But DEET is corrosive to plastics and buttons; and there are safety concerns from skin applications. Permethrin, picaridin and essential oils may be less corrosive and potentially safer than DEET; but their efficiency, effectiveness and longevity need to be investigated. Along with permethrin and essential oils, some new botanicals are potential alternatives to 10% DEET for treating fabrics for lasting arthropod repellency.

Isolongifolenone is an odorless sesquiterpene found in the South American Tauroniro tree, *Humiria balsamifera*. It can be synthesized with few impurities from turpentine oil feedstock. Isomers of isolongifolenone repel *Aedes* and *Anopheles* mosquitoes more effectively than DEET; and repel blacklegged ticks, *Ixodes scapularis*, and lone star ticks, *Amblyomma americanum*, as effectively as DEET. A 5% isolongifolenone concentration provides 100% mosquito and tick repellency for 3 hours; this is a higher repellency than DEET. But DEET provides more hours of repellency than isolongifolenone.

In bed bug petri dish assays, 5% DEET is 100% repellent and lasts longer than isolongifolenone, which starts losing its repellency after 3 hours. In arena tests with host cues, 25% DEET keeps surfaces repellent to bed bugs for 2 weeks; 10% DEET protects surfaces for 9 hours; 5% DEET loses its repellency after 9 hours.

Steam Heat and Cold Kill Bed Bugs

Steam, heat and cold temperature treatments can prove useful in bed bug IPM, said Roger Gold (Texas A&M Univ, Minnie Belle Heep Bldg, College Station, TX 77843; r-gold@tamu.edu) and Robert Puckett. Length of exposure was a key variable in these experiments using the bed bug strain Earl. Future studies will look at sublethal effects (e.g. on behavior, reproduc-

Conference Notes

tion) of heat and cold in surviving bed bugs.

In the heat experiments, 10 of each bed bug life stage were placed inside a plastic shoebox containing an arena simulating a bed with mattress fabric from a Sealy® factory. A Jiffy® model J-4000-DM steamer applied drifts of unpressurized steam heat to the fabric for 10, 20 or 30 seconds via placing the steam head edge on the fabric; future studies will look at varying the width of the steam head to increase efficiency. Samples were left up to 7 hours, and weighed to see if they gained weight from the water; a negative, as water is associated with mildew. There was no fabric weight gain from the water; and no mildew, mold or staining on fabric left for two weeks in an attic.

After 7 hours, the brief (10-30 seconds) steaming killed 87-94% of adult and nymph bed bugs on the mattress fabric; temperatures reached 80°C (176°F) in the arena. A few bed bugs managed to escape the steam plume in the arena; but a PCO protocol could close down the escape routes. The 10-30 second steam exposure essentially poached all the bed bug eggs; so there was zero bed bug egg hatch. Steam time is kept necessarily brief, as bed bug nymphs and adults can move fast and must not be given time to escape. The heat conveyed has to be intense, as killing is by heat contact.

Cold is also effective at killing bed bugs. But very low temperatures and significant exposure durations are necessary. Cold is probably less likely to be used than heat or steam by PCOs, as longer time periods mean more labor costs; and days can be involved, depending on the protocol. Subzero freezers were used in the cold tests.

Bed bug life stages were exposed to temperatures ranging from -20°C (-4°F) to -80°C (-112°F) for 1, 2, 5 and 20 minutes. Observations were made at 1, 4, 8 and 24 hours, up to 7 days. Bed bug mortality was 30% after 20 minutes exposure to -20°C (-4°F). However, bed bug mortality was 100% after 20 minutes exposure to -40°C (-40°F). Bed bug mor-

talities were 95% after 10 minutes exposure to -40°C (-40°F); there was some egg hatch at 1 and 2 minutes. Bed bug mortality was 100% after 5, 10 or 20 minutes exposure to -60°C (-76°F).

Metarhizium, Fungicidal Pheromones and Bed Bugs

"Overall, our results suggest that entomopathogenic fungi present a potential method for targeting bed bugs as part of a wider integrated program," said Kevin Ulrich (Univ of Maryland, College Park, MD 20742; kru@umd.edu), who tested both pyrethroid-resistant and pyrethroid-susceptible bed bugs, *Cimex lectularius*. The most effective strain in this study, *Metarhizium anisopliae* strain V1630, should be formulated in a 0.1% dish detergent solution for maximum effectiveness."

Choice of adjuvant for the fungal conidia was very important. In contrast to 0.1% dish detergent, neither 0.01% Tween nor 0.01% sunflower oil provided significantly more bed bug mortality than the control. Interestingly, V1630 was the only *M. anisopliae* strain significantly better than the control in killing bed bugs. V1630 even bested a transgenic *Metarhizium* strain expressing an insect-specific scorpion neurotoxin.

"The fungus has the potential to be a cost effective and relatively straight forward weapon against bed bugs," said Ulrich. However, "the fungicidal characteristics of the bed bug (alarm) pheromones (*E*)-2-hexenal and (*E*)-2-octenal and other cuticular components" could limit control via bed bug contact with the fungal conidia. But oral "ingestion of fungal spores by bed bugs greatly increased overall mortality."

American Cockroach IPM

"Entomopathogenic fungi and diatomaceous earth can play vital roles in eco-friendly control of cockroaches," said Waqas Wakil (Univ of Agriculture, Faisalabad, Pakistan; arid1972@yahoo.com). In laboratory trials, four concentrations (15, 30, 45 and 60 ppm) of the entomopathogenic (insect-killing) fungus

Metarhizium anisopliae were applied topically; both alone and in combination with diatomaceous earth (SilicoSec®) against adult American cockroaches, *Periplaneta americana*. Over a 28 day period in which dead cockroaches were removed daily, mortality from SilicoSec increased over time. At 14 days, the combination of *Metarhizium anisopliae* and SilicoSec caused significantly more American cockroach mortality than either alone. Thus, IPM programs using high concentrations of both products can potentially achieve high cockroach mortality.

Bug Bombs Fail, Baits Work for Cockroach IPM

Also known as "bug bombs," total release foggers (TRFs) typically combine pyrethroid insecticides and a synergist with flammable propellants that in New York City alone are associated with 48 fires and house explosions per year, said Coby Schal (North Carolina State Univ, Campus Box 7613, Raleigh, NC 27695; coby_schal@ncsu.edu). The CDC (Centers for Disease Control and Prevention) recorded 460 TRF-related events in a recent year. Desperate consumers often try TRFs against bed bugs, though they do not work well for that purpose. Consumer mishaps and misuse of TRFs as low-cost alternatives to PCO services prompted comparisons between TRFs and cockroach baits.

TRFs penetrated sealed cabinets to reach dishes and left pesticide residues; but the trap catch data showed that the cockroaches survived. Dose-response curves calculated from cockroaches in open cages exposed to TRFs indicated that pyrethroid insecticide resistance was a factor in TRF treatment failure. Indeed, there was a 200-fold increase in resistance to cypermethrin. In some treatments, cockroach populations actually increased after TRF use. In contrast, professional and consumer cockroach baits like MaxForce® gave high reductions in cockroach populations.

Want Long-Term Pigeon Control?



OvoControl® P

“BIRTH CONTROL” FOR BIRDS

www.ovocontrol.com
or 858.759.8012

EcoWise Certified



Hire an EcoWise Certified Professional
and Ask for EcoWise Service

EcoWise Certified Pest Management Professionals

- Emphasize prevention, exclusion, and non-chemical methods
- Do not apply perimeter sprays that could pollute water
 - Minimize pesticide applications
- Use baits and reduced risk pesticides

To find a company, go to the EcoWise Certified website
www.ecowisecertified.org

Companies and professionals who need certification contact

EcoWise Certified, 510-524-2567 or email birc@igc.org

An Online Certification Course is available on the BIRC website,
www.birc.org. Just use the link www.birc.org/EcoWiseCertified.htm

An EcoWise Certified Technician uses Low Impact IPM Methods such as Baits and Traps for Cockroach Management. *Photos courtesy Pestec IPM Providers*



Every gardener, landscape professional, and small farmer in America should have a copy of this book.

- Completely new edition of the classic 1995 version
- 200 photographs and 100 drawings
- Seasoned authors with over 100 years combined experience
- In-depth chapter on Integrated Pest Management (IPM)
- Includes a natural approach to pest management
- The guidebook to planning and creating a safer garden through sustainable and Integrated Pest Management (IPM) practices.

The Gardener's Guide to Common-Sense Pest Control

William Olkowski, Sheila Daar, Helga Olkowski

Coauthored and edited by Stephen Ash

Paperback

EAN: 9781600855009, Product #071424

8 x 10, 400 pages, 200 full-color photographs, 100 drawings

\$24.95 U.S., \$28.95 Can.



The Taunton Press
Inspiration for hands-on living®



Available wherever books are sold
and at www.tauntonstore.com/pest

CLIMBU^{IP}

INSECT INTERCEPTOR

DETECTION STANDARD AGAINST WHICH OTHERS ARE MEASURED

- people are the best bed bug bait...use under furniture legs where people are sleeping
- early, reliable detection of bed bugs
- all stages are clearly visible against the white surface
- talc-coated interior makes surface too slick to climb out...it is not acting as a pesticide
- easy to clean and easy to retail
- bed bugs are attracted to the fibrous exterior surface to climb up
- tells if bed bugs are going to or coming from furniture
- aids in protecting furniture from reinfestation
- can be reused for months
- made in USA
- patent pending



Manufactured by: Susan McKnight, Inc.
P.O. Box 11406 Memphis, TN 38111
www.insect-interceptor.com • susan@insect-interceptor.com
901-848-3831

"Pest Controls Mother Nature Would Use" NATURE'S CONTROL

Specializing in Beneficial Insects and Organic Pest Controls for Over 20 Years!

- Ladybugs, Spider Mite Predators, Nematodes, Lacewings, and many more "Hired Bugs".
- Mighty Myco Mycorrhizae.
- Magnifiers, Yellow & Blue Traps.
- Quantity Discounts.
- Orders Arrive in 1-2 Days.
- Live Delivery Guaranteed!
- Friendly, Knowledgeable Staff.
- Check our website for the distributor nearest you, or call for your free "Hired Bugs" brochure.

NATURE'S CONTROL

PHONE: (541) 245-6033

FAX: (800) 698-6250

P.O. BOX 35

MEDFORD, OR 97501

www.naturescontrol.com



Crop
Production
Services



A Fatal Attraction For Sucking Insects



Sucking insects are wildly attracted to the color yellow. HOPPERFINDER is coated with a non-drying, water resistant adhesive and holds pests until they die. HOPPERFINDER is an excellent IPM tool that offers these advantages:

- Attracts leafhoppers, aphids, white flies and other insects.
- Compatible with organic and IPM practices.
- Cost effective.
- No Pesticides – Non-toxic to the environment.

For more information call (800) 829-7671 or visit our website at www.cpsagu.com

• PredaLure

Controlled release. Attracts predators/parasites to crop for control of aphids, mites, leafhopper, many others.

• Honey Bee Lure

Controlled release dispenser attracts bees for increased pollination. No spray. No mess.

• Mycostop Biofungicide

• Pentra-Bark Surfactant

• Agri-Fos Fungicide

• Insect Traps

Stink Bugs, Oriental Beetles, Onion Maggot, Thrips, more.

• Pine Beetle Control

• Mycorrhiza

High Potency. Undiluted. University tested.


AGBIO

887-268-2020

Fax 303-469-9598

www.agbio-inc.com

Since 1990



It's a breakthrough in pest control. There's no getting around that.

Introducing GRANDEVO®, a unique, broad-spectrum, biological insecticide/miticide. You've got tough pest management challenges. GRANDEVO meets them head on with a dynamic new approach. It controls today's worst sucking and chewing pests—with complex modes of action that enhance resistance management. Added up, it's a breakthrough unlike anything you've ever seen.

Broad spectrum control. Complex modes of action. Application flexibility.
Go to www.marronebio.com/grandevo for more.

GRANDEVO®

Changing pest management.

 **Marrone®**
Bio Innovations
www.marronebio.com/grandevo



Always read and follow label directions.
©2013 Marrone Bio Innovations, Inc. All rights reserved. Grandevo, the Grandevo logo, Marrone Bio Innovations, and the Marrone Bio Innovations logo are trademarks or registered trademarks of Marrone Bio Innovations, Inc. U.S. Patent No. 7,244,607. Additional patents pending.
18643-03/13

PUT THE POWER IN YOUR HANDS WITH THE ELECTRO-GUN FOR TERMITE CONTROL



**SUSAN FORTUNA, PRESIDENT,
SAYS TEAM UP WITH THE LADY
WITH THE GUN**

CA DPR REG #
:55850-50001-AA



Celebrating 34 years of excellence!


The Discreet Treat

One of the tools of the trade.

The Power of: Higher Profits
Versatility
Greater Customer Satisfaction
IPM & Green Technology

Etex Ltd.
(800) 543-5651
www.etex-ltd.com
egunlady@etex-ltd.com
Established February 1979

PESTEC



Specialists in Structural IPM

- Consulting
- Exclusion • Sanitation • Steam • Vacuuming • Baits

Call us at 925/757-2945; www.ipmprovider.com



**The Ultimate in
Biological Pest Control**

Guardian Nematodes™ Lawn Patrol™

(Steinernema spp. & Heterorhabditis spp. beneficial nematodes)

Application rate: 1 million per 2,000/3,000 sq.ft. of greenhouse
24 million per acre

Pests: Controls over 250 root zone pests including:

- | | | |
|----------------------|--------------------------|-------------------------|
| * Cutworms | * Fungus gnats | * Corn rootworm |
| * Black vine weevils | * White grubs | * Thrips |
| * Sod webworms | * Strawberry root weevil | * Japanese beetle grubs |

Other beneficial items: Encarsia formosa, Phytoseiulus persimilis, Mesoseiulus longipes, Neoseiulus californicus, Aphidoletes aphidimyza, Aphidius, Amblyseius cucumeris, Chrysopa carnea (lacewings), Hippodamia convergens (ladybugs), Nosema locustae (Nolo Bait), Orius, Mealybug predators, etc. Sticky ribbons, Sticky cards, Insect Screens and much more!



**Call TOLL-FREE 1-800-634-6362
for a FREE Catalog**

HYDRO-GARDENS, INC.
Your Total Greenhouse Supplier!
<http://www.hydro-gardens.com>
email: hgi@hydro-gardens.com

P.O. Box 25845, Colorado Springs, CO 80936 * FAX 719-495-2266




**Serving entomology for
more than 60 years**

BioQuip offers a wide selection of pest management equipment including traps, protective clothing, videos, slides, software, books, and more. We also provide thousands of other products and books for entomology and related sciences.

*Contact us to receive our 214-page CD
catalog at no charge.*

*Visit our web site to view monthly product
and book specials, and new products.*



BioQuip Products

2321 Gladwick St.

Rancho Dominguez, CA 90220

Ph: (310) 667-8800

Fax: (310) 667-8808

Email: bqinfo@bioquip.com

www.bioquip.com



Regalia® switches on a plant's natural defenses to fight fungal and bacterial disease.

Regalia is a powerful, proven tool for controlling a broad spectrum of fungal and bacterial diseases, including downy mildew, powdery mildew, botrytis, and walnut blight. Regalia switches on natural defense mechanisms so plants produce – and accumulate – higher levels of natural proteins and other compounds that fight disease development. OMRI® Listed and NOP compliant for organic production, with a four-hour REI and zero-day PHI.

Find out more at www.marronebio.com/regalia

Just add 

 **Marrone®**
Bio Innovations

Always read and follow label directions. Regalia is a registered trademark of Marrone Bio Innovations. © 2012, Marrone Bio Innovations, Inc. 01/12-17646 IPM

Bio-Integral Resource Center

B • I • R • C

P.O. Box 7414, Berkeley, California 94707

ADDRESS SERVICE REQUESTED

NON-PROFIT ORG.
U.S. POSTAGE
PAID
Berkeley, CA
Permit #442

Please renew your membership and help support BIRC. THANK YOU!



Printed with vegetable-based inks
On Processed Chlorine-free paper
100% post-Consumer Waste content

Classified Ads

PRODUCTS



Beneficial Nematode Products for:
Lawn & Garden Insects
Greenhouses & Horticultural Insects
Termites, German Roaches
Turf Insects
Bulk Nematodes Available

BioLogic

For full information see
www.biologicco.com

PO Box 177
Willow Hill, PA 17271

Tel. 717-349-2789


PRODUCTS



RINCON-VITOA
INSECTARIES, INC.
Biological Solutions Since 1950



Aphid Control



Mealybug Control



Fly Control

800-248-BUGS - 805-643-5407 - Ventura, CA - rinconvitova.com

Canadian Producer of Nematodes



BENEFICIAL NEMATODES
Canadian and "Organic"

BENEFICIAL INSECTS
Distributor for Applied Bionomics
Largest Variety Available

Monitoring Tools and Traps

3737 Netherby Road, Stevensville, ON, L0S 1S0
905-382-2904; 905-382-4418(fax)
nic@niagara.com

www.naturalinsectcontrol.com



3244 Gravenstein Hwy. N
Sebastopol, CA 95472

P.O. Box 460, Graton, CA 95444
(707) 823-9125 • FAX (707) 823-1734
info@harmonyfarm.com

*For sustainability...
from the earth to the sun!*

Organic Pest Control

- Beneficial Insects/Nematodes
- Takedown/Spinosad/Neem oil
- Organocide Fungicide/Insecticide
- Insecticidal Soap
- GF 120 Fruit Fly Bait
- JMS Stylet Oil



Custom Design/Build Compost Tea Brewers

We design and build to suite your farm,
nursery and garden needs

- ▶ 20 - 2,000 Gallon Systems
- ▶ AirDrive Technology
- ▶ Easy Cleaning and Maintenance
- ▶ 2 Yr. Warranty on all Parts
- ▶ On-line and Phone Support
- ▶ Order 1-877-432-6249

Visit our website at www.willametteorganics.com/brewmaster

Garlic Barrier
Aphids, Spider Mites, and Earwigs
are quickly controlled by OMRI listed
Garlic Barrier.

See our website:
www.garlicbarrier.com

CLASSIFIED AD RATES—1x rate: \$1 per word. 2x-3x: 80¢ per word. 4x or more: 80¢ per word. Write ad, calculate the cost. **BUSINESS CARD AD RATES** (2 x 3.5")—1x rate: \$55. 2x-3x: \$45 each time. 4x or more \$40 each time. Business card ads must be camera-ready; or BIRC will typeset your ad for \$40. **ALL ADS MUST BE PREPAID.** Send ads and payment to **IPMP Classified Ads**, PO Box 7414, Berkeley, CA 94707. Ads must be received five weeks prior to date of issue in which it is to appear. We reserve the right to refuse materials we feel are inappropriate.