

**REPORT OF *Erpetogomphus sabaeticus* WILLIAMSON, 1918 (ODONATA: GOMPHIDAE)  
FEEDING ON *Diaphorina citri* KUWAYAMA (HEMIPTERA: LIVIIDAE)**

**Takumasa Kondo**

*Corporación Colombiana de Investigación Agropecuaria (Corpoica), Centro de Investigación Palmira, Colombia. E-mail: takumasa.kondo@gmail.com*

**Fredy Palacino-Rodríguez**

*Grupo de Investigación en Odonatos de Colombia, Laboratorio de Artrópodos, Centro Internacional de Física, Universidad Nacional, sede Bogotá / Departamento de Biología, Universidad El Bosque, Bogotá, Colombia. E-mail: odonata17@hotmail.com*

**Richard Danilo Peña-Cuellar**

*Universidad Nacional, sede Palmira, Colombia. E-mail: rdpenac@unal.edu.co*

**RESUMEN**

La libélula *Erpetogomphus sabaeticus* Williamson, 1918 (Odonata: Anisoptera: Gomphidae) se reporta por primera vez depredando a *Diaphorina citri* Kuwayama (Hemiptera: Liviidae). Se provee una lista compilada de más de 63 especies de enemigos naturales de *D. citri* distribuidos en siete órdenes y diecisiete familias de artrópodos (insectos y arañas).

*Palabras clave.* Anisoptera, depredador, enemigos naturales, libélula, listado.

**SUMMARY**

The dragonfly *Erpetogomphus sabaeticus* Williamson, 1918 (Odonata: Anisoptera: Gomphidae) is for the first time reported preying on *Diaphorina citri* Kuwayama (Hemiptera: Liviidae). A compiled list of more than 63 species distributed in seven orders and seventeen families of arthropod (insects and spiders) natural enemies of *D. citri* is provided.

*Key words.* Anisoptera, checklist, dragonfly, natural enemies, predator.

**INTRODUCTION**

The Asian citrus psyllid, *Diaphorina citri* Kuwayama, 1908 (Hemiptera: Liviidae) is an insect vector of the phloem-limited bacteria (*Candidatus Liberibacter* spp.) which cause citrus greening disease (Huanglongbing) or HLB, one of the world's most serious diseases of citrus (Aubert 1990, McClean & Schwartz 1970, Halbert & Manjunath 2004, Mead & Fasulo 2010). Apart from being a vector of HLB, *D. citri* can cause direct damage to its host plant by sucking large amounts of sap, injecting toxins that cause malformation of leaves and shoots and by inducing sooty molds that grow on the honeydew they excrete (Michaud

2004). The Asian citrus psyllid *D. citri* is an economically important citrus pest in many citrus growing regions of the world (Halbert & Manjunath 2004, Manjunath *et al.* 2008). *Diaphorina citri* has a worldwide distribution, and has been recorded from Asia (Afghanistan, Bangladesh, Cambodia, China, Hong Kong, India, Indonesia, Japan, Lao, Macau, Malaysia, Myanmar, Nepal, Pakistan, Philippines, Saudi Arabia, Singapore, Sri Lanka, Taiwan, Thailand, Vietnam), Africa (Mauritius, Réunion), the Caribbean (Bahamas, Belize, Cayman Islands, Cuba, Dominican Republic, Guadeloupe, Jamaica, Puerto Rico, and St. Thomas), Central America (Honduras, Mexico) and from South America (Argentina, Brazil, Colombia,

Paraguay, Uruguay and Venezuela) (Augier *et al.* 2006, Cermeli *et al.* 2007, EPPO 2005, ICA 2010, Kondo & Simbaqueba 2014, Kondo *et al.* 2012). The Asian citrus psyllid was reported in the New World for the first time in Brazil in 1942 (Costa-Lima 1942), and it was only until fairly recently that it was reported from Colombia in 2007. Since then, *D. citri* has been recorded in Colombia from the Departments of Antioquia, Atlántico, Bolívar, Caldas, Cauca, Cesar, Córdoba, Cundinamarca, Magdalena, Meta, Norte de Santander, Quindío, Risaralda, Santander, Sucre, Tolima, and Valle del Cauca (ICA 2010, Ebratt-Ravelo *et al.* 2011a, b, Kondo & Simbaqueba 2014, Kondo *et al.* 2012). In Colombia, *D. citri* is commonly found on *Citrus* spp., *Murraya paniculata* (L.) Jack and *Swinglea glutinosa* (Blanco) Merr. (Rutaceae) (T. Kondo, pers. obs.).

In recent years, populations of *D. citri* have increased in citrus orchards in rural areas and house gardens in urban areas in Colombia. In order to find biological control options to control *D. citri*, natural enemies of *D. citri* were searched in the department of Valle del Cauca, Colombia. During this study, a dragonfly of the family Gomphidae (Odonata: Anisoptera) was captured preying upon an adult *D. citri* in a citrus grove. The adults of some odonate species visit field crops such as cotton and rice in search of prey, helping to control pests of these crops (Rathod *et al.* 2012), including species in the Gomphidae family (Rathod *et al.* 2012, Ghahari *et al.* 2009). This is the first report of an insect belonging to the order Odonata preying on *D. citri*.

## MATERIALS AND METHODS

Field trips were carried out in the Department of Valle del Cauca, between February and May 2014, in the municipalities of Candelaria, El Cerrito, Florida, Palmira, Pradera and Zarzal, in order to collect natural enemies of *D. citri* in citrus (*Citrus* spp.) orchards, and hedges of *M. paniculata* and *S. glutinosa*. Flying insects were collected with an insect net through sweeping, preserved in 70 % ethyl alcohol and labeled with their collecting data. The dragonfly was identified using the keys by Garrison (1994) and Garrison *et al.* (2006).

## RESULTS

The dragonfly was identified as *Erpetogomphus sabaeticus* Williamson, 1918 (Odonata: Anisoptera: Gomphidae) (Fig. 1), a species found in Colombia, Panama and Venezuela (Paulson 2009). In Colombia, *E. sabaeticus* has been recorded from the Departments of Antioquia, Magdalena, Tolima and from Valle del Cauca (present study). *Erpetogomphus sabaeticus* has been found in streams and adjacent forests (Williamson 1918, Garrison 2004), but this is the first record of this species in a cultivated area, namely, a citrus orchard. This is the first report on the eating habits of *E. sabaeticus*, and also the first report of a dragonfly feeding on *D. citri* (Fig. 1).

**Specimen Repository. MCUB-He:** Museo de Ciencias Universidad El Bosque, Bogotá, Colombia.

**Natural enemies of *Diaphorina citri*.** Despite the importance of *D. citri* as a vector of HLB, very few studies on natural enemies of *D. citri* have been carried out in Colombia, with the exception of those that report the presence of the ectoparasitoid *Tamarixia radiata* Waterston (Hymenoptera: Encyrtidae) in various regions of mainland Colombia (Ebratt *et al.* 2011a, b, Kondo *et al.* 2012). Herein we have compiled a list based on records of natural enemies of *D. citri* from the literature plus a gomphid dragonfly collected in the present study. The bulk of natural enemies of *D. citri* are generalist predators. With the addition of *E. sabaeticus*, the list of natural enemies of the class Arthropoda is increased to more than 63 species, distributed in seven orders and seventeen families, namely, Araneae [Anyphaenidae (1 sp.), Miturgidae (1 sp.), Oxyopidae (1 sp.), Salticidae (2 spp.)], Coleoptera [Carabidae (1 sp.), Coccinellidae (28 spp.), Histeridae (1 sp.)], Diptera [Syrphidae (6 spp.)], Hemiptera [Anthocoridae (2 spp.), Geocoridae (1 sp.), Reduviidae (2 spp.)], Hymenoptera [Encyrtidae (1 sp.), Eulophidae (1 sp.), Formicidae (2 spp.)], Neuroptera [Chrysopidae (11 spp.), Hemerobiidae (1 sp.)] and Odonata [Gomphidae (1 sp.)] (Table 1). Of the 63 species of arthropods that have been reported preying on *D. citri*, 7.9 % (5 spp.) are spiders and 92.1 % (58 spp.) are insects.




Figure 1. Remains of *Diaphorina citri* (Hemiptera: Liviidae) found inside mouthparts of the gomphid dragonfly *Erpetogomphus sabaeticus* (Odonata: Anisoptera: Gomphidae).

**Material studied.** *Erpetogomphus sabaeticus* Williamson, 1918 (Odonata: Anisoptera: Gomphidae), 1 adult male ( $\sigma$ ). Colombia: Valle del Cauca, Candelaria, 26.iii.2014, coll. Richard D. Peña Cuellar, collected while feeding on Asian citrus psyllid, *Diaphorina citri*.

Of the 58 species of insects, almost half (48.3 %; 28 spp.) are lady beetles (Coleoptera: Coccinellidae), with the most commonly cited species being *Cyclonedaa sanguinea* (Linnaeus) (Chong *et al.* 2010, González *et al.* 2001, 2002, González-Cárdenas *et al.* 2012, Hall *et al.* 2008, Lozano-Contreras *et al.* 2012, Michaud 2002, 2004, Miranda-Salcedo & López-Arroyo 2010, Rodríguez-Palomera *et al.* 2012, Rodríguez-Toledo *et al.* 2008), *Olla v-nigrum* (Mulsant) (Hall *et al.* 2008, Lozano-Contreras *et al.* 2012, Michaud 2002, 2004, Miranda-Salcedo & López-Arroyo 2010, Rodríguez-Palomera *et al.* 2012), *Chilocorus cacti* (Linnaeus) (González *et al.* 2001, 2002, Lozano-Contreras *et al.* 2012, Rodríguez-Palomera *et al.* 2012, Rodríguez-Toledo *et al.* 2008) and *Harmonia axyridis* (Pallas) (Chong *et al.* 2010, Hall *et al.* 2008, Michaud 2002, 2004). The multicolored Asian lady beetle, *H. axyridis* was recently reported from Colombia (Brown *et al.* 2011, Amat-García *et al.* 2011, Kondo & Gonzalez 2013) where it has become well established (Kondo & Gonzalez 2013). Based on museum insect specimens, Kondo & Gonzalez (2013) estimated the introduction of *H. axyridis* in Colombia to

have occurred in 1989 or earlier, making it the second oldest record of *H. axyridis* in South America after it was deliberately released in Argentina in 1986. The next group of abundant natural enemies are the lacewings (Neuroptera: Chrysopidae) with 20.7 % (12 spp.). Amongst the lacewings *Chrysoperla rufilabris* (Burmeister) (Lozano-Contreras *et al.* 2012, Michaud 2002, 2004, Miranda-Salcedo & López-Arroyo 2010) is the most cited species followed by *Mallada boninensis* (Okamoto) (Chien & Chu 1996, Lin *et al.* 1973, Yang *et al.* 2006) and *Chrysopa* spp. (González *et al.* 2001, 2002, Hall *et al.* 2008, Lozano-Contreras *et al.* 2012, Rodríguez-Toledo *et al.* 2008). On the third place are the hover flies (Diptera: Syrphidae) with 10.3 % (6 spp.), with members of the genera *Allograpta* Osten Sacken (Aubert 1987, Hall *et al.* 2008, Michaud 2002, 2004) and *Ocyptamus* Macquart (González *et al.* 2001, 2002, Michaud 2004, Rodríguez-Toledo *et al.* 2008) being the most common according to the number of bibliographical citations. The rest of natural enemies include the carabid beetle *Egaploa crenulata* (Dejean) (Coleoptera: Carabidae) (Al-Ghamdi 2000), a histerid beetle *Saprinus*

*chalcites* (Illiger) (Coleoptera: Histeridae) (Al-Ghamdi 2000), two species of minute pirate bugs, *Anthocoris* sp. and *Orius* sp. (Hemiptera: Anthocoridae) (Michaud 2002), a bigeyed bug *Geocoris* sp. (Hemiptera: Geocoridae) (Chien & Chu 1996), two species of assassin bugs, *Zelus longipes* (Miranda-Salcedo & López-Arroyo 2010, Hall et al. 2008) and *Zelus* sp. (Michaud 2002) (Hemiptera: Reduviidae), two species of parasitoid wasps *Diphorencyrtus aligarhensis* Shafee, Alam & Agarwal (Hymenoptera: Encyrtidae) (Aubert 1987, Aubert & Quilici 1984, Chien & Chu 1996, Chu & Chien, 1991, García-Darderes 2009, Hoy & Nguyen 1998, Michaud 2002, Yang et al. 2006) and *Tamarixia*

*radiata* (Waterston) (Hymenoptera: Eulophidae) (Baños-Díaz et al. 2013; Chien & Chu, 1996, Chu & Chien 1991, Ebratt-Ravelo et al. 2011a, b, García-Darderes 2009, González et al. 2001, 2002, Hoy & Nguyen 1998, Kondo et al. 2012, Michaud 2002, 2004, Miranda et al. 2011, Miranda-Salcedo & López-Arroyo 2010, Rodríguez-Toledo et al. 2008, Yang et al. 2006), two ant species, namely *Dorymyrmex bureni* (Trager) and *Pseudomyrmex gracilis* (Fabricius) (Hymenoptera: Formicidae) and a brown lacewing *Micromus posticus* (Walker) (Neuroptera: Hemerobiidae) (Table 1).

**Table 1.** Arachnids and insects reported as natural enemies of *Diaphorina citri*.

Taxonomy (Order/Family/Species)	Country and reference
<b>Araneae</b>	
<b>Anypshaenidae</b>	
<i>Hibana velox</i> (Becker, 1879)	USA (Michaud 2002, 2004)
<b>Miturgidae</b>	
<i>Cheiracanthium inclusum</i> (Hentz, 1847)	USA (Michaud 2002)
<b>Oxyopidae</b>	
<i>Oxyope</i> sp.	USA (Michaud 2002)
<b>Salticidae</b>	
<i>Hentzia palmarum</i> (Hentz, 1832)	USA (Michaud 2002)
<i>Marpissa tigrina</i> Tikader, 1965	India (Sanda 1991)
<b>Coleoptera</b>	
<b>Carabidae</b>	
<i>Egaploa crenulata</i> (Dejean, 1829)	Saudi Arabia (Al-Ghamdi 2000)
<b>Coccinellidae</b>	
<i>Arawana</i> sp.	Mexico (Lozano-Contreras et al. 2012)
<i>Azya orbignera</i> Mulsant, 1850	Mexico (Lozano-Contreras et al. 2012)
<i>Brachiacantha dentipes</i> (Fabricius, 1801)	USA (Michaud 2004)
<i>Brumus suturalis</i> (Fabricius, 1798)	India (Husain & Nath 1927, Pruthi & Mani 1945)
<i>Cheiromenes sexmaculata</i> (Fabricius, 1781)	China (Lin et al. 1973), Taiwan (Chien & Chu 1996), India (Husain & Nath 1927, Pruthi & Mani 1945)
<i>Chilocorus cacti</i> (Linnaeus, 1767)	Cuba (González et al. 2001, 2002, Rodríguez-Toledo et al. 2008), Mexico (Lozano-Contreras et al. 2012, Rodríguez-Palomera et al. 2012)
<i>Chilocorus nigrita</i> (Fabricius, 1798)	India (Husain & Nath 1927, Pruthi & Mani 1945)
<i>Chilocorus stigma</i> (Say, 1835)	Mexico (González-Cárdenas et al. 2012), USA (Chong et al. 2010).
<i>Coccinella repanda</i> Thunberg, 1781	India (Husain & Nath 1927, Pruthi & Mani 1945)
<i>Coccinella septempunctata</i> (Linnaeus, 1758)	India (Husain & Nath 1927, Pruthi & Mani 1945)
<i>Coelophora inaequalis</i> (Fabricius, 1775)	USA (Michaud 2002, 2004)
<i>Coleomeguilla maculata</i> (De Geer, 1775)	USA (Michaud 2002)
<i>Curinus coeruleus</i> (Mulsant, 1850)	Mexico (González-Cárdenas et al. 2012), USA (Chong et al. 2010, Hall et al. 2008, Michaud 2002, 2004)
<i>Cyclonedda devestita</i> (Mulsant, 1850)	Venezuela (Solano et al. 2014)

<i>Cyclonedda sanguinea</i> (Linnaeus, 1763)	Cuba (González <i>et al.</i> 2001, 2002, Rodríguez-Toledo <i>et al.</i> 2008), USA (Chong <i>et al.</i> 2010; Michaud 2002, 2004), Mexico (González-Cárdenas <i>et al.</i> 2012, Hall <i>et al.</i> 2008, Lozano-Contreras <i>et al.</i> 2012, Miranda-Salcedo & López-Arroyo 2010, Rodríguez-Palomera <i>et al.</i> 2012)
<i>Delphastus</i> sp.	Mexico (Lozano-Contreras <i>et al.</i> 2012)
<i>Exochomus childreni</i> Mulsant, 1850	USA (Michaud 2002, 2004)
<i>Exochomus cubensis</i> Dimmock, 1906	Cuba (González <i>et al.</i> 2001, 2002)
<i>Harmonia axyridis</i> (Pallas, 1773)	USA (Chong <i>et al.</i> 2010, Hall <i>et al.</i> 2008, Michaud 2002, 2004)
<i>Hippodamia convergens</i> (Guérin-Méneville, 1842)	Mexico (González-Cárdenas <i>et al.</i> 2012, Miranda-Salcedo & López-Arroyo 2010)
<i>Nephus</i> sp.	Mexico (Rodríguez-Palomera <i>et al.</i> 2012)
<i>Olla v-nigrum</i> (Mulsant, 1866)	USA (Hall <i>et al.</i> 2008, Michaud 2002, 2004), Mexico (Lozano-Contreras <i>et al.</i> 2012, Miranda-Salcedo & López-Arroyo 2010, Rodríguez-Palomera <i>et al.</i> 2012)
<i>Pentilia</i> sp.	Mexico (Rodríguez-Palomera <i>et al.</i> 2012)
<i>Rodolia cardinalis</i> (Mulsant, 1850)	USA (Michaud 2004)
<i>Scymnus distinctus</i> Casey, 1924	Cuba (González <i>et al.</i> 2001, 2002)
<i>Scymnus</i> spp.	Brazil (Gravena <i>et al.</i> 1996)
<i>Serangium</i> sp.	Taiwan (Chien & Chu 1996)
<i>Zagloba</i> sp.	Mexico (Lozano-Contreras <i>et al.</i> 2012)
<b>Histeridae</b>	
<i>Saprinus chalcites</i> (Illiger, 1807)	Saudi Arabia (Al-Ghamdi 2000)
<b>Diptera</b>	
<b>Syrphidae</b>	
<i>Allograpta obliqua</i> (Say, 1823)	USA (Michaud 2002, 2004)
<i>Allograpta</i> sp. (probably <i>A. obliqua</i> Say)	USA (Hall <i>et al.</i> 2008)
<i>Allograpta</i> spp.	Nepal (Aubert 1987), Reunion island (Aubert 1987)
<i>Ocyptamus</i> sp.	Cuba (González <i>et al.</i> 2001, 2002, Rodríguez-Toledo <i>et al.</i> 2008)
<i>Ocyptamus fuscipennis</i> (Say, 1823)	USA (Michaud 2004)
<i>Pseudodororus clavatus</i> (Fabricius, 1794)	USA (Michaud 2002)
<b>Hemiptera</b>	
<b>Anthocoridae</b>	
<i>Anthocoris</i> sp.	USA (Michaud 2002)
<i>Orius</i> sp.	USA (Michaud 2002)
<b>Geocoridae</b>	
<i>Geocoris</i> sp.	Taiwan (Chien & Chu 1996)
<b>Reduviidae</b>	
<i>Zelus</i> sp.	USA (Michaud 2002)
<i>Zelus longipes</i> (Linnaeus, 1767)	Mexico (Miranda-Salcedo & López-Arroyo 2010), USA (Hall <i>et al.</i> 2008)
<b>Hymenoptera</b>	
<b>Encyrtidae</b>	
<i>Diphorencyrtus aligarhensis</i> Shafee, Alam & Agarwal, 1975	Argentina (García-Darderes 2009), China (Yang <i>et al.</i> 2006), India, Philippines (Aubert 1987), Reunion island (Aubert & Quilici 1984), USA (Hoy & Nguyen 1998, Michaud 2002), Taiwan (Chien & Chu 1996, Chu & Chien 1991), Vietnam (Aubert 1987)
<b>Eulophidae</b>	
<i>Tamarixia radiata</i> (Waterston, 1922)	Argentina (García-Darderes 2009); Colombia (Ebratt-Ravelo <i>et al.</i> 2011a, b, Kondo <i>et al.</i> 2012), China (Yang <i>et al.</i> 2006), Cuba (Baños-Díaz <i>et al.</i> 2013, González <i>et al.</i> 2001, 2002, Miranda <i>et al.</i> 2011, Rodríguez-Toledo <i>et al.</i> 2008), USA (Hoy & Nguyen 1998; Michaud 2002, 2004), Mexico (Miranda-Salcedo & López-Arroyo 2010), Taiwan (Chien & Chu 1996; Chu & Chien 1991)
<b>Formicidae</b>	
<i>Dorymyrmex bureni</i> (Trager, 1988)	USA (Michaud 2004)

<i>Pseudomyrmex gracilis</i> (Fabricius, 1804)	USA (Michaud 2004)
<b>Neuroptera</b>	
<b>Chrysopidae</b>	
<i>Ceraeochrysa claveri</i> (Navás, 1911)	Mexico (Lozano-Contreras et al. 2012)
<i>Ceraeochrysa cubana</i> (Hagen, 1861)	Mexico (Lozano-Contreras et al. 2012)
<i>Ceraeochrysa everes</i> (Banks, 1920)	Mexico (Lozano-Contreras et al. 2012)
<i>Ceraeochrysa</i> sp. nr. <i>cincta</i> (Schneider, 1851)	Mexico (Lozano-Contreras et al. 2012)
<i>Ceraeochrysa</i> spp.	USA (Michaud 2002, 2004), Mexico (Rodríguez-Palomera et al. 2012)
<i>Ceraeochrysa valida</i> (Banks, 1895)	Mexico (Lozano-Contreras et al. 2012)
<i>Chrysopa septempunctata</i> Wesmael, 1841	China (Yang et al. 2006)
<i>Chrysopa</i> spp.	Cuba (González et al. 2001, 2002, Rodríguez-Toledo et al. 2008), Mexico (Lozano-Contreras et al. 2012), USA (Hall et al. 2008)
<i>Chrysoperla rufilabris</i> (Burmeister, 1839)	USA (Michaud 2002, 2004), Mexico (Lozano-Contreras et al. 2012; Miranda-Salcedo & López-Arroyo 2010)
<i>Chrysoperla</i> spp.	USA (Michaud 2004), Mexico (Lozano-Contreras et al. 2012)
<i>Mallada boninensis</i> (Okamoto, 1914)	China (Lin et al. 1973, Yang et al. 2006), Taiwan (Chien & Chu 1996)
<b>Hemerobiidae</b>	
<i>Micromus posticus</i> (Walker, 1853)	USA (Michaud 2002)
<b>Odonata</b>	
<b>Gomphidae</b>	
<i>Erpetogomphus sabaeticus</i> Williamson, 1918	Present study

## DISCUSSION

A general characteristic of arthropods reported as natural enemies of *D. citri* is that all recorded species are generalist predators, for the exception of the parasitoids *T. radiata* and *D. aligarhensis*. Michaud (2004) reported eight different ant species that were observed on *D. citri* infested terminals, usually harvesting honeydew, of which only two species, namely *Dorymyrmex bureni* and *Pseudomyrmex gracilis* were observed carrying away *D. citri* nymphs. Thus, in Table 1, we only listed these two ant species that had some evidence of predation (i.e., carrying away *D. citri* nymphs), as the other six species listed by Michaud (2004) were only observed nearby the psyllids or tending them (i.e., feeding on the honeydew).

Very few studies have looked into the importance of dragonflies as natural enemies of plant pests, although they may be mentioned as having some degree of predatory role. For example, in a study carried in the Philippines, insects in the order Odonata are listed as being recorded feeding on the brown planthopper, *Nilaparvata lugens* (Stål) (Hemiptera: Delphacidae), a well-known pest of rice plants (*Oryza*

*sativa* L.) (Chiu 1979). However, most studies on dragonflies as natural enemies deal with controlling mosquito larvae. Sebastian et al. (1990) reported a great reduction on a population of *Aedes aegypti* (L.) (Diptera: Culicidae) through augmentative releases of nymphs of a dragonfly, *Crocothemis servilia* (Drury, 1773) (Odonata: Libellulidae) during the rainy season in Yangon (Rangoon), Myanmar. Furthermore, in a study carried out in the suburbs of Omiya City, Japan, during June-July 1979, Urabe et al. (1990) reported that nymphs of the Autumn Darter, *Sympetrum frequens* (Selys 1883) (Odonata: Libellulidae) plays an important role as a natural enemy of the larvae of the mosquito, *Anopheles sinensis* Wiedemann, 1828 (Diptera: Culicidae).

Although, hitherto dragonflies have not been listed as natural enemies of *D. citri*, these generalist predators may contribute to the reduction of populations of *D. citri* in orchards that are found near marshes, lakes, ponds, streams, and wetlands. Beneficial insects may help control insect pest populations by enhancing their efficacy and local abundance through modification of the environment or existing management practices, a practice known as conservation biological control (Barbosa 1998). This pest

management method using dragonflies as natural enemies may be efficient in such areas near water bodies where there is little use of pesticides.

## CONCLUSIONS

With the addition of *E. sabaeticus*, the number of known arthropod natural enemies is increased to more than 63 species. Further studies are needed to determine the predation rate of *E. sabaeticus* on *D. citri* populations in areas

where the distribution of the two species overlap.

## ACKNOWLEDGEMENTS

We are thankful to Antonio Gómez Anaya and Rodolfo Novelo Gutiérrez of the Department of Entomology, Instituto de Ecología, Xalapa, Veracruz, México, for kindly reviewing an earlier draft of the manuscript. Thanks also to anonymous reviewers who helped improve the manuscript.

## LITERATURE CITED

- Al-Ghamdi, K.M.S. 2000. A field study on synchrony between the populations of citrus *Psylla, Diaphorina citri* (Kuwayama) [sic.] (Homoptera: Psyllidae) and its natural enemies in western Saudi Arabia. Bull. Fac. Agric., Cairo University, 51: 227-238.
- Amat-García, G., E. Amat-García & E. Ariza-Marín. 2011. Insectos invasores en los tiempos de cambio climático. Revista Innovación y Ciencia, 18: 45-53.
- Aubert, B. 1987. *Trioza erytreae* del Guercio and *Diaphorina citri* Kuwayama (Homoptera: Psylloidea), the two vectors of citrus greening disease: Biological aspects and possible control strategies. Fruits, 42: 149-162.
- Aubert, B. 1990. Integrated activities for the control of huanglongbing greening and its vector *Diaphorina citri* Kuwayama in Asia. Pp. 133–144, in Proceedings of the Fourth FAO-UNDP International Asia Pacific Conference on Citrus Rehabilitation, 4–10 February 1990 (B. Aubert, S. Tontyaporn & D. Buangsuvon, eds). Chiang Mai, Thailand,
- Aubert, B. & S. Quilici, 1984. Biological control of the African and Asian citrus psyllids (Homoptera: Psylloidea) through euphorid and encyrtid parasites (Hymenoptera: Chalcidoidea) in Reunion Island. Pp. 100–108, in Proceedings of the 9th Conference of International Organization of Citrus Virologists. International Organization of Citrus Virologists (L. W. Timmer, S. M. Garnsey & L. Navarro, eds.). Valencia, Spain.
- Augier, L., G. Gastaminza, M. Lizondo, M. Argañaraz & E. Willink. 2006. Presencia de *Diaphorina citri* (Hemiptera: Psyllidae) en el Noroeste Argentino (NOA). Revista de la Sociedad Entomológica Argentina, 65: 67-68.
- Baños-Díaz, H. L., I. Miranda & M. A. Martínez. 2013. Biología y tabla de vida de *Tamarixia radiata* Waterston bajo condiciones controladas. Revista de protección vegetal, 28: 120-126.
- Barbosa, P. 1998. Conservation Biological Control. Academic Press, San Diego, California, USA.
- Brown, P. M. J., C. E. Thomas, E. Lombaert, D. L. Jeffries, A. Estoup & L.L. Handley. 2011. The global spread of *Harmonia axyridis* (Coleoptera: Coccinellidae): distribution, dispersal and routes of invasion. BioControl, 56: 623-641.
- Cermeli, M., P. Morales-Valle, J. Perozo & F. Godoy. 2007. Distribución del psílido asiático de los cítricos (*Diaphorina citri* Kuwayama (Hemiptera, Psyllidae) y presencia de *Tamarixia radiata* (Waterston) (Hymenoptera, Eulophidae) en Venezuela. Entomotrópica, 22: 181-184.
- Chien, C. C. & Y. I. Chu. 1996. Biological control of citrus psyllid, *Diaphorina citri* in Taiwan. Biological Pest Control in Systems of Integrated Pest Management—1996. Food and Fertilizer Technology Center Book Series No. 47, Taipei, Taiwan.
- Chiu, S. C. 1979. Biological control of the brown planthopper. Brown Planthopper: Threat to rice production in Asia, 335-355.

- Chong, J. H., A. L. Roda & C. M. Mannion. 2010. Density and Natural Enemies of the Asian Citrus Psyllid *Diaphorina citri* (Hemiptera: Psyllidae), in the Residential Landscape of Southern Florida. *Journal of Agricultural and Urban Entomology*, 27: 33-49.
- Chu, Y. I. & C. C. Chien. 1991. Utilization of natural enemies to control of psyllid vectors transmitting citrus greening. Pp. 135–145, in Integrated Control of Plant Virus Diseases. Proceedings of the International Workshop TARI, Taichung, Taiwan, April 9-14, 1990 (K. Kiritani, H.J. Su & Y.I. Chu, eds.), Food and Fertilizer Technology Center for the Asian and Pacific Region. Taipei, Taiwan.
- Costa-Lima, A. M. 1942. Homopteros. Insetos do Brazil. Arquivos da Escola Superior de Agricultura e Medicina Veterinaria. Rio de Janeiro, 3: 1-327.
- Ebratt-Ravelo, E. E., L. T. Rubio-González, V. A. Costa, E. M. Zambrano-Gómez, A. P. Castro-Ávila & M. Y. Santamaría Galindo, 2011a. Primer registro de *Tamarixia radiata* (Waterston, 1922) (Hymenoptera: Eulophidae) en Colombia. *Revista Facultad Nacional de Agronomía, Medellín*, 64: 6141-6146.
- Ebratt-Ravelo, E. E., L. T. Rubio-González, V. A. Costa, Castro Ávila A. P., E. M. Zambrano-Gómez & J. E. Ángel-Díaz. 2011b. *Diaphorina citri* (Kuwayama, 1907) and *Tamarixia radiata* (Waterston, 1922) in citrus crops of Cundinamarca, Colombia. *Agronomía Colombiana*, 29: 487-493.
- European and Mediterranean Plant Protection Organization (EPPO). 2005. Data Sheets on Quarantine Pests: *Diaphorina citri*. [http://www.eppo.org/QUARANTINE/insects/Diaphorina\\_citri/DIAACI.ds.pdf](http://www.eppo.org/QUARANTINE/insects/Diaphorina_citri/DIAACI.ds.pdf) (consultada 12/10/2014).
- García-Darderes, C. S. 2009. *Diaphorina citri* Kuwayama (Hemíptera: Psyllidae), vector de la bacteria que causa el Huanglongbing (HLB – Greening). Ministerio de la Producción Secretaría de Agricultura, Ganadería, Pesca y Alimentación. Servicio Nacional de Sanidad y Calidad Agroalimentaria SENASA. Buenos Aires, Argentina.
- Garrison, R. 1994. A revision of the New World genus *Erpetogomphus* Hagen in Selys (Odonata: Gomphidae). *Tijdschrift voor Entomologie*, 137: 173-269.
- Garrison, R., N. Von Ellenrieder & J. Louton. 2006. Dragonfly genera of the New World. An illustrated and annotated key to the Anisoptera. The Johns Hopkins University Press, Baltimore.
- Ghahari, H., M. Tabari, H. Sakenin, H. Ostovan & S. Imani. 2009. Odonata (Insecta) from Northern Iran, with comments on their presence in rice fields. *Munis Entomology & Zoology*, 4: 148-154.
- González, C., D. Hernández & J. Rodríguez. 2001. Influencia de los enemigos naturales en el comportamiento de *Diaphorina citri* Kuw. (Homoptera, Psyllidae) en los cítricos de Cuba. Pág. 273. Presented at the IV Seminario Científico Internacional de Sanidad Vegetal. Varadero, Cuba.
- González, C., D., Hernández, R. I. Cabrera & J. R. Tapia. 2002. *Diaphorina citri* Kuw., inventario y comportamiento de los enemigos naturales en la citricultura cubana. Paper presented at the taller sobre plagas emergentes de los cítricos. Resúmenes de las sesiones orales. Instituto de Investigaciones de Cítricos y otros Frutales. La Habana, Cuba.
- González-Cárdenas, J. C., I. E. Castellanos Sturemark, L. J. Fucikovsky Zac, M. López Herrera & G. Sánchez Rojas. 2012. Coccinélidos como potenciales enemigos naturales de *Diaphorina citri* (Hemiptera: Psyllidae) en un huerto de cítricos en Tuxpan, Veracruz, México. *Revista Científica UDO Agrícola* 12 (4): 855-860.
- Gravena, S., M. J. G. Beretta, P. E. B. Paiva, R. Gallão & P. T. Yamamoto. 1996. Seasonal abundance and natural enemies of *Diaphorina citri* (Hemiptera: Psyllidae) in citrus orchards of São Paulo State, Brazil. Pag. 414, in Proceedings 13th Conference of the International Organization of Citrus Virologists (IOCV) (J.V. da Graça, P. Moreno & R.K. Yokomi, eds). University of California, Riverside.
- Halbert, S. E. & K. L. Manjunath. 2004. Asian citrus psyllids (Sternorrhyncha: Psyllidae) and greening disease of citrus: a literature review and assessment of risk in Florida. *Florida Entomologist*, 87: 330-353.

- Hall, D. G., M. G. Hentz & R. C. Adair Jr. 2008. Population ecology and phenology of *Diaphorina citri* (Hemiptera: Psyllidae) in two Florida citrus groves. Environmental Entomology, 37: 914-924.
- Hoy, M. A. & R. Nguyen. 1998. Citrus psylla: here in Florida - an action plan. Pest Alert. <http://extlab7.entnem.ufl.edu/PestAlert/hoy-0615.htm>
- Husain, M. A. & D. Nath. 1927. The citrus psylla (*Diaphorina citri*, Kuw.) [Psyllidae: Homoptera]. Memoirs of the Department of Agriculture in India. Entomology Series, 10: 27.
- ICA [Instituto Colombiano Agropecuario]. 2010. Situación actual de HLB (Huanglonbing) y su vector el psilido asiático de los cítricos (*Diaphorina citri* Kuwayama) en Colombia. Boletín Epidemiológico. [http://www.ica.gov.co/Areas/Agricola/Servicios/Epidemiologia-Agricola/BOLETINES/Nacionales/2010/BN\\_Diaphorina\\_2010.aspx](http://www.ica.gov.co/Areas/Agricola/Servicios/Epidemiologia-Agricola/BOLETINES/Nacionales/2010/BN_Diaphorina_2010.aspx)(consultada15/10/2014).
- Kondo, T. & F. G. González. 2013. The multicolored Asian lady beetle, *Harmonia axyridis* (Pallas, 1773) (Coleoptera: Coccinellidae), a not so new invasive insect in Colombia and South America. Insecta Mundi, 0283: 1-7.
- Kondo, T., E. M. Quintero, M. Campuzano, K. A. G. Wyckhuys & J. Heraty. 2012. First report of *Tamarixia radiata* (Waterston) (Hymenoptera: Encyrtidae), a parasitoid of the Asian citrus psyllid *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae) in the department of Valle del Cauca, Colombia. Boletín del Museo de Entomología de la Universidad del Valle, 13: 48-51.
- Kondo, T. & R. Simbaqueba-Cortés. 2014. *Sarucallis kahawaluokalani* (Kirkaldy) (Hemiptera: Aphelinidae), a new invasive aphid on San Andres island and mainland Colombia, with notes on other recent invasive species. Insecta Mundi, 0362: 1-10.
- Lin, S. J.; Y. F. Ke & C. C. Tao. 1973. Bionomics observation and integrated control of citrus Psylla, *Diaphorina citri* Kuwayama. Journal of Horticultural Society of China, 19: 234-242.
- Lozano-Contreras, M. G., M. Guadalupe & J. J. Argumedo. 2012. Identificación de enemigos naturales de *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae) en el Estado de Yucatán, Mexico. Fitosanidad, 16: 5-11.
- Manjunath, K. L., S. E. Halbert, C. Ramadugu, S. Webb & R. F. Lee. 2008. Detection of 'Candidatus Liberibacter asiaticus' in *Diaphorina citri* and its importance in the management of citrus Huanglongbing in Florida. Phytopathology, 98: 387-396.
- McClean, A. P. D. & R. E. Schwartz. 1970. Greening of blotchy-mottle disease in citrus. Phytophylactica, 2: 177-194.
- Mead, F. W. & T. R. Fasulo. 2010. Asian citrus psyllid, *Diaphorina citri* Kuwayama (Insecta: Hemiptera: Psyllidae). FDACS/DPI Entomology. Circular 180. University of Florida, Gainesville, FL.
- Michaud, J. P. 2002. Biological control of Asian citrus psyllid, *Diaphorina citri* (Hemiptera: Psyllidae) in Florida: a preliminary report. Entomological News, Philadelphia, 113: 216-222.
- Michaud, J. P. 2004. Natural mortality of Asian citrus psyllid (Homoptera: Psyllidae) in central Florida. Biological Control, 29: 260-269.
- Miranda, I., H. Baños-Díaz, Y. Pérez-Aranda & M. A. Martínez. 2011. Patrón espacial y parámetros de crecimiento de *Diaphorina citri* Kuwayama y su parasitoide *Tamarixia radiata* Waterston sobre *Murraya paniculata* Linn. Revista de protección vegetal, 26: 100-104.
- Miranda-Salcedo, M. A. & J. I. López-Arroyo. 2010. Avance de investigación para el manejo del psilido asiático de los cítricos *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae) en Michoacán. Pp. 73-85, in Memorias del 3er Congreso Nacional Mitigación del daño ambiental en el Sector Agropecuario de México, México.
- Pruthi, H. S. & M. S. Mani. 1945. Our knowledge of the insect and mite pests of citrus in India and their control. Imperial Council of Agricultural Research, Scientific Monograph 16. Government of India Press, Calcutta.
- Paulson, D. R. 2009. *Erpetogomphus sabaleticus*. The IUCN Red List of Threatened Species. Version 2014.2. <http://www.iucnredlist.org>

- Rathod, P. P., S. S. Pawar, N. A. Manwar & I. A. Raja. 2012. Diversity and abundance of dragonflies and damselflies (Order - Odonata) in agroecosystems around the Amravati city (m.s.), India in monsoon season. Golden Research Thoughts, 3: 1-3.
- Rodríguez-Palomera, M., J. Cambero-Campos, A. Robles-Bermúdez, C. Carvajal-Cazola & O. Estrada-Virgen. 2012. Natural enemies associated to *Diaphorina citri* Kuwayama (Hemiptera: Psyllidae) in *Citrus latifolia* Tanaka, in the State of Nayarit, Mexico. Acta Zoológica Mexicana, nueva serie, 28: 625-629.
- Rodríguez-Toledo, O., C. C. González-Fernández, C. L. Marrero-Artabe & L. Robledo-Ortega. 2008. Bases para el manejo de *Diaphorina citri* Kuw. (Hemiptera: Psyllidae) en fomento de toronjo Marsh en la localidad de Jagüey Grande. Retrieved from CD de Monografías 2008, Universidad de Matanzas "Camilo Cienfuegos", Cuba.
- Sanda, G. L. 1991. Mode of hunting and functional response of the spider *Marpissa tigrina* Tikader (Salticidae: Arachnida) to the density of its prey, *Diaphorina citri*. Entomon, 16: 279-282.
- Sebastian, A., M-M. Sein, M-M. Thu & P. S. Corbet. 1990. Suppression of *Aedes aegypti* (Diptera: Culicidae) using augmentative release of dragonfly larvae (Odonata: Libellulidae) with community participation in Yangon, Myanmar. Bulletin of Entomological Research, 80, pp 223-232. doi:10.1017/S0007485300013468.
- Solano, Y. A., E. Arcaya & G. González. 2014. Actividad depredadora de *Cycloneda devestita* (Mulsant) (Coleoptera: Coccinellidae) en el estado Lara, Venezuela. Boletín de la Sociedad Entomológica Aragonesa, (S.E.A), 54: 423-424.
- Urabe, K., T. Ikemoto & S. Takei. 1990. Studies on *Sympetrum frequens* (Odonata: Libellulidae) nymphs as natural enemies of the mosquito larvae, *Anopheles sinensis*, in rice fields. 4. Prey-predator relationship in the rice field areas. Japanese Journal of Sanitary Zoology, 41(3): 265-272.
- Williamson, E. B. 1918. Results of the University of Michigan-Williamson expedition to Colombia 1916-17. I. Two interesting new Colombian Gomphines (Odonata). Occasional Papers of the Museum of Zoology, University of Michigan, 52: 1-14.
- Yang, Y., M. Huang, G. Andrew, C. Beattie, Y. Xia, G. Ouyang & J. Xiong. 2006. Distribution, biology, ecology and control of the psyllid *Diaphorina citri* Kuwayama, a major pest of citrus: A status report for China. International Journal of Pest Management, 52: 343-352.

Recibido abril 14, 2015, publicado julio 2015.