

## Candidate gene makers for *Candidatus Liberibacter asiaticus* for detecting citrus greening disease

MADHUGIRI NAGESWARA-RAO<sup>1,2,\*</sup>, MIKE IREY<sup>3</sup>, STEPHEN M GARNSEY<sup>1</sup> and SIDDARAME GOWDA<sup>1,\*</sup>

<sup>1</sup>University of Florida, IFAS, Citrus Research and Education Center, 700 Experiment Station Road, Lake Alfred, FL 33850, USA

<sup>2</sup>Department of Plant Sciences, The University of Tennessee, 252 Ellington Plant Sciences, 2431 Joe Johnson Dr., Knoxville, TN 37996, USA

<sup>3</sup>US Sugar Corporation, Clewiston, FL 33440, USA

\*Corresponding author (Fax, +1-863-9564631; Emails, gowda@ufl.edu, mnrbhav@yahoo.com)

Citrus Huanglongbing (HLB) also known as citrus greening is one of the most devastating diseases of citrus worldwide. The disease is caused by *Candidatus Liberibacter* bacterium, vectored by the psyllid *Diaphorina citri* Kuwayama and *Trioza erytreae* Del Guercio. Citrus plants infected by the HLB bacterium may not show visible symptoms sometimes for years following infection. The aim of this study was to develop effective gene-specific primer pairs for polymerase chain reaction based method for quick screening of HLB disease. Thirty-two different gene-specific primer pairs, across the *Ca. Liberibacter asiaticus* genome, were successfully developed. The possibility of these primer pairs for cross-genome amplification across '*Ca. Liberibacter africanus*' and '*Ca. Liberibacter americanus*' were tested. The applicability of these primer pairs for detection and differentiation of *Ca. Liberibacter* spp. is discussed.

[Nageswara-Rao M, Irely M, Garnsey SM and Gowda S 2013 Candidate gene makers for *Candidatus Liberibacter asiaticus* for detecting citrus greening disease. *J. Biosci.* **38** 1–9] DOI 10.1007/s12038-013-9315-x

Citrus greening (Huanglongbing, HLB) is the most destructive disease of citrus, debilitating the productive capacity of citrus trees wherever it is present worldwide (Halbert and Manjunath 2004; Bové 2006). HLB has been shown to be present internationally, from its first description in the early 1900s in China to its appearance in Florida, USA, as recently as 2005 (Bové 2006). It is estimated that nearly 100 million trees in 40 countries are affected by this disease especially in Asian and Southeast Asian countries like India, China, Indonesia, Philippines, as well as the Arabian Peninsula, and Africa (Bové 1986; Halbert and Manjunath 2004; Gottwald *et al.* 2007) and more recently in Brazil and USA. The fast progression of the disease, in the orchards where the bacterial inoculum or the psyllids have not been efficiently managed, may lead to the infected tree decline in 3 to 13 years (Gottwald *et al.* 1989, 2007). In March 2004, HLB was recognized for the first time in the Americas in São Paulo State, Brazil, that resulted in removal of nearly 3 million HLB-affected sweet orange trees (NRC 2010).

Florida being the second largest citrus producer in the world, its \$9.3 billion annual economic benefit of the citrus industry would be lost or significantly diminished due to severe spread of HLB disease statewide (NRC 2010).

The etiologic causal agent of HLB is a fastidious, phloem-limited, gram-negative bacterium (Garnier *et al.* 1984), restricted to phloem sieve tubes of its natural host, citrus. Despite several attempts, the HLB-causing bacterium has not been obtained in pure culture (Das 2004; Li *et al.* 2006). Based on the nucleotide sequence of its 16S ribosomal RNA gene (rDNA), the HLB pathogen has been characterized as a new genus belonging to the alpha subdivision of the proteobacteria, with the *Candidatus* status '*Candidatus Liberibacter*' (Jagoueix *et al.* 1994). Three *Candidatus* spp. of the pathogen are currently known with most widespread being the Asian species '*Ca. Liberibacter asiaticus*' found in all HLB-affected countries except Africa. The African species *Ca. Liberibacter africanus* and the American species *Ca. L. americanus* are restricted to Africa and Brazil,

**Keywords.** Citrus greening; *Candidatus Liberibacter*; cross-amplification; DNA markers; polymerase chain reaction

**Table 1.** Nucleotide sequence of forward and reverse primers used for the detection of *Candidatus* Liberibacter species

Serial No.	Primer name	GenBank Entry	Gene target	Primer sequence (5'-3')	Amplicon size
1	HLB135 HLB136	EF164805.1	tRNA methyl RNA protein chain elongation factor EF-Tu (tuFB)	F: TACTAAATACGACTCACTATAGGGCTTGTGTGCAGGGCAAGATGTGT R: CCAGATGCAACGCCAACTTCACCTT	1 kb
2	HLB141 HLB142	EF164805.1	tRNA methyl RNA protein chain elongation factor EF-Tu (tuFB)	F: AATCCTTGTGTGAGGGTTCAAATCCC R: TACTAAATACGACTCACTATAGGGACTATATAACC	2.5 kb
3	HLB157 HLB158	JF811345.1	Outer membrane protein	F: CACCGTAGAAGGGCATAATTGAT R: CATGCGATTACCTATACGAAAACC	1.3 kb
4	HLB159 HLB160	FJ489642.1	Outer membrane protein	F: CACCATGATAGTTCCTATTACTG R: AAACCTGCTGAATTTTATCACCCCTC	900 bp
5	HLB163 HLB164	AB473570.1	RNA polymerase beta subunit	F: CGCTTCATGTAGAAAGTTGTG R: CCTACAGGTGGCTGACTCAT	500 bp
6	HLB175 HLB176	CP001677.4	ATP-binding protein; Prophase-related gene cluster-1	F: CCTTAATACGACTCACTATAGCGCGTATGCAATACGAGCGGCA R: CGAGACCCGGGAATACACAAAATACAGCTAAAAATCACTG	1.5 kb
7	HLB177 HLB179	CP001677.4	Prophase-related gene cluster-1, elongation factor EF-Tu	F: CTGATATTGTGAACCATGGGAACATCCTCAC R: GATAGACCAACAGATCAGCGACAAACCCGACGGG	1.5 kb
8	HLB178 HLB179	CP001677.4	Prophase-related gene cluster-1, elongation factor EF-Tu	F: GTGAGGATGTTCCCATGGTTCACAAATACAG R: GATAGACCAACAGATCAGCGACAAACCCGACGGG	825 bp
9	CL420	CLIBASIA_03420	Integral membrane protein MvIN	F: GAGCTTAATTAATGAAGATTATTCGAAACTTTTCTTACGGTATG R: TTATTTTCGAATCATTTGTTGCAATGGGGATAGAAAATC	1.5 kb
10	CL30	CLIBASIA_03030	Zinc uptake ABC transporter, permease protein	F: GAGCTTAATTAATGAAGATTATTCGAAACTTTTCTTACGGTATG R: TTATTTTATAGAAAAGCTTAATAATAAATAAGAAATAAG	723 bp
11	CL025	CLIBASIA_03025	Putative high-affinity zinc uptake system ATP-binding	F: GAGCTTAATTAATGAAGATTATTCGAAACTTTTCTTACGGCTTG R: TTAGTGATCATGTTGTGATTATGAATAGCTAATATTTCTG	780 bp
12	CL20	CLIBASIA_03020	Zinc uptake ABC transporter	F: GAGCTTAATTAATGAAGATTATTCGAAACTTTTCTTACGGTATG R: CTAAGAACACAGTTCTTAGCTATTGAATTAGACATAC	885 bp
13	CL555	CLIBASIA_01555	Hemolysin protein	F: GAGCTTAATTAATGAAGATTATTCGAAACTTTTCTTACGGTATG R: TTAAATTTTAAATGAACAAGACGTGGAATACGCACACGAC	963 bp
14	CL965	CLIBASIA_02965	ABC transporter, membrane-spanning protein	F: GAGCTTAATTAATGAAGATTATTCGAAACTTTTCTTACGGTATG R: TCAATCGTACTTCTCTGATATTTTCTCATGATATACATAG	1.5 kb
15	CL096	CLIBASIA_02960	Phosphate ABC transporter, permease protein PstA	F: GAGCTTAATTAATGAAGATTATTCGAAACTTTTCTTACGGTATG R: TTACCAACCGCTTTTAAAGCGGATTCCTCAACCAATAACAIAG	1.3 kb
16	CL250	CLIBASIA_02250	Extracellular solute-binding protein	F: GAGCTTAATTAATGAAGATTATTCGAAACTTTTCTTACGGTATG R: CTATCGCTTGTATTTGGTCAATATCTCGTTGGCAGTAATA	600 bp
17	CL335	CLIBASIA_01355	Type I secretion membrane fusion protein, HlyD	F: GAGCTTAATTAATGAAGATTATTCGAAACTTTTCTTACGGTATG R: TTACCTTCAAATATTTAGGCCAGTTATCGGTAATG	1.3 kb
18	CL55	CLIBASIA_03055	Component of type IV pilus	F: GAGCTTAATTAATGAAGATTATTCGAAACTTTTCTTACGGTATG R: CTACAAATGACGACTCTGTTTCTCCAGAGATTGAAAGACTTC	1.4 kb
19	CL80	CLIBASIA_03080	Peptidase A24A prepilin type IV	F: GAGCTTAATTAATGAAGATTATTCGAAACTTTTCTTACGGTATG R: TTACGCAGAAAGGCCCACTTAAAGCAACTTTAAACCAAGTATG	530 bp

Table 1 (continued)

Serial No.	Primer name	GenBank Entry	Gene target	Primer sequence (5'-3')	Amplicon size
20	CL165	CLIBASIA_05165	Hypothetical protein	F: GACCTTAATTAATGATCAAAAAGCAACAAGCATTCCATG R: TTATTTGTCAGTGTCTTTTCTGTTTTGTCTG	480 bp
21	CL195	CLIBASIA_05195	Hypothetical protein	F: GAGCTTAATTAATGAACGCTAGCCATATTAATACCCGTTTG R: TTATCCTCTAGCCAAAAGGATCTTTCCCG	615 bp
22	CL200	CLIBASIA_05200	Hypothetical protein	F: GAGCTTAATTAATGGCAAAAAGATTCGCTCTTTTATTAATTG R: CTACCAGAACGGATATCATCTGCTGTAAT	564 bp
23	CL605	CLIBASIA_05605	Hypothetical protein	F: GAGCTTAATTAATGAGTTTTCTTAGATTCGAGTGTAAAATTCGGTTC R: TCATCGTACATGACCCCTGATAAAAAGACTCG	360 bp
24	CL620	CLIBASIA_05620	Hypothetical protein	F: GAGCTTAATTAATGTACGCTCACAAATACACAAAAAGAAATTG R: TTAGTCAATGTTAGTTAACTTCGGAAGCTCAAG	500 bp
25	CL635	CLIBASIA_05635	Hypothetical protein	F: GAGCTTAATTAATGGGACAATTAAGCAATATTACCTCGAAGAG R: TCAAACTGACTGGGTAAATACTTCTTTATG	420 bp
26	CL665	CLIBASIA_05665	Hypothetical protein	F: GAGCTTAATTAATGGGAAAGAAAAGTTTTAACACCTGAAGAAAAGG R: TTAACCTCTCTTATAAATAGTTCTTCTATGG	417 bp
27	CL875	CLIBASIA_02875	Putative flagellar motor switch protein	F: GAGCTTAATTAATGATCAAAAAGTCTCAAAAATTTCTTAC R: TTAGTCAATGTTAAAATTTCTATTACCCGCACTCCAAAAG	430 bp
28	CL910	CLIBASIA_02910	Flagellar MS-ring protein	F: GAGCTTAATTAATGGCTATTTTAGATCAATTTGTCAGTTTTTC R: CTAGCTAATAATGTTGAGTATATCGATCTTCTATTTT	1.7 kp
29	HLB180 HLB181	CP001677.4	Transcriptional regulator	F: GCCGTCGGTGTGTCGCTGATCTGTTGGTCTATC R: AACGGTTAGTTCACGCTAGAGGATATACTAAAAACGG	500 bp
30	CL345	CLIBASIA_01345	Serralysin	F: GAGCTTAATTAATGCATAATATAAAAACCGTTTATACAACTG R: TCAGGAAAAATCATGATTTATATCATGCATAGAAATATAAATC	2.0 kb
31	OMP1	CLIBASIA_00995	Outer membrane protein	F: TGCAGGCTCGGTACCCACAAACTAA R: TGCATCATAGCCACCTGTCACGTG	475 bp
32	OMP2	CLIBASIA_02425	Outer membrane protein	F: ACTCGCCTCTTCTGTTCTGCTC R: AATCCTCCGACTTCAACACCGGACA	474 bp

respectively (Teixeira *et al.* 2005). *Ca. Liberibacter asiaticus*, a heat-tolerant species, and '*Ca. Liberibacter africanus*', a heat-sensitive species, are vectored by both *Diaphorina citri* Kuwayama and *Trioza erytreae* Del Guercio, while '*Ca. Liberibacter americanus*', a heat-tolerant species, is vectored only by *D. citri* Kuwayama (McClellan and Oberholzer 1965; Capoor *et al.* 1967; Bové 2006).


The typical symptoms of HLB disease are blotchy mottling of leaves (NRC 2010) often resembling zinc deficiency symptoms which leads to the typical appearance of yellowing of shoots in the tree. The infected trees show the open growth, stunning twig dieback appearance and severe fruit drop (da Graca and Korsten 2004). Additionally, symptoms of mottling and chlorosis of leaves, and small misshapen fruits with aborted seeds are characteristics of HLB-infected trees. The causal agent of HLB in Florida is *Ca. Liberibacter asiaticus* while both *Ca. Liberibacter asiaticus* and *Ca. Liberibacter americanus* isolates of HLB disease have been detected in Brazil (Bové 2006; NRC 2010). In some instances, more than one species of *Ca. Liberibacter* have been observed in plants as mixed infections (Bové 2006). Primers based on 16S RNA gene have been developed to differentiate between species of *Ca. Liberibacter*. However, availability of additional marker primer pairs to other gene loci would help ascertain the infection by *Ca. Liberibacter* spp. In this study, efforts have been made to develop various candidate gene makers specific to *Ca. Liberibacter asiaticus* for early detection of HLB disease. The effectiveness of the primer pairs developed were also tested for cross-species amplification, if any, against the other two HLB-causing species *Ca. Liberibacter africanus* and *Ca. Liberibacter americanus*.

Total nucleic acids from HLB-infected citrus was extracted from 100 mg of tissue (leaf and/or inner bark tissue from healthy as well as HLB-infected plants, Tatineni *et al.* 2008) and also from HLB-infected psyllids. Total nucleic acid samples of *Ca. Liberibacter africanus* were kindly provided by Prof G Pietersen (University of Pictoria, South Africa) and total nucleic acid samples from *Ca. Liberibacter asiaticus* as well as *Ca. Liberibacter americanus* from Brazil were kindly provided by Mr Luis Matos (Dominican Institute of Agriculture and Forestry Research, Dominican Republic).

Conventional PCR with SpeedSTAR HS DNA polymerase (Takara Bio, WI) was used to examine the presence of the HLB bacterium in citrus as well as psyllid samples. The PCR reaction (20  $\mu$ L volume) consisted of 1  $\mu$ L of template DNA, 0.2  $\mu$ M each oligonucleotide (gene-specific forward and reverse primers; table 1), 0.25 mM dNTPs, 1 $\times$  buffer (FBII; Takara Bio, USA), and 0.125  $\mu$ L (5 U/ $\mu$ L) of SpeedSTAR HS DNA polymerase. Amplification was carried out using the following protocol (primer pairs 1–19, table 1): 94°C for 2 min; followed by 10 cycles at 94°C for

10 s, 54°C for 10 s, and 72°C for 90 s; followed by 25 cycles at 94°C for 10 s, 58°C for 10 s, and 72°C for 90 s; followed by final extension at 72°C for 5 min. For primer pairs 20–32 (table 1), the PCR cycle was modified slightly with the extension being carried out for 60 s instead of 90 s. PCR reaction (15  $\mu$ L) was analysed through 1.5% agarose gel in 1 $\times$  Tris-acetate-EDTA buffer (40 mM Tris, 20 mM acetic acid, 1 mM EDTA, pH 8.5) and PCR amplicons were visualized using ethidium bromide. The PCR products were eluted from the gel and purified using GeneClean® Kit (MP Biomedicals, USA) following the manufacturer's instructions and sent to core sequencing facility, Interdisciplinary Center for Biotechnology Research (ICBR), University of Florida, Gainesville, FL, for gene sequence confirmation. For confirmation of *Ca. Liberibacter asiaticus*, *Ca. Liberibacter africanus* as well as *Ca. Liberibacter americanus* bacterium, the total nucleic acids samples were also screened against HLB gene-specific 16S rDNA primer pairs developed by Teixeira *et al.* (2005). The HLB-free, healthy citrus samples were used as a negative control in PCR amplifications. All the experiments were performed in triplicates and the results obtained were reconfirmed.

Availability of increasing number of nucleotide (genome) sequences in the public domain provides a fast and efficient approach to develop gene-specific PCR molecular markers (Chen *et al.* 2008; Soneji *et al.* 2010). Based on *Ca. Liberibacter asiaticus* whole genome sequence information (Duan *et al.* 2009; GenBank accession number NC\_012985; NZ\_ABQW01000001 to NZ\_ABQW01000034), 50 gene-specific primer-pairs for HLB pathogen were developed using the online free primer development software, PrimerQuest<sup>SM</sup>-Integrated DNA Technologies, USA. Of the 50 primer pairs, 32 primer pairs (table 1) amplified specific amplicons from the total nucleic acids isolated from *Ca. Liberibacter asiaticus*-infected citrus plants as well as infected psyllids (figure 1). No PCR products were amplified from the healthy citrus plants as well as psyllids. The GenBank accession number, primer name, loci studied, and the expected PCR product size are mentioned in table 1. The study also successfully standardized a common PCR thermocycler program for all the primer pairs developed. The PCR amplicon sizes observed on the agarose gel electrophoresis were compatible with the *Ca. Liberibacter asiaticus*-infected citrus plants as well as infected psyllids. The PCR amplification of all the primer pairs were also screened against the *Ca. Liberibacter asiaticus* infected samples obtained from Brazil and the amplicon sizes were found to be compatible with *Ca. Liberibacter asiaticus* infected samples from Florida (figure 2). The amplified gene fragments from agarose gel were gel-eluted (for both the American as well as Brazilian *Ca. Liberibacter* samples), gene-cleaned (GeneClean® Kit; MP Biomedicals, USA), sequenced and the sequence information were reconfirmed with respect to their GenBank accession numbers (table 1). The nucleotide




**Figure 1.** PCR amplification of *Candidatus Liberibacter asiaticus* specific sequences. (A, B, C, D) In HLB infected citrus plants. (E) In HLB infected psyllids. (The numbers in the lane corresponds to primer pairs in table 1); M: 1 kb DNA ladder; W: Water.

sequences were also aligned using ClustalW2 – multiple sequence alignment (<http://www.ebi.ac.uk/Tools/msa/clustalw2/>) and 98.5-100% sequence homology was observed between the American and Brazilian *Ca. Liberibacter* samples.

Total nucleic acids from citrus infected with *Ca. Liberibacter asiaticus* and *Ca. Liberibacter americanus* obtained from Brazil and the *Ca. Liberibacter africanus* from South Africa were reconfirmed by PCR experiments using primer pairs HLB75<sup>(+)</sup>/HLB177T<sup>(-)</sup> (*Ca. Liberibacter asiaticus*), HLB77<sup>(+)</sup>/HLB177T<sup>(-)</sup> (*Ca. Liberibacter africanus*) and HLB73<sup>(+)</sup>/HLB166<sup>(-)</sup> (*Ca. Liberibacter americanus*) (table 2). The PCR amplification from these primer pairs, developed from the *Ca. Liberibacter* 16S

rRNA genes, allowed us for a direct identification of all the three species.

In order to understand the cross-compatibility of the primer pairs developed in this study, across the *Ca. Liberibacter* spp., viz *Ca. Liberibacter asiaticus*, *Ca. Liberibacter africanus*, and *Ca. Liberibacter americanus*, cross-species amplification studies were carried out. Of the 32 primer pairs, only three primer pairs (1, 7, and 10; table 1; figure 3) showed cross-genome amplification in all the three *Ca. Liberibacter* spp. and three primer pairs (3, 5 and 9; table 1) showed cross-genome amplification between *Ca. Liberibacter asiaticus* and *Ca. Liberibacter africanus* and no PCR product amplicons were observed for the *Ca.*



**Figure 2.** PCR amplification of *Candidatus Liberibacter asiaticus* samples. (A) HLB samples from Florida. (B) HLB samples from Brazil. The numbers in the lane corresponds to primer pairs in table 1. W: water. M: 1 kb DNA ladder.


**Table 2.** Nucleotide sequence of forward and reverse primers used for the amplification of *Candidatus Liberibacter* 16S rDNA (Teixeira et al. 2005)

Serial No.	Primer name	Primer sequence (5'—3')	PCR product size	HLB bacterium
1	HLB73 HLB116	F: AGTCGAGCGAGTACGCAAGTACT R: CAACTTAATGATGGCAAATATAG	1027 bp	<i>Ca. Liberibacter americanus</i>
2	HLB75 HLB177T	F: CGCGTATGCAATACGAGCGGCA R: GCCTCGCGACTTCGCAACCCAT	1027 bp	<i>Ca. Liberibacter asiaticus</i>
3	HLB77 HLB177T	F: GCGCGTATTTTATACGAGCGGCA R: GCCTCGCGACTTCGCAACCCAT	1027 bp	<i>Ca. Liberibacter africanus</i>

*Liberibacter americanus*. High rates of transferability of molecular loci across species within a genus have been reported (Peakall et al. 1998; Scott et al. 2001; Clauss et al. 2002; Dirlewanger et al. 2002; Gaitán-Solís et al. 2002; Jones et al. 2002; Williamson et al. 2002; Thiel et al. 2003; Eujayl et al. 2004; Saha et al. 2004). It has also been observed that the molecular loci may exhibit decreasing variation as the evolutionary distance increases from the species used to develop the primer pairs (Sun and Kirkpatrick 1996; White and Powell 1997; Peakall et al. 1998; Roa et al. 2000; Thiel et al. 2003). For some of the primer pairs, 1, 3 and 10, more than one amplicon was observed. Amplification of more than one amplicon using EST-SSR markers has been reported in previous studies and was attributed to the possible amplification of both orthologous and paralogous copies of the target region (Varshney et al. 2005; Sim et al. 2009), or homoloci from different genomes (Gupta et al. 2003; Saha et al. 2004). The cross-genome amplicons for the primer pairs, 1, 3, 5, 7, 9 and 10 (table 1), for all the three *Ca. Liberibacter* species, were gel-eluted, gene-cleaned (GeneClean® Kit; MP Biomedicals, USA), sequenced and the sequence information were blasted against the GenBank database. Except the *Ca. Liberibacter asiaticus* for which the primers were originally developed, no sequence homology was observed in *Ca. Liberibacter africanus*, and *Ca. Liberibacter americanus* for the primer pairs. Studies

have also observed loss of sequence homology when markers developed from one species were screened on related species (Asp et al. 2007; Sim et al. 2009; Yu et al. 2011). While cross-amplification of different types of molecular loci even across genera is possible, more attention should be given to the nature of the amplified fragments before inferring synteny or orthology in genetic diversity studies (table 3). As new sequencing technologies are emerging, longer nucleotide sequence reads at much lower costs than what is available right now are anticipated, making genome sequencing the method of choice for cross-genome/genera studies (Soneji et al. 2010; Zeid et al. 2010).

In this study, the effectiveness of the application of PCR primer pairs specific to *Ca. Liberibacter asiaticus*, causing HLB disease in citrus, was studied. In the absence of any durable resistant citrus cultivars, the management of HLB at least in the immediate future is to eradicate the source of pathogen, the infected citrus trees and the transmitting psyllid vector (Halbert and Manjunath 2004; Pelz-Stelinski et al. 2010). Management of citrus greening disease requires rapid and large scale detection, followed by removing of infected trees, and this practice is hindered due to lack of quick, sensitive and large-scale detection methods. PCR-based methods currently being used are based on the 16S rRNA gene region (Jagoueix et al. 1997; Teixeira et al. 2005; Das et al.

**Figure 3.** Cross-species amplification of *Candidatus Liberibacter* species. M: 1 kb DNA ladder. Gel A (primer 10 from table 1), B (primer 1 from table 1), C (primer 7 from table 1): Lane 1, *Candidatus Liberibacter asiaticus* from Florida; Lane 2, *Candidatus Liberibacter asiaticus* from Brazil; Lane 3, *Candidatus Liberibacter africanus*; Lane 4, *Candidatus Liberibacter americanus*.

**Table 3.** A few representative studies with no sequencing information presented in the cross-species/genome studies

Serial No.	Species name	Cross-species/genome studies
1	<i>Aphis fabae</i>	Gauffre and Coeur d'acier (2006)
2	<i>Asparagus racemosus</i>	Ginwall <i>et al.</i> (2011)
3	<i>Blattella germanica</i>	Booth <i>et al.</i> (2007)
4	Citrus	Dong <i>et al.</i> (2006)
5	<i>Jatropha curcas</i>	Mittal and Dubey (2010)
6	Rice, Maize, Soybean, Wheat	Guo <i>et al.</i> (2003); Bandopadhyay <i>et al.</i> (2004)
7	<i>Microtus arvalis</i>	Walser and Heckel (2008)
8	<i>Liriodendron chinense</i>	Yao <i>et al.</i> (2008)
9	<i>Locusta migratoria</i>	Zhang <i>et al.</i> (2003)
10	<i>Pogona vitticeps</i>	Quinn <i>et al.</i> (2010)
11	<i>Rhizobia</i>	Ya-mei <i>et al.</i> (2008)
12	<i>Ruditapes variegata</i>	An <i>et al.</i> (2009)
13	<i>Casuarina</i> spp.	Yasodha <i>et al.</i> (2005)
14	<i>Suta</i> spp.	Stapley <i>et al.</i> (2005)
15	<i>Elaeocarpus</i> spp.	Jones <i>et al.</i> (2002)

2007; Das 2009). Because these conserved regions are homologous to sequences of the host and/or citrus associated endophytes, reliable and specific detection and diagnosis of HLB is of concern (Lin *et al.* 2008). Thus, the candidate gene markers developed in this study, across the wide genome of *Ca. Liberibacter asiaticus*, will essentially benefit the citrus industry by providing simple, sensitive and rapid detection method for large-scale detection/management of HLB under field, nursery, bud-wood certification and epidemiological conditions. It will also be useful for the identification of *Ca. Liberibacter asiaticus* where more than one species of HLB pathogen is present.

### Acknowledgements

This research was supported by the Florida Citrus Production Research Advisory Council, Florida, USA. The authors also thank Dr. William O Dawson for permission to use his laboratory and greenhouse facilities.

### References

- An SH, Kim SM and Park JY 2009 Isolation and characterization of microsatellite markers for the clam *Ruditapes philippinarum* and cross-species amplification with the clam *Ruditapes variegata*. *Conserv. Genet.* **10** 1821–1823
- Asp T, Frei UK, Didion T, Nielsen KK and Lubberstedt T 2007 Frequency, type, and distribution of EST-SSRs from three genotypes of *Lolium perenne*, and their conservation across orthologous sequences of *Festuca arundinacea*, *Brachypodium distachyon*, and *Oryza sativa*. *BMC Plant Biol.* **7** 36
- Bandopadhyay R, Sharma S, Rustgi S, Singh R, Kumar A, Balyan HS and Gupta PK 2004 DNA polymorphism among 18 species of *Triticum-Aegilops* complex using wheat EST-SSRs. *Plant Sci.* **166** 349–356
- Booth W, Bogdanowicz SM, Prodöhl PA, Harrison RG, Schal C and Vargo EL 2007 Identification and characterization of 10 polymorphic microsatellite loci in the German cockroach, *Blattella germanica*. *Mol. Ecol. Notes.* **7** 648–650
- Bové JM 1986 Greening in Arab Peninsula: Towards new techniques for its detection and control. *FAO Plant Prot. Bull.* **34** 7-14
- Bové JM 2006 Huanglongbing: A destructive, newly-emerging, century-old disease of citrus. *J. Plant Pathol.* **88** 7–37
- Capoor SP, Rao DG and Viswanath SM 1967 *Diaphorina citri* Kuway., a vector of the Huanglongbing disease of citrus in India. *Indian J. Agric. Sci.* **37** 572–576
- Chen C, Bowman KD, Choi YA, Dang PM, Nageswara-Rao M, Huang S, Soneji JR, McCollum TG and Gmitter FG Jr 2008 EST-SSR genetic maps for *Citrus sinensis* and *Poncirus trifoliata*. *Tree Genet. Genomes* **4** 1–10
- Clauss MJ, Cobban H and Mitchell-Olds T 2002 Cross-species microsatellite markers for elucidating population genetic structure in *Arabidopsis* and *Arabis* (Brassicaceae). *Mol. Ecol.* **11** 591–601
- Da Graca JV and Korsten L 2004. Citrus Huanglongbing: Review, present status and future strategies; in *Diseases of fruits and vegetables* vol I (ed) SAMH Naqui (Dordrecht, The Netherlands: Kluwer Academic Publishers) pp 229–245
- Das AK 2004 Rapid detection of *Candidatus Liberibacter asiaticus*, the bacterium associated with citrus Huanglongbing (Greening) disease using PCR. *Curr. Sci.* **87** 1183–1185
- Das AK 2009 Molecular identification and characterization of citrus greening bacterium, *Candidatus Liberibacter asiaticus* associated with decline of Nagpur mandarin orange in Vidarbha region, Maharashtra. *Curr. Sci.* **96** 890–892
- Das AK, Rao CN and Singh S 2007 Presence of citrus greening (Huanglongbing) disease and its psyllid vector in the North-Eastern region of India confirmed by PCR technique. *Curr. Sci.* **92** 1759–1763
- Dirlwanger E, Cosson P, Tavaud M, Aranzana MJ, Poizat C, Zanetto A, Arús P and Laigret F 2002 Development of microsatellite markers in peach [*Prunus persica* (L.) Batsch] and their use in genetic diversity analysis in peach and sweet cherry (*Prunus avium* L.). *Theor. Appl. Genet.* **105** 127–138
- Dong J, Guang-Yan Z and Qi-Bing H 2006 Analysis of microsatellites in *Citrus unigenes*. *Acta Genetica Sinica* **33** 345–353
- Duan Y, Zhou L, Hall DG, Li W, Doddapaneni H, Lin H, Liu L, Gabriel D, Vahling CM, Williams K, Dickerman A, Sun Y and Gottwald TR 2009 Complete genome sequence of citrus huanglongbing bacterium, *Candidatus Liberibacter asiaticus*-obtained through metagenomics. *Mol. Plant-Microbe Interact.* **22** 1011–1020
- Eujayl I, Sledge MK, Wang L, May GD, Chekhovskiy K, Zwonitzer JC and Mian MAR 2004 *Medicago truncatula* EST-SSRs reveal cross-species genetic markers for *Medicago* spp. *Theor. Appl. Genet.* **108** 414–422

- Gaitán-Solis E, Duque MC, Edwards KJ and Tohme J 2002 Microsatellite repeats in common bean (*Phaseolus vulgaris*): Isolation, characterization, and cross-species amplification in *Phaseolus* spp. *Crop Sci.* **42** 2128–2136
- Garnier M, Danel N and Bové JM 1984 The greening organism is a gram-negative bacterium; in *Proc. 9th Conf. Intl. Org. Citrus Virologists* (eds) SM Garnsey, LW Timmer and JA Dodds (Riverside, CA, USA: University of California) pp 115–124
- Gauffre B and Coeur d'acier A 2006 New polymorphic microsatellite loci, cross-species amplification and PCR multiplexing in the black aphid, *Aphis fabae* Scopoli. *Mol. Ecol. Notes* **6** 440–442
- Ginwall HS, Mittal N, Maurya SS, Barthwal S and Bhatt P 2011 Genomic DNA isolation and identification of chloroplast microsatellite markers in *Asparagus racemosus* Willd. through cross-amplification. *Indian J. Biotechnol.* **10** 33–38
- Gottwald TR, Aubert B and Zhao X-Y 1989 Preliminary analysis of citrus greening (Huanglongbing) epidemics in the People's Republic of China and French Reunion Island. *Phytopathol.* **79** 687–693
- Gottwald TR, da Graça JV and Bassanezi RB 2007 Citrus Huanglongbing: The pathogen and its impact. *Plant Health Progress* doi:10.1094/PHP-2007-0906-01-RV
- Guo L, Tang J, Li H and Jia J 2003 Analysis of microsatellites in major crops assessed by computational and experimental approaches. *Mol. Breed.* **12** 245–261
- Gupta PK, Rustgi S, Sharma S, Singh R, Kumar N and Balyan HS 2003 Transferable EST-SSR markers for the study of polymorphism and genetic diversity in bread wheat. *Mol. Genet. Genom.* **270** 315–323
- Halbert SE and Manjunath KL 2004 Asian citrus psyllids (Sternorrhyncha: Psyllidae) and greening disease of citrus: A literature review and assessment of risk in Florida. *Florida Entomol.* **87** 330–352
- Jagoueix S, Bové JM and Garnier M 1994 The phloem-limited bacterium of greening disease of citrus is a member of the a-subdivision of the Proteobacteria. *Int. J. Syst. Bacteriol.* **44** 379–386
- Jagoueix S, Bové JM and Garnier M 1997 Comparison of the 16S/23S ribosomal intergenic regions of “*Candidatus Liberibacter asiaticum*” and “*Candidatus Liberibacter africanum*” the two species associated with citrus Huanglongbing (greening) disease. *Int. J. Syst. Bacteriol.* **47** 224–227
- Jones RC, McNally J and Rossetto M 2002 Isolation of microsatellite loci from a rainforest tree *Elaeocarpus grandis* (Elaeocarpaceae), and amplification across closely related taxa. *Mol. Ecol. Notes* **2** 179–181
- Li W, Hartung JS and Levy LE 2006 Quantitative real time PCR for detection and identification of ‘*Candidatus Liberibacter species*’ associated with citrus Huanglongbing. *J. Microbiol. Methods* **66** 104–115
- Lin H, Doddapaneni H, Bai X, Yao J, Zhao X and Civerolo EL 2008 Acquisition of uncharacterized sequences from *Candidatus Liberibacter*, an unculturable bacteria, using an improved genomic walking. *Mol. Cell. Probes* **22** 30–37
- McClellan APD and Oberholzer PCJ 1965 Citrus psylla, a vector of the greening disease of sweet orange. *S. Afr. J. Agric. Sci.* **8** 297–298
- Mittal N and Dubey AK 2010 A novel set of highly polymorphic chloroplast, microsatellite and ISSR markers for the biofuel crop *Jatropha curcas*. *EurAsia J. BioSci.* **4** 119–131
- National Research Council 2010 Strategic planning for the Florida citrus industry: Addressing citrus greening <http://www.nap.edu/catalog/12880.html>
- Peakall R, Gilmore S, Keys W, Morgante M and Rafalski A 1998 Cross-species amplification of soybean (*Glycine max*) simple sequence repeats (SSRs) within the genus and other legume genera, implications for the transferability of SSRs in plants. *Mol. Biol. Evol.* **15** 1275–1287
- Pelz-Stelinski KS, Bransky RH and Rogers ME 2010 Transmission of *Candidatus Liberibacter asiaticus* by the Asian citrus psyllid, *Diaphorina citri*. *J. Economic Entomol.* **103** 1531–1541
- Quinn AE, Ezaz T, Sarre SD, Graves JAM and Georges A 2010 Extension, single-locus conversion and physical mapping of sex chromosome sequences identify the Z microchromosome and pseudo-autosomal region in a dragon lizard, *Pogona vitticeps*. *Heridity* **104** 410–417
- Roa AC, Chavarriaga-Aguirre P, Duque MC, Maya M, Bonierbale MW, Iglesias C and Tohme J 2000 Cross-species amplification of cassava (*Manihot esculenta*) Euphorbiaceae microsatellites: allelic polymorphism and degree of relationships. *Am. J. Bot.* **87** 1647–1655
- Saha MC, Mian MAR, Eujayl I, Zwonitzer JC, Wang L and May GD 2004 Tall fescue EST-SSR markers with transferability across several grass species. *Theor. Appl. Genet.* **109** 783–791
- Scott IAW, Hayes CMJ, Koegh S and Morrison SF 2001 Isolation and characterization of novel microsatellite markers from the Australian water skin *Eulamprus kosciuskoi* and cross-species amplification in other members of the species-group. *Mol. Ecol. Notes* **1** 28–30
- Sim SC, Yu JK, Jo Y, Sorrells ME and Jung G 2009 Transferability of cereal EST-SSR markers to ryegrass. *Genome* **52** 431–437
- Soneji JR, Nageswara-Rao M, Sudarshana P, Panigrahi J and Kole C 2010 Current status on on-going genome initiatives; in *Principles and practices of plant genomics* vol 3 *Structural and functional genomics* (eds) C Kole and AG Abbott (Enfield, New Hampshire, USA: Science Publishers Inc. & New York, USA: CRC Press, Taylor and Francis Group) pp 305–353
- Stapley J, Hayes CM, Webb JK and Koegh JS 2005 Novel microsatellite loci identified from the Australian eastern small-eyed snake (Elapidae: *Rhinocephalus nigrescens*) and cross species amplification in the related genus *Suta*. *Mol. Ecol. Notes* **5** 54–56
- Sun HS and Kirkpatrick BW 1996 Exploiting dinucleotide microsatellites conserved among mammalian species. *Mamm. Gen.* **7** 128–132
- Tatineni S, Sagaram US, Gowda S, Robertson C, Dawson WO, Iwanami T and Wang N 2008 In planta distribution of ‘*Candidatus Liberibacter asiaticus*’ as revealed by polymerase chain reaction (PCR) and real-time PCR. *Phytopathology* **98** 592–599
- Teixeira DC, Danet JL, Eveillard S, Martins EC, Jesus Jr WC, Yamamoto PT, Lopes SA, Bassanezi EB, Ayres AJ, Saillard C and Bové JM 2005 Citrus Huanglongbing in São Paulo, Brazil: PCR detection of the ‘*Candidatus Liberibacter*’ species associated with the disease. *Mol. Cell. Probes* **19** 173–179



- Thiel T, Michalek W, Varshney R and Graner A 2003 Exploiting EST databases for the development and characterization of gene-derived SSR-markers in barley (*Hordeum vulgare* L.). *Theor. Appl. Genet.* **106** 411–422
- Varshney RK, Graner A and Sorrells ME 2005 Genic microsatellite markers in plants: features and applications. *Trends Biotechnol.* **23** 48–55
- Walser B and Heckel W 2008 Microsatellite markers for the common vole (*Microtus arvalis*) and their cross-species utility. *Conserv. Genet.* **9** 479–481
- White G and Powell W 1997 Isolation and characterization of microsatellite loci in *Swietenia humilis* (Meliaceae): An endangered tropical hardwood species. *Mol. Ecol.* **6** 851–860
- Williamson JE, Huebinger RM, Sommer JA, Louis Jr EE and Barber RC 2002 Development and cross-species amplification of 18 microsatellite markers in the Sumatran tiger (*Panthers tigris sumatrae*). *Mol. Ecol. Notes* **2** 110–112
- Ya-mei G, Yi-qiang H, Hui T, Dong-mei S, Yan-jie W and Wei-dong W 2008 Analysis of simple sequence repeats in genomes of *Rhizobia*. *Agric. Sci. China* **7** 1189–1195
- Yao X, Zhang J, Ye Q and Huang H 2008 Characterization of 14 novel microsatellite loci in the endangered *Liriodendron chinense* (Magnoliaceae) and cross-species amplification in closely related taxa. *Conserv. Genet.* **9** 483–485
- Yasodha R, Ghosh M, Sumathi R and Gurumurthy K 2005 Cross-species amplification of eucalyptus SSR markers in Casuarinaceae. *Acta. Bot. Croat.* **64** 115–120
- Yu F, Wang B, Feng S, Wang J, Li W and Wu Y 2011 Development, characterization, and cross-species/genera transferability of SSR markers for rubber tree (*Hevea brasiliensis*). *Plant Cell Rep.* **30** 335–344
- Zeid M, Yu JK, Goldowitz I, Dentond ME, Costich DE, Jayasuriya CT, Saha M, Elshire R, Benschel D, Breseghello F, Munkvold J, Varshney RK, Belay G and Sorrells ME 2010 Cross-amplification of EST-derived markers among 16 grass species. *Field Crops Res.* **118** 28–35
- Zhang D, Yan L, Ji Y, Kang LE, Godfrey HM and Huang Z 2003 Isolation, characterization and cross-species amplification of eight microsatellite DNA loci in the migratory locust (*Locusta migratoria*). *Mol. Ecol. Notes* **3** 483–486

MS received 14 August 2012; accepted 12 March 2013

Corresponding editor: MAN MOHAN JOHRI